



Ann. Funct. Anal. 2 (2011), no. 1, 13–18

ANNALS OF FUNCTIONAL ANALYSIS

ISSN: 2008-8752 (electronic)

URL: [www.emis.de/journals/AFA/](http://www.emis.de/journals/AFA/)

## UNIFORM BOUNDEDNESS PRINCIPLES FOR ORDERED TOPOLOGICAL VECTOR SPACES

SHUHUI ZHONG<sup>1\*</sup> AND RONGLU LI<sup>2</sup>

Communicated by J. Esterle

**ABSTRACT.** We obtain uniform boundedness principles for a new class of families of mappings from topological vector spaces to ordered topological vector spaces. The new class of families of mappings includes the family of linear mappings and many other families which consist of nonlinear mappings.

### 1. INTRODUCTION AND PRELIMINARIES

The uniform boundedness principle is well known as one of the foundation stones of functional analysis. Its basic form [3] asserts that a pointwise boundedness family of continuous linear operators from a Banach space  $X$  to a normed space  $Y$  is equicontinuous at each  $x \in X$ , and uniformly bounded on each bounded subset of  $X$ . For its importance there has been a lot of work (see books [1, 16]) on uniform boundedness principles since Banach and Steinhaus gave the above version of uniform boundedness principles in 1927. Especially, today we can find various versions [2], [4]-[11], [13]-[15] of uniform boundedness principles in many different mathematical fields.

In this paper, we will give two uniform boundedness principles for a new class of families of mappings from topological vector spaces to ordered topological vector spaces, and point out that the new class of families of mappings includes the family of linear mappings and many other families which consist of nonlinear mappings. Our results have strongly generalized the above version of uniform

---

*Date:* Received: 4 January 2011; Accepted: 27 January 2011.

\* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 46A30; Secondary 46A08.

*Key words and phrases.* Uniform boundedness principles, UB, order, topological vector spaces.

boundedness principle, but are very different from any of the results in relevant literature.

In the whole paper, all topological vector spaces are Hausdorff and over the real scalar field  $\mathbb{R}$ . For a topological vector space  $X$ , denote by  $\mathcal{N}(X)$  the family of neighborhoods of  $0 \in X$ .

**Definition 1.1.** Let  $X, Y$  be topological vector spaces. A family  $\Gamma$  of mappings from  $X$  to  $Y$  is said to be UB if there exist  $U \in \mathcal{N}(X)$ ,  $\delta, M > 0$  and  $\varphi : (0, \delta) \rightarrow (0, +\infty)$  with  $\lim_{t \rightarrow 0} \varphi(t) = 0$  such that for all  $f \in \Gamma$ :

- (1)  $f(0) = 0$ ;
- (2) For all  $u \in U$  and  $x \in X$ , there exist  $r, s \in [-M, M]$  such that  $f(x - u) = rf(x) + sf(u)$ ;
- (3) For all  $u \in U$  and  $0 < t < \delta$ , there exists a real  $s$  such that  $|s| \leq \varphi(t)$  and  $f(tu) = sf(u)$ .

*Remark 1.2.* The family  $\Gamma$  which consists of linear mappings from  $X$  to  $Y$  is UB with  $U = X$ ,  $M = 1$  and  $\varphi(t) = t$ .

**Example 1.3.** Let  $(X, \|\cdot\|)$  be a normed space. Then the mapping  $\|\cdot\| : X \rightarrow [0, +\infty)$  belongs to the UB family of mappings from  $X$  to  $\mathbb{R}$  with  $U = X$ ,  $M = 1$  and  $\varphi(t) = t$ .

In fact, since  $\|x\| - \|u\| \leq \|x - u\| \leq \|x\| + \|u\|$  for all  $x, u \in X$  we have  $\|x - u\| = \|x\| + s\|u\|$  for some  $|s| \leq 1$ .

**Example 1.4.** Let  $(X, \|\cdot\|)$  be a normed space and  $Y$  a vector space. For a linear mapping  $T$  from  $X$  to  $Y$ , define  $f_T : X \rightarrow Y$  by

$$f_T(x) = \frac{1}{1 + \|x\|} T(x), \quad \forall x \in X.$$

Then the family  $\Gamma = \{f_T : X \rightarrow Y | T : X \rightarrow Y \text{ is linear}\}$  is UB with  $U = \{x \in X : \|x\| < 1\}$ ,  $M = 2$  and  $\varphi(t) = 2t$ .

For each  $f_T \in \Gamma$ ,  $f_T(0) = 0$  is obvious. For  $x \in X$  and  $u \in U$ ,

$$\begin{aligned} f_T(x - u) &= \frac{1}{1 + \|x - u\|} T(x - u) = \frac{1}{1 + \|x - u\|} T(x) - \frac{1}{1 + \|x - u\|} T(u) \\ &= \frac{1 + \|x\|}{1 + \|x - u\|} f_T(x) - \frac{1 + \|u\|}{1 + \|x - u\|} f_T(u) \end{aligned}$$

and

$$\frac{1 + \|x\|}{1 + \|x - u\|} < 2 = M, \quad \frac{1 + \|u\|}{1 + \|x - u\|} < 2 = M$$

since  $2(1 + \|x - u\|) \geq 2 + \|x - u\| + \|x\| - \|u\| \geq 1 + \|x\| + 1 - \|u\| > 1 + \|x\|$  and  $2(1 + \|x - u\|) \geq 2 > 1 + \|u\|$ .

For  $u \in U$  and  $t > 0$ ,

$$f_T(tu) = \frac{1}{1 + \|tu\|} T(tu) = \frac{1}{1 + \|tu\|} tT(u) = \frac{1 + \|u\|}{1 + \|tu\|} t f_T(u)$$

where

$$\left| \frac{1 + \|u\|}{1 + \|tu\|} t \right| < 2|t| = \varphi(t) \text{ with } t > 0.$$

*Remark 1.5.* Each  $f_T \in \Gamma$  in Example 1.4 is nonlinear.

*Remark 1.6.* The family of demi-linear mappings [9] from  $X$  to  $Y$  related to  $\gamma$  and  $U$  is UB with  $U$ ,  $\delta = 1$ ,  $M = 1 + |\gamma(-1)|$ ,  $\varphi = |\gamma|_{|(0,\delta)} : (0, \delta) \rightarrow (0, +\infty)$ .

**Definition 1.7.** [12] Let  $Y$  be a topological vector space and an ordered vector space.  $Y$  is called an ordered topological vector space if the positive cone  $C = \{y \in Y : y \geq 0\}$  is closed in  $Y$ .

For topological vector spaces  $X$  and  $Y$ , a family  $\Gamma \subset Y^X$  is said to be pointwise bounded if for each  $x \in X$  the set  $\{f(x) : f \in \Gamma\}$  is bounded in  $Y$ . If for each bounded subset  $B \subset X$  the set  $\{f(x) : f \in \Gamma, x \in B\}$  is bounded in  $Y$ , then  $\Gamma$  is said to be uniformly bounded on each bounded subset of  $X$ .

And for  $Y$  an ordered topological vector space, a family  $\Gamma \subset Y^X$  is said to be pointwise order bounded if for each  $x \in X$  the set  $\{f(x) : f \in \Gamma\}$  is order bounded in  $Y$ . If for each bounded subset  $B \subset X$  the set  $\{f(x) : f \in \Gamma, x \in B\}$  is order bounded in  $Y$ , then  $\Gamma$  is said to be uniformly order bounded on each bounded subset of  $X$ .

For more notations and terminologies in topological vector spaces or ordered topological vector spaces, we can refer to [16] or [12].

## 2. MAIN RESULTS

**Theorem 2.1.** *Let  $X$  be a topological vector space of second category and  $Y$  an ordered topological vector space with an order unit  $e$ . If the UB family  $\Gamma$  of continuous mappings from  $X$  to  $Y$  with  $U \in \mathcal{N}(X)$ ,  $\delta$ ,  $M > 0$  and  $\varphi : (0, \delta) \rightarrow (0, +\infty)$  satisfying  $\lim_{t \rightarrow 0} \varphi(t) = 0$  is pointwise order bounded, then  $\Gamma$  is uniformly order bounded on each bounded subset of  $X$ .*

*Proof.* Let  $U$  be balanced and closed, and  $x \in X$ . Then there exists  $\theta \in (0, 1)$  such that  $\theta x \in U$  and  $\{f(\theta x) : f \in \Gamma\}$  is order bounded in  $Y$ . Pick  $a, b \in Y$  with

$$\{f(\theta x) : f \in \Gamma\} \subset [a, b] := \{y \in Y : a \leq y \leq b\}.$$

Since  $e$  is an order unit in  $Y$ , there exists  $n \in \mathbb{N}$  for which

$$[a, b] \subset [-ne, ne] = n[-e, e].$$

Thus

$$\{f(\theta x) : f \in \Gamma\} \subset n[-e, e].$$

Since  $\lim_{t \rightarrow 0} \varphi(t) = 0$  there exists  $m \in \mathbb{N}$  such that  $1/m < \delta$  and  $\varphi(1/m) \leq (1/n)$ . But  $f((1/m)\theta x) = s_f f(\theta x)$ , where  $|s_f| \leq \varphi(1/m) \leq (1/n)$  and  $s_f n[-e, e] \subset [-e, e]$  for all  $f \in \Gamma$ . Then

$$\{f((1/m)\theta x) : f \in \Gamma\} = \{s_f f(\theta x) : f \in \Gamma\} \subset \cup_{f \in \Gamma} (s_f n[-e, e]) \subset [-e, e]$$

and  $(1/m)\theta x \in \cap_{f \in \Gamma} f^{-1}([-e, e])$ .

Obviously,  $(1/m)\theta x \in U$ . Let

$$W = U \cap (\cap_{f \in \Gamma} f^{-1}([-e, e])).$$

From the above deducing process, we know  $X = \cup_{n=1}^{\infty} nW$ . By Definition 1.7, the positive cone  $C = \{y \in Y : y \geq 0\}$  is closed in  $Y$ . So  $[-e, e]$  is also closed in  $Y$ . Then  $f^{-1}([-e, e])$  is closed in  $X$  since each  $f \in \Gamma$  is continuous. Thus

$$W = U \cap (\cap_{f \in \Gamma} f^{-1}([-e, e]))$$

is closed in  $X$ . But  $X$  is a topological vector space of second category. There exists  $n_0 \in \mathbb{N}$  with  $n_0W$  having nonempty interior. So  $W$  has nonempty interior and  $W - W = \{x - z : x, z \in W\} \in \mathcal{N}(X)$ .

Let  $x, z \in W$  and  $f \in \Gamma$ . Then  $f(x), f(z) \in [-e, e]$  and

$$f(x - z) = r_f f(x) + s_f f(z)$$

where  $|r_f| \leq M$  and  $|s_f| \leq M$ . Pick  $p \in \mathbb{N}$  with  $p > M$ . It is easy to know  $r_f f(x), s_f f(z) \in p[-e, e]$ . Thus

$$f(x - z) = r_f f(x) + s_f f(z) \in 2p[-e, e].$$

So  $f(W - W) \subset 2p[-e, e]$  for all  $f \in \Gamma$ .

Let  $B \subset X$  be bounded, and without loss of generality suppose that  $W - W$  is balanced. Since  $(W - W) \cap U \in \mathcal{N}(X)$ , pick  $q \in \mathbb{N}$  for which

$$(1/q)B \subset (W - W) \cap U.$$

Observe that  $q$  is independent of every individual  $x \in B$ . For  $x \in B$  and  $f \in \Gamma$ ,  $x = q(x/q)$  where  $(x/q) \in (W - W) \cap U$  and  $f(-(x/q)) \in 2p[-e, e]$ .

$$\begin{aligned} f(x) &= f\left(q \frac{x}{q}\right) = f\left[(q-1) \frac{x}{q} + \frac{x}{q}\right] \\ &= r_{1f} f\left[(q-1) \frac{x}{q}\right] + s_{1f} f\left(-\frac{x}{q}\right) \\ &= r_{1f} r_{2f} f\left[(q-2) \frac{x}{q}\right] + r_{1f} s_{2f} f\left(-\frac{x}{q}\right) + s_{1f} f\left(-\frac{x}{q}\right) \\ &\quad \dots \dots \\ &= r_{1f} \cdots r_{q-1f} f\left(\frac{x}{q}\right) + (r_{1f} \cdots r_{q-2f} s_{q-1f} + \cdots + r_{1f} s_{2f} + s_{1f}) f\left(-\frac{x}{q}\right) \\ &= r_{1f} \cdots r_{q-1f} r_{qf} f(0) + (r_{1f} \cdots r_{q-1f} s_{qf} + \cdots + r_{1f} s_{2f} + s_{1f}) f\left(-\frac{x}{q}\right) \\ &= (r_{1f} \cdots r_{q-1f} s_{qf} + \cdots + r_{1f} s_{2f} + s_{1f}) f\left(-\frac{x}{q}\right), \end{aligned}$$

where

$$|r_{if}| \leq M, \quad i = 1, 2, \dots, q-1$$

and

$$|s_{if}| \leq M, \quad i = 1, 2, \dots, q.$$

Obviously,

$$\begin{aligned} |r_{1f}r_{2f}\cdots r_{q-1f}s_{qf} + \cdots + r_{1f}s_{2f} + s_{1f}| &\leq M^q + M^{q-1} + \cdots + M^2 + M \\ &< p^q + p^{q-1} + \cdots + p^2 + p \\ &= \frac{p(1-p^q)}{1-p}. \end{aligned}$$

So

$$f(x) = (r_{1f}r_{2f}\cdots r_{q-1f}s_{qf} + \cdots + r_{1f}s_{2f} + s_{1f})f(-x/q) \in \frac{p(1-p^q)}{1-p}2p[-e, e]$$

for all  $f \in \Gamma$  and  $x \in B$ . Thus,  $\Gamma$  is uniformly order bounded on  $B \subset X$  and is uniformly order bounded on each bounded subset of  $X$ .  $\square$

**Theorem 2.2.** *Let  $X$  be a topological vector space of second category and  $Y$  an ordered topological vector space with the positive cone having nonempty interior. If the UB family  $\Gamma$  of continuous mappings from  $X$  to  $Y$  with  $U \in \mathcal{N}(X)$ ,  $\delta, M > 0$  and  $\varphi : (0, \delta) \rightarrow (0, +\infty)$  satisfying  $\lim_{t \rightarrow 0} \varphi(t) = 0$  is pointwise bounded or pointwise order bounded, then  $\Gamma$  is uniformly order bounded on each bounded subset of  $X$ .*

*Proof.* Let  $U$  be balanced and closed, and  $C = \{y \in Y : y \geq 0\}$  the positive cone in  $Y$ . By the hypothesis,  $C$  is closed in  $Y$  and there exists  $y_0$  in the interior of  $C$ . Obviously,  $C - y_0$  and  $y_0 - C$  are both neighborhoods of  $0 \in Y$ . Then

$$V = (C - y_0) \cap (y_0 - C)$$

is also a neighborhood of  $0 \in Y$ , and a closed and balanced convex set. Let  $x \in X$ . There exists  $\theta \in (0, 1)$  such that  $\theta x \in U$ . And since  $\{f(\theta x) : f \in \Gamma\}$  is bounded in  $Y$  there exists  $n \in \mathbb{N}$  such that  $\{f(\theta x) : f \in \Gamma\} \subset nV$ . In another case, we know  $\{f(\theta x) : f \in \Gamma\}$  is order bounded in  $Y$ . Then there exist  $a, b \in Y$  such that

$$\{f(\theta x) : f \in \Gamma\} \subset [a, b] := \{y \in Y : a \leq y \leq b\}.$$

But  $C - y_0$  and  $y_0 - C$  absorbs  $[a, b]$ . Pick  $m \in \mathbb{N}$  with  $m > n$  and

$$[a, b] \subset [-my_0, my_0] = m[-y_0, y_0].$$

Hence  $\{f(\theta x) : f \in \Gamma\} \subset m[-y_0, y_0] = mV$ . Observing  $nV \subset mV$  we obtain  $\{f(\theta x) : f \in \Gamma\} \subset mV$  in two cases.

Since  $\lim_{t \rightarrow 0} \varphi(t) = 0$  there exists  $p \in \mathbb{N}$  such that  $1/p < \delta$  and  $\varphi(1/p) \leq (1/m)$ . But

$$f((1/p)\theta x) = s_f f(\theta x), \quad \text{where } |s_f| \leq \varphi(1/p) \leq (1/m).$$

Then  $|ms_f| \leq 1$  for all  $f \in \Gamma$  and

$$\{f((1/m)\theta x) : f \in \Gamma\} = \{s_f f(\theta x) : f \in \Gamma\} \subset \cup_{f \in \Gamma} s_f mV \subset V$$

Thus  $(1/m)\theta x \in \cap_{f \in \Gamma} f^{-1}(V)$ . Let

$$W = U \cap (\cap_{f \in \Gamma} f^{-1}(V))$$

where

$$V = (C - y_0) \cap (y_0 - C) = [-y_0, y_0].$$

Obviously,  $(1/m)\theta x \in U$ . Hence,  $X = \cup_{n=1}^{\infty} nW$ . Since  $V$  is closed in  $Y$ , each  $f \in \Gamma$  is continuous and  $U$  is closed in  $X$ , we know  $W$  is closed in  $X$ . But  $X$  is of second category. So  $W - W = \{x - z : x, z \in W\} \in \mathcal{N}(X)$ .

As the same in the proof of Theorem 2.1, we can obtain the result.  $\square$

*Remark 2.3.* In Theorem 2.2, if the positive cone  $C$  in  $Y$  is normal in the weak topology, then  $\Gamma$  is uniformly bounded on each bounded subset of  $X$ . In fact,  $Y' = C' - C'$  where  $C' = \{f \in Y' : f(y) \geq 0 \text{ for all } y \in C\}$ . Then an order bounded subset in  $Y$  must be bounded in  $Y$ . See details for [12].

**Acknowledgement.** The authors wish to thank the editor and referee for their very useful suggestions.

## REFERENCES

1. N. Adasch, B. Ernst and D. Keim, *Topological Vector Spaces*, Lecture Notes in Math. Vol. 639, Springer-Verlag, Berlin, 1978.
2. A.D. Arvanitakis and J.D. Stein, *A generalization of uniform boundedness principle*, Internat. J. Pure Appl. Math. **43** (2008), 507–528.
3. S. Banach, *Theorie des Operations Lineaires*, Warszawa, 1932.
4. A. Bourass, B. Ferrahi and B.M. Schreiber, *A Random multivalued uniform boundedness principle*, Set-Valued Anal. **13** (2005), 105–124.
5. C.A. Faure and A. Frolicher, *The uniform boundedness principle for bornological vector spaces*, Arch. Math. **62** (1994), 270–277.
6. R. Henrion and A. Seeger, *Uniform boundedness of norms of convex and nonconvex process*, Numerical Functional Anal. **29** (2008), 551–573.
7. U. Kohlenbach, *A logical uniform boundedness principle for abstract metric and hyperbolic spaces*, Electronic Notes in Theoretical Computer Sci. **165** (2006), 81–93.
8. J. Kupka, *Uniform boundedness principles for regular borel vector measures*, J. Austral. Math. Soc. Ser. A **29** (1980), 206–218.
9. R. Li, S. Zhong and L. Li, *Demi-linear analysis I—basic principles*, J. Korean Math. Soc. **46** (2009), 643–656.
10. O. Naguard, *A strong boundedness principle in Banach spaces*, Proc. Amer. Math. Soc. **129** (2000), 861–863.
11. W. Roth, *A uniform boundedness theorem for locally convex cones*, Proc. Amer. Math. Soc. **125** (1998), 1973–1982.
12. H.H. Schaefer, *Topological Vector Spaces*, Springer-Verlag, New York, Heidelberg, Berlin, 1980.
13. M. Song and J. Fang, *Resonance theorems for family of quasi-homogeneous operators in fuzzy normed linear spaces*, Fuzzy Sets and Systems **159** (2008), 708–719.
14. C. Swartz, *The uniform boundedness principle for order bounded operators*, Internat. J. Math. Math. Sci. **12** (1989), 487–492.
15. L. Vesely, *Local uniform boundedness principle for families of  $\epsilon$ -monotone operators*, Non-linear Anal. **24** (1995), 1299–1304.
16. A. Wilansky, *Modern Methods in Topological Vector Spaces*, McGraw-Hill, New York, 1978.

<sup>1</sup> DEPARTMENT OF MATHEMATICS, TIANJIN UNIVERSITY, TIANJIN 300072, CHINA.  
E-mail address: [zhshuhui@yahoo.com.cn](mailto:zhshuhui@yahoo.com.cn)

<sup>2</sup> DEPARTMENT OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN 150001, CHINA.  
E-mail address: [rongluli@yahoo.com.cn](mailto:rongluli@yahoo.com.cn)