

AN OMEGA THEOREM ON DIFFERENCES OF TWO SQUARES, II

M. KÜHLEITNER

ABSTRACT. Let $\rho(n)$ denote the number of pairs $(u, v) \in \mathbb{N} \times \mathbb{Z}$ with $u^2 - v^2 = n$. Due to a formula of Sierpinski, $\rho(n)$ is closely related to the classical divisor function $d(n)$. We establish a lower bound for the remainder term in the asymptotic expansion for the Dirichlet summatory function of $\rho(n)$.

1. INTRODUCTION

As in part I of this paper [8], let $\rho(n)$ denote the number of pairs $(u, v) \in \mathbb{N} \times \mathbb{Z}$ with $u^2 - v^2 = n$. For the more general case where the square is replaced by a “ k ”-th power $k \geq 2$ see Krätzel [6], [7] and the recent paper of Nowak [9]. Due to an elementary formula of Sierpinski, our function $\rho(n)$ is closely related to the classical divisor function $d(n)$ by

$$(1) \quad \rho(n) = d(n) - 2d\left(\frac{n}{2}\right) + 2d\left(\frac{n}{4}\right),$$

where $d(\cdot) = 0$ for non-integers, due to Sierpinski.

For a large real variable x , we consider the remainder term $\theta(x)$ in the asymptotic formula

$$T(x) = \sum_{n \leq x} \rho(n) = \frac{x}{2} \log x + (2\gamma - 1)\frac{x}{2} + \theta(x),$$

where γ denotes throughout this paper the Euler-Mascheroni constant.

Upper bounds for $\theta(x)$ can be readily established as a trivial generalization of the corresponding results for the Dirichlet divisor problem. It is known that

$$D(x) = x \log x + (2\gamma - 1)x + \Delta(x)$$

with

$$\Delta(x) \ll x^{23/73} (\log x)^{461/146}.$$

Received August 4, 1997.

1980 *Mathematics Subject Classification* (1991 *Revision*). Primary 11N37.

Key words and phrases. Divisor problem, Dirichlet summatory function, asymptotic expansion.

(See Huxley [5] for this upper bound and the textbook of Krätzel [6] for an enlightening survey of the theory of Dirichlet's divisor problem and the definition of the O - and the Ω - symbols.)

Concerning lower estimates, the author proved in [8], on the basis of [1] and Hafner's method [3], that

$$\theta(x) = \Omega_+ \left((x \log x)^{1/4} (\log \log x)^{(3+2 \log 2)/4} \exp(-A \sqrt{\log \log \log x}) \right).$$

The aim of the present article is an Ω_- - result for $\theta(x)$, corresponding to that of Corrádi and Kátai [1] for the divisor problem.

Theorem.

$$T(x) = \frac{x}{2} \log x + (2\gamma - 1) \frac{x}{2} + \theta(x),$$

with

$$\theta(x) = \Omega_- \left(x^{1/4} \exp \left(c (\log \log x)^{1/4} (\log \log \log x)^{-3/4} \right) \right),$$

where c is a positive absolute constant.

2. NOTATIONS AND LEMMAS

For large real x we define P_x as the set of all primes less than or equal to x , and Q_x the set of all square-free integers composed only of primes from P_x . We write $|P_x|$ for the cardinality of P_x and $M = 2^{|P_x|}$ for the cardinality of Q_x . We then have

$$|P_x| \asymp \frac{x}{\log x} \quad \text{and} \quad M \ll \exp \left(c_1 \frac{x}{\log x} \right),$$

for some positive constant c_1 . The largest integer in Q_x is bounded by e^{2x} , since for $q \in Q_x$, we have

$$\log q \leq \sum_{p \leq x} \log p \leq 2x.$$

Let S_x be the set of numbers defined by

$$S_x = \left\{ \mu = \sum_{q \in Q_x} r_q \sqrt{q} \quad \text{where } r_q \in \{0, \pm 1\} \text{ and at least two } r_q \neq 0 \right\}.$$

Finally let

$$\eta(x) = \inf \left\{ |\sqrt{n} + 2\mu| \quad \text{with } n \in \mathbb{N}_o \text{ and } \mu \in S_x \right\},$$

and

$$q(x) = -\log(\eta(x)).$$

By a slight modification of the method used for the corresponding result in Gangadharan [2], one readily shows the following lemma.

Lemma 1. For $x \rightarrow \infty$ we have

$$x \ll q(x) \ll \exp\left(c_2 \frac{x}{\log x}\right),$$

for some positive constant c_2 .

Lemma 2. There exists a positive constant c_3 such that

$$\sum_{q \in Q_x} \frac{d(q)}{q^{3/4}} \gg \exp\left(c_3 \frac{x^{1/4}}{\log x}\right).$$

Proof. By the definition of Q_x , we have

$$\begin{aligned} \sum_{q \in Q_x} \frac{d(q)}{q^{3/4}} &= \prod_{p \leq x} (1 + 2p^{-3/4}) = \exp\left(\sum_{p \leq x} \log(1 + 2p^{-3/4})\right) \\ &\geq \exp\left(\sum_{p \leq x} p^{-3/4} + O(1)\right) \gg \exp\left(c_3 \frac{x^{1/4}}{\log x}\right). \quad \square \end{aligned}$$

As in Gangadharan [2] define for real z ,

$$V(z) = 2\left(\cos\left(\frac{z}{2}\right)\right)^2 = 1 + \frac{e^{iz} + e^{-iz}}{2},$$

and

$$T_x(u) = \prod_{q \in Q_x} V\left(u\sqrt{q} - \frac{5\pi}{4}\right).$$

Lemma 3. We have

- (1) $0 \leq T_x(u) \leq 2^M$, for all u ,
- (2) $T'_x(u) \ll M 2^M e^x$, for all u ,
- (3) $T_x(u) = T_0 + T_{1,x} + T_{2,x} + T_{3,x}$ where,

$$T_0 = 1,$$

$$T_{1,x} = \frac{e^{5\pi i/4}}{2} \sum_{q \in Q_x} e^{-iu\sqrt{q}}$$

$$T_{3,x} = \sum_{\mu \in S_x} h_\mu e^{iu\mu},$$

$T_{2,x}$ is the complex conjugate of $T_{1,x}$ and $|h_\mu| \leq 1/4$.

Proof. The proof of Lemma 3 is straightforward by the definition of $V(z)$ and $T_x(u)$.

3. PROOF OF THE THEOREM

We start with the well known Voronoi identity for

$$\Delta_1(x) \stackrel{\text{def}}{=} \int_0^x \Delta(t) dt = \frac{x}{4} + \frac{x^{3/4}}{2\sqrt{2}\pi^2} \sum_{n=1}^{\infty} \frac{d(n)}{n^{5/4}} \sin(4\pi\sqrt{nx} - \frac{\pi}{4}) + O(1).$$

Inserting this in

$$\theta(x) = \Delta(x) - 2\Delta\left(\frac{x}{2}\right) + 2\Delta\left(\frac{x}{4}\right),$$

and substituting $T = 4\pi\sqrt{x}$, we get

$$\begin{aligned} E_1(T) &\stackrel{\text{def}}{=} \int_0^T E(t) t dt \\ &= T^{3/2} \sum_{n=1}^{\infty} \frac{d(n)}{n^{5/4}} \left(\sin(T\sqrt{n} - \pi/4) - 2^{5/4} \sin(T\sqrt{n/2} - \pi/4) \right. \\ &\quad \left. + 2^{3/2} \sin(T\sqrt{n/4} - \pi/4) \right), \end{aligned}$$

with

$$E(t) = 2\pi\sqrt{2\pi} \left(\theta(t^2/16\pi^2) - 1/4 \right).$$

Define

$$P(x) = \exp\left(a \frac{x}{\log x}\right)$$

such that

$$q(x) \leq P(x) \quad \text{and} \quad M^2 \leq P(x),$$

and let

$$\sigma_x = \exp(-2P(x)).$$

Next define for fixed x ,

$$\gamma_x = \sup_{u>0} \frac{-2\pi\sqrt{2\pi} \theta(u^2/16\pi^2)}{u^{1/2+1/P(x)}}.$$

We may assume that $\gamma_x < \infty$, otherwise more than Theorem 1 would be true.

Thus

$$(2) \quad \gamma_x u^{1/2+1/P(x)} + A + E(u) \geq 0,$$

for all u , where $A = 2\pi\sqrt{2\pi}/4$.

Let

$$J_x = \sigma_x^{5/2} \int_0^{\infty} (\gamma_x u^{1/2+1/P(x)} + A + E(u)) u \exp(-\sigma_x u) T_x(u) du.$$

The next lemma provides an asymptotic expansion for J_x .

Lemma 4. For $x \rightarrow \infty$,

$$J_x = e^2 \Gamma\left(\frac{5}{2}\right) \gamma_x - \frac{1}{4} \Gamma\left(\frac{5}{2}\right) \sum_{q \in Q_x} \frac{d(q)}{q^{3/4}} + o(\gamma_x) + o(1).$$

Proof. Do deal with the first two terms of J_x , we observe that, for $r = 1$ or $r = \frac{3}{2} + \frac{1}{P(x)}$,

$$\begin{aligned} \int_0^\infty u^r \exp(-\sigma_x u) T_x(u) du &= \Gamma(1+r) \sigma_x^{-(1+r)} \\ &+ \sum_{i=1,2,3} \int_0^\infty u^r \exp(-\sigma_x u) T_{i,x}(u) du \end{aligned}$$

where $1 \leq r \leq \frac{3}{2} + \frac{1}{P(x)}$.

The part of $T_{1,x}$ contributes exactly,

$$\begin{aligned} \frac{e^{5\pi i/4}}{2} \Gamma(1+r) \sum_{q \in Q_x} \frac{1}{(\sigma_x + i\sqrt{q})^{1+r}} &\ll \sum_{q \in Q_x} q^{-(1+r)/2} \\ &\ll \sum_{q \in Q_x} 1 \ll M \ll \sqrt{P(x)} = o(\sigma_x^{-5/2}). \end{aligned}$$

The contribution of $T_{2,x} = \overline{T_{1,x}}$ is obviously no more than this. Finally $T_{3,x}$ contributes

$$\begin{aligned} \sum_{\mu \in S_x} \frac{h_\mu}{(\sigma_x + i\mu)^{1+r}} &\ll 3^M \eta(x)^{-(1+r)} \\ &\ll \exp(M \ln 3 + (1+r)(-\log \eta(x))) \ll \exp(3P(x)) = o(\sigma_x^{-5/2}). \end{aligned}$$

Next we deal with the contribution of $E(u)$ to J_x . Our first step is to integrate by parts to introduce $E_1(u)$ in the integral. Thus,

$$I \stackrel{\text{def}}{=} \int_0^\infty E(u) u \exp(-\sigma_x u) T_x(u) du = - \int_0^\infty E_1(u) \frac{d}{du} (\exp(-\sigma_x u) T_x(u)) du,$$

since $E_1(u) \ll u^{3/2}$ for large u and $E_1(0) = 0$. Inserting the series representation for $E_1(u)$ and integrating term by term, noting that the series converges absolutely for every u and uniformly on compact sets, we get

$$\begin{aligned} I &= - \sum_{n=1}^\infty \frac{d(n)}{n^{5/4}} \text{Im} (e^{-\pi i/4} I_n) + O\left(\int_0^\infty \left| \frac{d}{du} (\exp(-\sigma_x u) T_x(u)) \right| du\right) \\ &+ O\left(\int_0^\infty u^{1/2} \exp(-\sigma_x u) |T_x(u)| du\right), \end{aligned}$$

since

$$u^{3/2} \frac{d}{du} (\exp(-\sigma_x u) T_x(u)) = \frac{d}{du} (u^{3/2} \exp(-\sigma_x u) T_x(u)) - \frac{3}{2} u^{1/2} \exp(-\sigma_x u) T_x(u),$$

and

$$I_n \stackrel{\text{def}}{=} \int_0^\infty (e^{iu\sqrt{n}} - 2^{5/4} e^{iu\sqrt{n/2}} + 2^{3/2} e^{iu\sqrt{n/4}}) \frac{d}{du} (u^{3/2} \exp(-\sigma_x u) T_x(u)) du.$$

Estimating the contributions of the error terms, we see that

$$\begin{aligned} \int_0^\infty \left| \frac{d}{du} (\exp(-\sigma_x u) T_x(u)) \right| du &\leq \int_0^\infty |T_x(u)' - \sigma_x T_x(u)| \exp(-\sigma_x u) du \\ &\leq 4^M \sigma_x^{-1} + 2^M \\ &\ll \exp(c\sqrt{P(x)}) (1 + \exp(2P(x))) = o(\sigma_x^{-5/2}), \end{aligned}$$

and

$$\begin{aligned} \int_0^\infty u^{1/2} \exp(-\sigma_x u) |T_x(u)| du &\ll 2^M \int_0^\infty u^{1/2} \exp(-\sigma_x u) du \\ &\ll 2^M \sigma_x^{-3/2} \ll \exp(c\sqrt{P(x)} + 3P(x)) = o(\sigma_x^{-5/2}). \end{aligned}$$

We integrate I_n by parts once more and expand $T_x(u)$ as in (3) of Lemma 3, to get

$$\begin{aligned} I_n &= -i \sum_{k=0, \dots, 3} \int_0^\infty \left(\sqrt{n} e^{iu\sqrt{n}} - 2^{5/4} \sqrt{\frac{n}{2}} e^{iu\sqrt{n/2}} + 2^{3/2} \sqrt{\frac{n}{4}} e^{iu\sqrt{n/4}} \right) \\ &\quad \times u^{3/2} \exp(-\sigma_x u) T_{i,x}(u) du \\ &= I_0(n) + I_1(n) + I_2(n) + I_3(n), \end{aligned}$$

for short. We shall show that the main term of I_n comes from $I_1(n)$. In fact, the contribution of $I_0(n)$ is

$$\ll \sqrt{n} |\sigma_x - i\sqrt{n}|^{-5/2} \ll n^{-3/4},$$

that of $I_2(n)$ is

$$\ll \sqrt{n} \sum_{q \in Q_x} |\sigma_x - i(\sqrt{n} + \sqrt{q})|^{-5/2} \ll Mn^{-3/4}.$$

The contribution of $I_3(n)$ is bounded by

$$\begin{aligned} I_3(n) &\ll \sqrt{n} \sum_{\mu \in S_x} |\sigma_x - i(\sqrt{n} - \mu)|^{-5/2} \\ &\ll \begin{cases} \sqrt{n} 3^M (\eta(x))^{-5/2}, & \text{if } n \leq 2 \max\{|\mu| : \mu \in S_x\} \\ n^{-3/4} 3^M, & \text{else.} \end{cases} \end{aligned}$$

This $\max\{|\mu| : \mu \in S_x\}$ is bounded by Me^{cx} for some positive constant c . Hence the total contribution to I is bounded by

$$\begin{aligned} &\ll \sum_{n \leq 2Me^{cx}} \frac{d(n)}{n^{5/4}} \sqrt{n} 3^M \exp\left(-5 \log \frac{\eta(x)}{2}\right) + O\left(3^M \sigma_x^{-5/4} \sum_{n > 2Me^{cx}} \frac{d(n)}{n^2}\right) \\ &\ll 3^M \sigma_x^{-5/4} \sum_{n \leq 2Me^{cx}} n^{-3/4+\epsilon} + O(3^M \sigma_x^{-5/4}) \\ &\ll 3^M \sigma_x^{-5/4} (Me^{cx})^{1/4+\epsilon} \\ &= o(\sigma_x^{-5/2}). \end{aligned}$$

Therefore,

$$\begin{aligned} I &= -\frac{1}{2} \sum_{n=1}^{\infty} \frac{d(n)}{n^{5/4}} \operatorname{Im}\left(i \sum_{q \in Q_x} \int_0^{\infty} \left(\sqrt{n} e^{iu(\sqrt{n}-\sqrt{q})} - 2^{5/4} \sqrt{\frac{n}{2}} e^{iu(\sqrt{n/2}-\sqrt{q})}\right.\right. \\ &\quad \left.\left.+ 2^{3/2} \sqrt{\frac{n}{4}} e^{iu(\sqrt{n/4}-\sqrt{q})}\right) u^{3/2} \exp(-\sigma_x u) du\right) + o(\sigma_x^{-5/2}) \\ &= -\frac{1}{2} \sum_{q \in Q_x} \left(\frac{d(q)}{q^{5/4}} - 2^{5/4} \frac{d(2q)}{(2q)^{5/4}} + 2^{3/2} \frac{d(4q)}{(4q)^{5/4}}\right) \int_0^{\infty} \sqrt{q} u^{3/2} \exp(-\sigma_x u) du \\ &\quad + O\left(\sum_{n=1}^{\infty} \frac{d(n)}{n^{5/4}} \sum_{\substack{q \in Q_x \\ n \neq q}} \left|\int_0^{\infty} \sqrt{n} e^{iu(\sqrt{n}-\sqrt{q})} u^{3/2} \exp(-\sigma_x u) du\right|\right). \end{aligned}$$

For this last error term we get a bound exactly as above for $I_3(n)$ with M replacing the factor 3^M , since

$$\sqrt{n} - \sqrt{q} \gg (\sqrt{n} + \sqrt{q})^{-1} \gg e^{-x} \gg \exp(-P(x)),$$

for $n \leq 2 \max\{q : q \in Q_x\} \gg 2e^{2x}$ and $n \neq q$.

We get,

$$\begin{aligned} I &= -\frac{1}{2} \Gamma\left(\frac{5}{2}\right) \sigma_x^{-5/2} \left(\sum_{q \in Q_x} (d(q) - d(2q) + \frac{1}{2} d(4q))\right) q^{-3/4} + o(\sigma_x^{-5/2}) \\ &= -\frac{1}{4} \Gamma\left(\frac{5}{2}\right) \sigma_x^{-5/2} \sum_{q \in Q_x} d(q) q^{-3/4} + o(\sigma_x^{-5/2}), \end{aligned}$$

since

$$d(q) - d(2q) + \frac{1}{2}d(4q) = \frac{1}{2}d(q).$$

This completes the proof of Lemma 4. \square

Since $\sigma_x > 0$ and $J_x > 0$ by (2), we have

$$\exp\left(c \frac{x^{1/4}}{\log x}\right) \ll \sum_{q \in Q_x} d(q)q^{-3/4} \ll \gamma_x,$$

by Lemma 2 and the last assertion by Lemma 4.

Thus by the definition of γ_x there is a sequence u_x which tends to infinity with x , such that

$$-\theta(u_x^2) \gg u_x^{1/2} \exp\left(\frac{\log u_x}{P(x)} + c \frac{x^{1/4}}{\log x}\right),$$

since $\theta(u)$ is bounded for bounded u , which follows for small u from

$$\theta(u) = -\frac{u}{2} \log u - (2\gamma - 1)\frac{u}{2},$$

and is obvious for the other values of u .

Consider first the values of u_x for which

$$(3) \quad \frac{\log u_x}{P(x)} \leq c \frac{x^{1/4}}{\log x}.$$

Taking logarithms on both sides, we have

$$\log \log u_x \ll \frac{x}{\log x}.$$

Since $y^{1/4}(\log y)^{-3/4}$ is an increasing function of y for sufficiently large y , we have from (3)

$$\frac{(\log \log u_x)^{1/4}}{(\log \log \log u_x)^{3/4}} \ll \frac{x^{1/4}}{\log x},$$

from which the desired estimate follows.

Consider now those values of x for which

$$(4) \quad c \frac{x^{1/4}}{\log x} \leq \frac{\log u_x}{P(x)}.$$

We may assume that

$$\frac{(\log \log u_x)^{1/4}}{(\log \log \log u_x)^{3/4}} \gg \frac{\log u_x}{P(x)},$$

otherwise the estimate holds obviously. Taking logarithms on both sides gives

$$\log \log u_x \ll \frac{x}{\log x},$$

from which the estimate follows as above. This proves the theorem. \square

References

1. Corrádi I. K. and Kátai F., *Egy megjegyzés K. S. Gangadharan, "Two classical lattice point problems" című dolgozatához*, MTA III. Osztály Közleményei **17** (1967), 89–97.
2. Gangadharan K. S., *Two classical lattice point problems*, Proc. Cambridge Phil. Soc. **57** (1961), 699–721.
3. Hafner J. L., *New omega theorems for two classical lattice point problems*, Invent. Math. **63** (1981), 181–186.
4. ———, *New omega theorems in a weighted divisor problem*, J. Number Theory **28** (1988), 240–257.
5. Huxley M. N., *Exponential sums and lattice points II*, Proc. London Math. Soc. **66(3)** (1993), 279–301.
6. Krätzel E., *Lattice Points*, Berlin, 1988.
7. ———, *Primitive lattice points in special plane domains and a related three dimensional lattice point problem I*, Forschungsergebnisse FSU Jenam, N/87/11, 1987.
8. Kühleitner M., *An omega theorem on differences of two squares*, Acta Math. Univ. Comenianae **LXI(1)** (1992), 117–123.
9. Nowak W. G., *On differences of two k -th powers of integers*, (to appear).

M. Kühleitner, Institut für Mathematik, Universität für Bodenkultur, Gregor Mendel Straße 33, A-1180 Wien, Austria; *e-mail*: kleitner@mail.boku.ac.at