

# On Feasible Sets of Mixed Hypergraphs

Daniel Král<sup>\*</sup>

Department of Applied Mathematics and  
Institute for Theoretical Computer Science<sup>†</sup>  
Charles University,  
Malostranské náměstí 25  
118 00 Praha 1, Czech Republic  
kral@kam.ms.mff.cuni.cz

Submitted: Nov 20, 2003; Accepted: Feb 24, 2004; Published: Mar 4, 2004.

MR Subject classifications: Primary 05C15, Secondary 05C65, 05C85

Keywords: coloring of hypergraphs, mixed hypergraphs, algorithms for coloring

## Abstract

A mixed hypergraph  $H$  is a triple  $(V, \mathcal{C}, \mathcal{D})$  where  $V$  is the vertex set and  $\mathcal{C}$  and  $\mathcal{D}$  are families of subsets of  $V$ , called  $\mathcal{C}$ -edges and  $\mathcal{D}$ -edges. A vertex coloring of  $H$  is proper if each  $\mathcal{C}$ -edge contains two vertices with the same color and each  $\mathcal{D}$ -edge contains two vertices with different colors. The spectrum of  $H$  is a vector  $(r_1, \dots, r_m)$  such that there exist exactly  $r_i$  different colorings using exactly  $i$  colors,  $r_m \geq 1$  and there is no coloring using more than  $m$  colors. The feasible set of  $H$  is the set of all  $i$ 's such that  $r_i \neq 0$ .

We construct a mixed hypergraph with  $O(\sum_i \log r_i)$  vertices whose spectrum is equal to  $(r_1, \dots, r_m)$  for each vector of non-negative integers with  $r_1 = 0$ . We further prove that for any fixed finite sets of positive integers  $A_1 \subset A_2$  ( $1 \notin A_2$ ), it is NP-hard to decide whether the feasible set of a given mixed hypergraph is equal to  $A_2$  even if it is promised that it is either  $A_1$  or  $A_2$ . This fact has several interesting corollaries, e.g., that deciding whether a feasible set of a mixed hypergraph is gap-free is both NP-hard and coNP-hard.

## 1 Introduction

Graph coloring problems are intensively studied both from the theoretical point view and the algorithmic point of view. A *hypergraph* is a pair  $(V, \mathcal{E})$  where  $\mathcal{E}$  is a family of subsets

---

<sup>\*</sup>The author acknowledges partial support by GAČR 201/1999/0242, GAUK 158/1999 and KON-TAKT 338/99.

<sup>†</sup>Institute for Theoretical Computer Science is supported by Ministry of Education of Czech Republic as project LN00A056.

of  $V$  of size at least 2. The elements of  $V$  are called *vertices* and the elements of  $\mathcal{E}$  are called *edges*. A *mixed hypergraph*  $H$  is a triple  $(V, \mathcal{C}, \mathcal{D})$  where  $\mathcal{C}$  and  $\mathcal{D}$  are families of subsets of  $V$  of size at least 2. The elements of  $\mathcal{C}$  are called  *$\mathcal{C}$ -edges* and the elements of  $\mathcal{D}$  are called  *$\mathcal{D}$ -edges*. A *proper  $\ell$ -coloring*  $c$  of  $H$  is a mapping  $c : V \rightarrow \{1, \dots, \ell\}$  such that there are two vertices with Different colors in each  $\mathcal{D}$ -edge and there are two vertices with a Common color in each  $\mathcal{C}$ -edge. A proper coloring  $c$  is a *strict  $\ell$ -coloring* if it uses all  $\ell$  colors. A mixed hypergraph is *colorable* if it has a proper coloring. Mixed hypergraphs were introduced in [23]. The concept of mixed hypergraphs can find its applications in different areas, e.g. list-coloring of graphs [14], graph homomorphisms [9], coloring block designs [2, 3, 16, 17, 18], etc. The importance and interest of the concept is witnessed by a recent monograph on the subject by Voloshin [21]. As an example, we present here the following construction described in [14]: Let  $G$  be a graph and let  $L$  be a function which assigns each vertex a set of colors. A coloring  $c$  of  $G$  is a *proper list-coloring with respect to the lists  $L$*  if  $c(v) \in L(v)$  for each vertex  $v$  of  $G$  and  $c(u) \neq c(v)$  for each edge  $uv$  of  $G$ . Let  $\mathcal{L}$  be the union of the lists of all the vertices of  $G$ . Consider a mixed hypergraph  $H$  with the vertex set  $V(G) \cup \mathcal{L}$  and the following edges: a  $\mathcal{D}$ -edge  $\{u, v\}$  for each  $uv \in E(G)$ , a  $\mathcal{D}$ -edge  $\{x, y\}$  for any  $x, y \in \mathcal{L}$  ( $x \neq y$ ) and a  $\mathcal{C}$ -edge  $\{v\} \cup L(v)$  for each vertex of  $G$ .  $H$  has a proper coloring iff  $G$  has a proper list-coloring. Similar constructions have been found by the author [9] for graph homomorphisms, the channel assignment problem,  $L(p, q)$ -labelings of graphs and some other graph coloring problems.

The *feasible set*  $\mathcal{F}(H)$  of a mixed hypergraph  $H$  is the set of all  $\ell$ 's such that there exists a strict  $\ell$ -coloring of  $H$ . The (*lower*) *chromatic number*  $\chi(H)$  of  $H$  is the minimum number contained in  $\mathcal{F}(H)$  and the *upper chromatic number*  $\bar{\chi}(H)$  of  $H$  is the maximum number. The feasible set of  $H$  is *gap-free (unbroken)* if  $\mathcal{F}(H) = [\chi(H), \bar{\chi}(H)]$  where  $[a, b]$  is the set of all the integers between  $a$  and  $b$  (inclusively). If the feasible set of  $H$  contains a gap, we say it is *broken*. The *spectrum* of a mixed hypergraph  $H$  is the vector  $(r_1, \dots, r_{\bar{\chi}(H)})$  where  $r_\ell$  is the number of different strict  $\ell$ -colorings of  $H$ . Two colorings  $c_1$  and  $c_2$  are considered to be different if there is no permutation of colors changing one of them to the other, i.e., it is not true that  $c_1(u) = c_1(v)$  iff  $c_2(u) = c_2(v)$  for each two vertices  $u$  and  $v$ . We remark that the spectrum is usually defined to be a vector  $(r_1, \dots, r_n)$  where  $n$  is the number of vertices of the mixed hypergraph, but we prefer using the definition without trailing zeroes in the vector. If  $\mathcal{F}$  is a set of positive integers, we say that a mixed hypergraph  $H$  is a *realization* of  $\mathcal{F}$  if  $\mathcal{F}(H) = \mathcal{F}$ . A mixed hypergraph  $H$  is a *one-realization* of  $\mathcal{F}$  if it is a realization of  $\mathcal{F}$  and all the entries of the spectrum of  $H$  are either 0 or 1.

A necessary and sufficient condition on a set of positive integers to be the feasible set of a mixed hypergraph was proved in [6]:

**Theorem 1** *A set  $\mathcal{F}$  of positive integers is a feasible set of a mixed hypergraph iff  $1 \notin \mathcal{F}$  or  $\mathcal{F}$  is an interval. If  $1 \in \mathcal{F}$ , then all the mixed hypergraphs with this feasible set contain only  $\mathcal{C}$ -edges.*

In particular, there exists a mixed hypergraph such that its feasible set contains a gap. On the other hand, it was proved that feasible sets of mixed hypertrees [10], mixed

strong hypercacti [13] and of mixed hypergraphs with maximum degree two [11, 12] are gap-free. Feasible sets of mixed hypergraphs with maximum degree three need not to be gap-free. The feasible sets of planar mixed hypergraphs, i.e., hypergraphs whose bipartite incidence graphs of their vertices and edges are planar [4, 15], are exactly intervals  $[k_1, k_2]$ ,  $1 \leq k_1 \leq 4$ ,  $k_1 \leq k_2$  and sets  $\{2\} \cup [4, k]$ ,  $k \geq 4$  as proved in [7].

Necessary or sufficient conditions for a vector to be a spectrum of a mixed hypergraph were not addressed in detail so far. However, Voloshin [22] conjectured a sufficient condition for a vector to be the spectrum of a mixed hypergraph (Conjecture 2 in [22]): *If  $n_0, \dots, n_t$  is a sequence of positive integers such that  $n_i \geq (n_{i-1} + n_{i+1})/2$  for  $1 \leq i \leq t-1$  and  $\max\{n_{\lfloor t/2 \rfloor}, n_{\lceil t/2 \rceil}\} = \max_{0 \leq i \leq t} \{n_i\}$ , then there exists a mixed hypergraph  $H$  such that  $\chi(H) + t = \bar{\chi}(H)$  and  $H$  allows exactly  $n_i$  different strict  $(\chi(H) + i)$ -colorings ( $0 \leq i \leq t$ ).* We prove this conjecture. In fact, Theorem 3 implies that the only hypothesis needed is that  $n_0, \dots, n_t$  is a sequence of non-negative integers. Let us remark at this point that Conjecture 1 from [22] on co-perfect mixed hypergraphs was disproved in [8].

We study several problems posed in [22] (Problem 10, 11, Conjecture 2) and in [6]. In particular, we are interested in the size of the smallest (one-)realization of a given feasible set. In [6], two constructions of a mixed hypergraph with a given feasible set  $\mathcal{F}$  are presented, but both of them can have exponentially many vertices in terms of  $\max \mathcal{F}$  and  $|\mathcal{F}|$ . The second construction from [6] does not even give one-realization of  $\mathcal{F}$ . We present an algorithmic construction (Theorem 2) which gives a small one-realization for a given feasible set  $\mathcal{F}$ . The number of vertices of this realization is at most  $|\mathcal{F}| + 2 \max \mathcal{F} - 1$  and the number of edges is cubic in the number of vertices.

Theorem 2 from Section 2 can be restated as follows: Let  $(r_1, \dots, r_m)$  be a vector such that  $r_1 = 0$  and  $r_i \in \{0, 1\}$  for  $2 \leq i \leq m$ . Then, there exists a mixed hypergraph  $H$  such that the spectrum of  $H$  is  $(r_1, \dots, r_m)$ . Note that the condition  $r_1 = 0$  is the condition  $1 \notin \mathcal{F}$  mentioned earlier. We generalize this theorem in Section 3. We prove that for each vector  $(r_1, \dots, r_m)$  of non-negative integers such that  $r_1 = 0$  there exists a mixed hypergraph such that its spectrum is equal to  $(r_1, \dots, r_m)$  (Theorem 3). The number of vertices of the mixed hypergraph from Theorem 3 is  $2m + 2 \sum_{i=1, r_i \neq 0}^m (1 + \lfloor \log_2 r_i \rfloor)$  and the number of its edges is cubic in the number of its vertices. Theorem 3 provides an affirmative answer to Conjecture 2 from [22] which was mentioned above.

We deal with complexity questions related to feasible sets of mixed hypergraphs in Section 4. We prove that for any fixed finite sets of positive integers  $A_1 \subset A_2$ , it is NP-hard to decide whether the feasible set of a given mixed hypergraph  $H$  is equal to  $A_2$  even if it is promised that  $\mathcal{F}(H)$  is either  $A_1$  or  $A_2$ . This theorem has several interesting corollaries: It is NP-complete to decide whether a given mixed hypergraph is colorable, it is both NP-hard and coNP-hard for a fixed non-empty finite set of positive integers  $A$  to decide whether the feasible set of a mixed hypergraph is equal to  $A$ , it is both NP-hard and coNP-hard to decide whether the feasible set of a given mixed hypergraph is gap-free. This particular result was previously obtained in [11]. It was also known before that it is NP-hard to decide whether a given mixed hypergraph is uniquely colorable [20] and that it is NP-hard to compute the upper chromatic number even when restricted to several special classes of mixed hypergraphs [1, 10, 12, 19].

There is also no polynomial-time  $o(n)$ -approximation algorithm for the lower or the upper chromatic number unless  $P = NP$  where  $n$  is the number of vertices of an input mixed hypergraph. We remark there is an  $O(n^{\frac{(\log \log n)^2}{\log^3 n}})$ -approximation algorithm for the chromatic number of ordinary graphs [5]. Recall that an algorithm for a maximization (minimization) problem is said to be  $K$ -approximation algorithm if it always finds a solution whose value is at least  $\text{OPT}/K$  (at most  $K \cdot \text{OPT}$ ) where  $\text{OPT}$  is the value of the optimum solution.

## 2 Small realizations of feasible sets

If  $\mathcal{F} = \{m\}$ , then the complete graph of order  $m$  is the one-realization of  $\mathcal{F}$  with the fewest number of vertices. In this section, we present a one-realization with few vertices for the case  $|\mathcal{F}| \geq 2$ :

**Theorem 2** *Let  $\mathcal{F}$  be a finite non-empty set of positive integers with  $1 \notin \mathcal{F}$ . There exists a mixed hypergraph with at most  $|\mathcal{F}| + 2 \max \mathcal{F} - \min \mathcal{F}$  vertices whose feasible set is  $\mathcal{F}$  and every entry of its spectrum is 0 or 1. The number of the edges of this mixed hypergraph is cubic in the number of its vertices.*

**Proof:** The proof proceeds by induction on  $\max \mathcal{F}$ . If  $2 \notin \mathcal{F}$ , then let  $H'$  be a one-realization of  $\mathcal{F}' = \{i - 1 | i \in \mathcal{F}\}$ . Let  $H$  be the mixed hypergraph obtained from  $H'$  by adding a vertex  $x$  and  $\mathcal{D}$ -edges  $\{x, v\}$  for all  $v \in V(H')$ . This operation was also used in [6] under the name “elementary shift”. It is clear that proper  $\ell$ -colorings of  $H$  are in one-to-one correspondence with proper  $(\ell + 1)$ -colorings of  $H'$ , since the color of the vertex  $x$  has to be different from the color of any other vertex and it does not affect coloring of any edge except for the added  $\mathcal{D}$ -edges of size two. Hence,  $H$  is one-realization of  $\mathcal{F}$ . The number of vertices of  $H$  is at most  $1 + |\mathcal{F}'| + 2 \max \mathcal{F}' - \min \mathcal{F}' \leq |\mathcal{F}| + 2 \max \mathcal{F} - \min \mathcal{F}$ .

It remains to consider the case when  $\min \mathcal{F} = 2$ . The case of  $\mathcal{F} = \{2\}$  is described before Theorem 2. In the rest, we assume that  $\max \mathcal{F} > 2$ . For this purpose, we define a mixed hypergraph  $H$  with the vertex set  $\{v_2^+, \dots, v_m^+, v_1^-, \dots, v_m^-, v_1^\oplus\} \cup \{v_i^\oplus | i \in \mathcal{F} \setminus \{2, m\}\}$  where  $m = \max \mathcal{F}$ . Let  $\mathcal{F}(H) = \{c_1, \dots, c_k\}$ ,  $c_1 < \dots < c_k$ , and set  $c'_1 = 1$  and  $c'_i = c_i$  for  $2 \leq i \leq k$ . Next, we describe the edges of  $H$ . The mixed hypergraph  $H$  contains the following edges for each  $l$ ,  $2 \leq l \leq k$ :

$$\{v_i^-, v_j^+\} \text{ is a } \mathcal{D} \text{-edge for } c'_{l-1} \leq i \leq c'_l \text{ and } c'_{l-1} < j \leq c'_l \text{ with } i \neq j \quad (1)$$

$$\{v_i^-, v_{c'_{l-1}}^\oplus\} \text{ is a } \mathcal{D} \text{-edge for } c'_{l-1} < i \leq c'_l \quad (2)$$

$$\{v_i^+, v_i^-, v_j^+\} \text{ is a } \mathcal{C} \text{-edge for } c'_{l-1} < i, j \leq c'_l \text{ and } i \neq j \quad (3)$$

$$\{v_i^+, v_i^-, v_{c'_{l-1}}^\oplus\} \text{ is a } \mathcal{C} \text{-edge for } c'_{l-1} < i \leq c'_l \quad (4)$$

$$\{v_i^+, v_i^-, v_j^-\} \text{ is a } \mathcal{C} \text{-edge for } c'_{l-1} < i, j \leq c'_l \text{ and } i \neq j \quad (5)$$

$$\{v_{c'_{l-1}}^\oplus, v_{c'_{l-1}}^-, v_j^+\} \text{ is a } \mathcal{C} \text{-edge for } c'_{l-1} < j \leq c'_l \quad (6)$$

$$\{v_{c'_{l-1}}^\oplus, v_{c'_{l-1}}^-, v_j^-\} \text{ is a } \mathcal{C} \text{-edge for } c'_{l-1} < j \leq c'_l \quad (7)$$

$$\{v_{c'_l}^+, v_{c'_l}^-, v_{c'_l}^\oplus\} \text{ is a } \mathcal{C}\text{-edge if } l < k \quad (8)$$

$$\{v_{c'_{l-1}}^\oplus, v_{c'_l}^+, v_{c'_l}^\oplus\} \text{ is a } \mathcal{C}\text{-edge if } l < k \quad (9)$$

$$\{v_{c'_{l-1}}^-, v_{c'_l}^-, v_{c'_l}^\oplus\} \text{ is a } \mathcal{D}\text{-edge if } l < k \quad (10)$$

$$\{v_i^-, v_j^+, v_j^-\} \text{ is a } \mathcal{D}\text{-edge for } 1 \leq i \leq c'_l, c'_{l-1} < j \leq c'_l \text{ and } i \neq j \quad (11)$$

We prove that the mixed hypergraph  $H$  has the properties claimed in the statement of the theorem. Note that  $H$  is exactly the mixed hypergraph  $H^k$  defined in the next paragraph. In the rest, we slightly abuse the notation and we call the  $\mathcal{D}$ -edges described in (1) just  $\mathcal{D}$ -edges (1) and we call other kinds of edges in a similar way.

Let  $H^l$  be the mixed hypergraph obtained from  $H$  (for  $l \geq 2$ ) restricting to the vertices  $\{v_2^+, \dots, v_{c'_l}^+, v_1^-, \dots, v_{c'_l}^-, v_1^\oplus\} \cup \{v_i^\oplus \mid i \in \mathcal{F} \wedge 3 \leq i < c'_l\}$ . The edges of  $H^l$  are those edges of  $H$  which are fully contained in the vertex set of  $H^l$ . We prove that the following statements hold for all  $l$ ,  $2 \leq l \leq k$ :

1.  $\mathcal{F}(H^l) = \{c_1, \dots, c_l\}$
2. Any proper coloring  $c$  of  $H^l$  with  $c(v_{c'_l}^+) \neq c(v_{c'_l}^-)$  uses less than  $c_l$  colors. In addition,  $c$  satisfies that  $c(v_{c'_{l-1}}^\oplus) = c(v_{c'_l}^+)$ ,  $c(v_{c'_{l-1}}^-) = c(v_{c'_l}^-) \neq c(v_{c'_l}^+)$ .
3. Any proper coloring  $c$  of  $H^l$  with  $c(v_{c'_l}^+) = c(v_{c'_l}^-)$  uses exactly  $c_l$  colors. In addition, the coloring  $c$  colors the vertices  $v_1^-, \dots, v_{c'_l}^-$  with mutually different colors and the colors of  $v_i^-$  and  $v_i^+$  (and  $v_i^\oplus$  if it exists) are the same.
4. There is exactly one proper coloring using exactly  $\lambda$  colors for each  $\lambda \in \mathcal{F}(H^l)$ .

These four claims are proved simultaneously by induction on  $l$ .

We first deal with the case that  $l = 2$ . Let  $c$  be a proper coloring of  $H^2$ . If  $c(v_1^\oplus) \neq c(v_1^-)$ , then this coloring uses exactly two colors on the vertices  $v_1^-, \dots, v_{c'_2}^-, v_1^\oplus, v_2^+, \dots, v_{c'_2}^+$  due to the presence of  $\mathcal{C}$ -edges (6) and (7). Furthermore, the  $\mathcal{D}$ -edges (1) and (2) force that  $c(v_1^-) = c(v_2^-) = \dots = c(v_{c'_2}^-)$  and  $c(v_1^+) = c(v_2^+) = \dots = c(v_{c'_2}^+) = c(v_1^\oplus)$ . Thus, the vertices are colored as described in the second claim.

Let us now suppose that  $c(v_1^\oplus) = c(v_1^-)$ . If  $c(v_i^+) \neq c(v_i^-)$  for some  $2 \leq i \leq c'_2$ , then  $c(v_1^\oplus) \neq c(v_1^-)$  due to the presence of  $\mathcal{C}$ -edges (4) and (5) and  $\mathcal{D}$ -edges (1) and (2). Thus  $c(v_i^+) = c(v_i^-)$  for all  $2 \leq i \leq c'_2$ . The colors of  $c(v_i^-)$  for  $1 \leq i \leq c'_2$  are mutually distinct due to the presence of  $\mathcal{D}$ -edges (1) and (2). We may infer that any such coloring  $c$  assigns  $c'_2$  colors to the vertices  $v_1^-, \dots, v_{c'_2}^-, v_1^\oplus, v_2^+, \dots, v_{c'_2}^+$ . Hence, the coloring  $c$  uses exactly  $c_2 = c'_2$  colors. This finishes the proof of all the four claims for  $H^2$ . It is straightforward to check that the two given colorings of  $H^2$  are proper.

Let us prove the claims 1, 2, 3 and 4 for  $H^l$  ( $l \geq 3$ ) assuming them proved for  $H^{l-1}$ . Consider a proper coloring  $c$  of a mixed hypergraph  $H^l$ . If  $c(v_{c'_{l-1}}^-) \neq c(v_{c'_{l-1}}^+)$ , then the  $\mathcal{C}$ -edge (8) and the  $\mathcal{D}$ -edge (10) together with the second claim assure that  $c(v_{c'_{l-1}}^+) = c(v_{c'_{l-1}}^\oplus)$ . Note that in this case  $c$  uses less than  $c_{l-1}$  colors to color vertices of

$H^{l-1}$ . If  $c(v_{c'_{l-1}}^-) = c(v_{c'_{l-1}}^+)$ , then  $c(v_{c'_{l-2}}^-) = c(v_{c'_{l-2}}^\oplus) \neq c(v_{c'_{l-1}}^-)$ . Thus, the color  $c(v_{c'_{l-1}}^\oplus)$  is either  $c(v_{c'_{l-1}}^-) = c(v_{c'_{l-1}}^+)$  or  $c(v_{c'_{l-2}}^-) = c(v_{c'_{l-2}}^\oplus)$  due to the presence of the  $\mathcal{C}$ -edge (9) (and both is possible). In both cases, the coloring  $c$  must use exactly  $c_{l-1}$  colors to color vertices of  $H^{l-1}$ .

We distinguish two cases (similar to those above):  $c(v_{c'_{l-1}}^-) \neq c(v_{c'_{l-1}}^\oplus)$  and  $c(v_{c'_{l-1}}^-) = c(v_{c'_{l-1}}^\oplus)$ . If  $c(v_{c'_{l-1}}^-) \neq c(v_{c'_{l-1}}^\oplus)$ , then the same argumentation as used before yields that  $c(v_{c'_{l-1}}^-) = \dots = c(v_{c'_l}^-)$  and  $c(v_{c'_{l-1}}^\oplus) = c(v_{c'_{l-1}+1}^+) = \dots = c(v_{c'_l}^+)$ . On the other hand, if  $c(v_{c'_{l-1}}^-) = c(v_{c'_{l-1}}^\oplus)$ , then we can again infer that  $c(v_{c'_{l-1}}^\oplus) = c(v_{c'_{l-1}}^-)$ ,  $c(v_{c'_{l-1}+1}^+) = c(v_{c'_{l-1}+1}^-) \neq \dots \neq c(v_{c'_l}^+) = c(v_{c'_l}^-)$ . The colors  $c(v_{c'_{l-1}}^-), \dots, c(v_{c'_l}^-)$  are also mutually distinct because of the  $\mathcal{D}$ -edges (1) and (2) and they are different from colors  $c(v_1^-), \dots, c(v_{c'_{l-1}-1}^-)$  due to the presence of  $\mathcal{D}$ -edges (11) and the third claim used for  $H^{l-1}$ . This proves the first, the second and the third claim for  $H^l$ . We again leave a straightforward check that all the described colorings are proper. As to the fourth claim: If  $c(v_{c'_{l-1}}^-) = c(v_{c'_{l-1}}^\oplus)$ , then exactly  $c_{l-1}$  colors are used to color the vertices of  $H^{l-1}$  and new  $c_l - c_{l-1}$  colors are used to color the vertices of  $v_{c'_{l-1}+1}^+, \dots, v_{c'_l}^+$  and  $v_{c'_{l-1}+1}^-, \dots, v_{c'_l}^-$  due to the presence of  $\mathcal{D}$ -edges 11. On the other hand, if  $c(v_{c'_{l-1}}^-) \neq c(v_{c'_{l-1}}^\oplus)$ , there exists unique extension of any proper coloring of  $H^{l-1}$  to  $H^l$ . This finishes the proof of all the four claims on  $H^l$ .

We conclude that  $H = H^k$  has the desired properties. The bound on the number of edges follows from the fact that each edge has size at most three. ■

We immediately have the following corollary of Theorems 1 and 2:

**Corollary 1** *There exists a polynomial-time algorithm which for a given set  $\mathcal{F}$  decides whether it is a feasible set of some mixed hypergraph and if so it outputs a mixed hypergraph  $H$  such that  $\mathcal{F}(H) = \mathcal{F}$ .*

**Proof:** If  $1 \notin \mathcal{F}$ , then the algorithm returns the construction from Theorem 2. If  $1 \in \mathcal{F}$  and  $\mathcal{F}$  is not interval, then the algorithm returns that no such mixed hypergraph exists (Theorem 1). If  $1 \in \mathcal{F}$  and  $\mathcal{F}$  is an interval, then the algorithm outputs a mixed hypergraph consisting of  $\max \mathcal{F}$  vertices and no edges. ■

### 3 Realizations of spectra

We first slightly alter the construction from Theorem 2:

**Lemma 1** *Let  $\mathcal{F} = \{c_1, \dots, c_k\}$  be a set of positive integers with  $1 \notin \mathcal{F}$ . There exists a mixed hypergraph  $H^*$  with at most  $2(|\mathcal{F}| + \max \mathcal{F})$  vertices which is a one-realization of  $\mathcal{F}$ . Moreover,  $H^*$  contains  $3l$  vertices  $w_i^+, w_i^\oplus, w_i^\ominus$  ( $1 \leq i \leq k$ ) with the following property: Let  $c$  be any proper coloring of  $H^*$ , then*

- The vertices  $w_i^+$ ,  $w_i^\oplus$ ,  $w_i^\ominus$  are colored by  $c$  with exactly two colors for each  $i$ .
- $c(w_i^\ominus) = c(w_i^+) \neq c(w_i^\oplus)$  iff  $c$  uses exactly  $c_i$  colors.
- $c(w_i^\ominus) \neq c(w_i^+) = c(w_i^\oplus)$  iff  $c$  does not use exactly  $c_i$  colors.

**Proof:** If  $\mathcal{F} = \{2\}$ , then consider the following mixed hypergraph  $H^*$ :  $V(H^*) = \{w_1^+, w_1^\oplus, w_1^\ominus\}$  where  $\{w_1^\ominus, w_1^+\}$  is the only  $\mathcal{C}$ -edge of  $H^*$  and  $\{w_1^\ominus, w_1^\oplus\}$  is the only  $\mathcal{D}$ -edge of  $H^*$ .

Assume now that  $2 \in \mathcal{F}$  and  $\mathcal{F} \neq \{2\}$ . The case  $2 \notin \mathcal{F}$  is considered later. We extend the construction from the proof of Theorem 2. Let  $H^k$  be the mixed hypergraph obtained in the construction and let us continue using notation from the proof of Theorem 2. We add a vertex  $v_1^+$  together with a  $\mathcal{C}$ -edge  $\{v_1^+, v_1^-\}$  and a vertex  $v_{c_k}^\oplus$  together with a  $\mathcal{C}$ -edge  $\{v_{c_{k-1}}^\oplus, v_{c_k}^\oplus\}$ . It is routine to check that the following two claims hold:

- $c(v_{c_i}^-) = c(v_{c_i}^+) \neq c(v_{c_i}^\oplus)$  iff  $c$  uses exactly  $c_i$  colors.
- $c(v_{c_i}^+) = c(v_{c_i}^\oplus)$  iff  $c$  does not use exactly  $c_i$  colors.

Let  $w_i^+ = v_{c_i}^+$ ,  $w_i^- = v_{c_i}^-$  and  $w_i^\oplus = v_{c_i}^\oplus$ . We add new vertices  $w_i^\ominus$  for all  $1 \leq i \leq k$  to the mixed hypergraph together with  $\mathcal{C}$ -edges  $\{w_i^\ominus, w_i^+, w_i^\oplus\}$  and  $\{w_i^\ominus, w_i^+, w_i^-\}$  for all  $1 \leq i \leq k$ ,  $\mathcal{C}$ -edges  $\{w_i^\ominus, w_i^-, v_1^-\}$  for all  $2 \leq i \leq k$  and  $\mathcal{D}$ -edges  $\{w_i^\ominus, w_i^\oplus\}$  for all  $1 \leq i \leq k$ . We further add a  $\mathcal{C}$ -edge  $\{w_1^\ominus, w_1^-, v_2^-\}$ . The resulting mixed hypergraph is  $H^*$ .

Let  $c$  be a proper coloring of  $H^*$ . If  $c(w_i^+) \neq c(w_i^\oplus)$  (and thus  $c(w_i^+) = c(w_i^-)$ ), then the  $\mathcal{C}$ -edge  $\{w_i^\ominus, w_i^+, w_i^\oplus\}$  and the  $\mathcal{D}$ -edge  $\{w_i^\ominus, w_i^\oplus\}$  force the vertex  $w_i^\ominus$  to have the color  $c(w_i^+) = c(w_i^-)$  (and this extension is possible) — this describes the case when the coloring  $c$  uses exactly  $c_i$  colors. Let us assume further  $c(w_i^+) = c(w_i^\oplus)$ . If  $c(w_i^+) \neq c(w_i^-)$ , then the  $\mathcal{C}$ -edge  $\{w_i^\ominus, w_i^+, w_i^-\}$  and the  $\mathcal{D}$ -edge  $\{w_i^\ominus, w_i^\oplus\}$  force the vertex  $w_i^\ominus$  to have the color  $c(w_i^-)$  (and this extension is possible). If  $c(w_i^+) = c(w_i^\oplus) = c(w_i^-)$ , then the  $\mathcal{C}$ -edge  $\{w_i^\ominus, w_i^-, v_1^-\}$  (the  $\mathcal{C}$ -edge  $\{w_1^\ominus, w_1^-, v_2^-\}$  in case  $i = 1$ ) and the  $\mathcal{D}$ -edge  $\{w_i^\ominus, w_i^\oplus\}$  force the vertex  $w_i^\ominus$  to have the color  $c(v_1^-)$  (the color  $c(v_2^-)$ ). This requires that  $c(v_1^-) \neq c(w_i^+) = c(w_i^\oplus) = c(w_i^-)$  ( $c(v_2^-) \neq c(w_i^\oplus)$  where  $w_i^\oplus = v_1^\oplus$ , since  $i$  is 1 in this case). The last non-equality is assured by the presence of the  $\mathcal{D}$ -edge (11) ( $\mathcal{D}$ -edge (2)) in the construction of Theorem 2. This implies that each coloring  $c$  of  $H^k$  can be uniquely extended to  $H^*$ .

It is straightforward to check that all the three properties stated by the lemma hold. The second and the third one are established due to the presence of a  $\mathcal{C}$ -edge  $\{w_i^\ominus, w_i^+, w_i^-\}$  and a  $\mathcal{D}$ -edge  $\{w_i^\ominus, w_i^\oplus\}$  ( $1 \leq i \leq k$ ) and due to the analogous claims stated in the previous paragraph for  $v_{c_i}^+$  and  $v_{c_i}^\oplus$ . The first one is established by the presence of the  $\mathcal{C}$ -edges  $\{w_i^\ominus, w_i^+, w_i^-\}$  and  $\mathcal{D}$ -edges  $\{w_i^\ominus, w_i^\oplus\}$ .

The final case to consider is that  $2 \notin \mathcal{F}$ . In this case, we first construct a mixed hypergraph for  $\mathcal{F}' = \{i - 1 | i \in \mathcal{F}\}$  and then add a new vertex  $x$  together with  $\mathcal{D}$ -edges  $\{x, v\}$  for all vertices  $v$  as in the beginning of the proof of Theorem 2. ■

**Lemma 2** *Let  $l$  be a given positive integer. There is a mixed hypergraph  $H_l$  with three special vertices  $w^+$ ,  $w^\ominus$ ,  $w^\oplus$  which satisfies: Let  $c$  be any precoloring of  $w^+$ ,  $w^\ominus$  and  $w^\oplus$  using two colors such that  $c(w^\ominus) \neq c(w^\oplus)$ , then:*

- *Any extension of  $c$  to a proper coloring of  $H_l$  uses no additional colors.*
- *If  $c(w^\ominus) \neq c(w^+) = c(w^\oplus)$ , then  $c$  can be uniquely extended to a proper coloring of  $H_l$ .*
- *If  $c(w^\ominus) = c(w^+) \neq c(w^\oplus)$ , then  $c$  can be extended to exactly  $l$  different proper colorings of  $H_l$ .*

*The number of vertices of  $H_l$  does not exceed  $3 + 2\lfloor \log_2 l \rfloor$ .*

**Proof:** The proof proceeds by induction on  $l$ . The statement is trivial for  $l = 1$ . We distinguish two cases:

- **The number  $l$  is even.**

Let  $H_l$  be the mixed hypergraph obtained from  $H_{l/2}$  by adding a new vertex  $x$ , a  $\mathcal{C}$ -edge  $\{w^\oplus, w^\ominus, x\}$  and a  $\mathcal{D}$ -edge  $\{w^\oplus, w^+, x\}$ . If  $c(w^\ominus) \neq c(w^+) = c(w^\oplus)$ , then  $c$  can be extended uniquely to  $H_{l/2}$  and also to  $x$ , since the added edges force that  $c(x) = c(w^\ominus)$ . If  $c(w^\ominus) = c(w^+) \neq c(w^\oplus)$ , then  $c$  can be extended to  $l/2$  different proper colorings to  $H_{l/2}$  and it can be extended by setting  $c(x)$  to either  $c(w^\oplus)$  or  $c(w^\ominus)$ . Altogether, we obtain  $l$  different extensions.

- **The number  $l$  is odd.**

Let  $l = 2t + 1$  and consider the mixed hypergraph  $H_t$  with the properties described in the statement of the lemma. Let  $w'^+$ ,  $w'^\oplus$ ,  $w'^\ominus$  be the special vertices of  $H_t$ . The mixed hypergraph  $H_l$  is constructed as follows: We set the vertex  $w^\oplus$  to be  $w'^\ominus$  and  $w^\ominus$  to  $w'^\oplus$ . In addition, new vertices  $w^+$  and  $x$  are introduced. Now add  $\mathcal{C}$ -edges  $\{w^\oplus, w^\ominus, w'^+\}$  and  $\{w^\oplus, w^\ominus, x\}$ , and  $\mathcal{D}$ -edges  $\{w^\oplus, w^+, w'^+\}$  and  $\{w^\ominus, w'^+, x\}$ . If  $c(w^\ominus) \neq c(w^+) = c(w^\oplus)$ , then the added  $\mathcal{C}$ -edges and  $\mathcal{D}$ -edges force that  $c(w'^+) = c(w^\ominus)$  and  $c(x) = c(w^\oplus)$ . The coloring  $c$  can be uniquely extended to the remaining vertices of  $H_t$ . If  $c(w^\ominus) = c(w^+) \neq c(w^\oplus)$ , then  $c(w'^+)$  can be either  $c(w^\oplus)$  or  $c(w^\ominus)$ . We consider these two possibilities. If  $c(w'^+)$  is  $c(w^\ominus)$ , then  $c(x)$  has to be  $c(w^\oplus)$  and  $c$  can be uniquely extended to the remaining vertices of  $H_t$ . If  $c(w'^+)$  is  $c(w^\oplus)$ , then  $c(x)$  can be either  $c(w^\oplus)$  or  $c(w^\ominus)$  and  $c$  can be extended in  $t$  different ways to the remaining vertices of  $H_t$ . Thus,  $c$  can be extended altogether in  $2t + 1 = l$  different ways.

The bound on the number of vertices of  $H_l$  is obviously fulfilled in both the cases. ■

We combine Lemmas 1 and 2 to get the main result of this section:

**Theorem 3** *Let  $(r_1, \dots, r_m)$  be any vector of non-negative integers such that  $r_1 = 0$ . Then there exists a mixed hypergraph with at most  $2m + 2 \sum_{i=1, r_i \neq 0}^m (1 + \lfloor \log_2 r_i \rfloor)$  vertices such that its spectrum is equal to  $(r_1, \dots, r_m)$ . Moreover, the number of edges of this mixed hypergraph is cubic in the number of its vertices.*

**Proof:** Let  $\mathcal{F} = \{j | r_j \neq 0\}$  and let  $H^*$  be the mixed hypergraph from Lemma 1. We keep the notation of Lemma 1. We apply the following procedure for each  $c_i \in \mathcal{F}(H^*)$ : We add a copy of  $H_{r_{c_i}}$  from Lemma 2 to  $H^*$  and we identify vertices  $w_i^+$  and  $w^+$ ,  $w_i^\oplus$  and  $w^\oplus$  and  $w_i^\ominus$  and  $w^\ominus$ . Lemmas 1 and 2 now yield that the spectrum of the just constructed mixed hypergraph is  $(r_1, \dots, r_m)$ . The bound on the number of vertices easily follows from counting the number of the vertices of  $H^*$  and the vertices of  $H_{r_{c_i}}$  (and realizing that some of the vertices have been identified). The bound on the number of edges follows from the fact that each edge has size at most three. ■

## 4 Complexity results

The main theorem of this section is proved in a similar way as Theorem 3, except that instead of Lemma 2, we use the following lemma:

**Lemma 3** *Let  $\Phi$  be a given formula with clauses of size three and let  $n$  be the number of variables and  $m$  the number of clauses of  $\Phi$ . There exists a mixed hypergraph  $H_\Phi$  with three special vertices  $w^+$ ,  $w^\ominus$ ,  $w^\oplus$  which satisfies: Let  $c$  be any precoloring of  $w^+$ ,  $w^\ominus$  and  $w^\oplus$  using two colors such that  $c(w^\ominus) \neq c(w^\oplus)$ , then:*

- *Any extension of  $c$  to a proper coloring of  $H_\Phi$  uses no additional colors.*
- *If  $c(w^\ominus) \neq c(w^+) = c(w^\oplus)$ , then  $c$  can always be extended to a proper coloring of  $H_\Phi$ .*
- *If  $c(w^\ominus) = c(w^+) \neq c(w^\oplus)$ , then  $c$  can be extended to a proper colorings of  $H_\Phi$  iff  $\Phi$  is satisfiable.*

$H_\Phi$  has  $2n + 3$  vertices and  $3n + m$  edges.

**Proof:** Let  $x_1, \dots, x_m$  be the variables of the given formula. Let  $H_\Phi$  be a mixed hypergraph with vertices  $w^+, w^\ominus, w^\oplus, v_1^T, v_1^F, \dots, v_n^T, v_n^F$  and the following edges:

- $\mathcal{C}$ -edges  $\{w^\oplus, w^\ominus, v_i^T\}$  and  $\{w^\oplus, w^\ominus, v_i^F\}$  for  $1 \leq i \leq n$ ,
- $\mathcal{D}$ -edges  $\{v_i^T, v_i^F\}$  for  $1 \leq i \leq n$  and
- $\mathcal{D}$ -edges  $\{w^\ominus, w^+, w_i^X, w_j^Y, w_k^Z\}$  for each clause of the formula containing the variables  $x_i, x_j$  and  $x_k$  where  $X = T$  if the occurrence of  $x_i$  in the clause is positive and  $X = F$  otherwise;  $Y$  and  $Z$  are set in the same manner.

The bounds on the size of  $H_\Phi$  are clearly fulfilled.

Any extension of a precoloring  $c$  of the vertices  $w^+, w^\ominus, w^\oplus$  with  $c(w^\ominus) \neq c(w^\oplus)$  to the vertices  $v_1^T, v_1^F, \dots, v_n^T, v_n^F$  uses only the colors  $c(w^\ominus)$  and  $c(w^\oplus)$  due to the presence of  $\mathcal{C}$ -edges  $\{w^\oplus, w^\ominus, v_i^T\}$  and  $\{w^\oplus, w^\ominus, v_i^F\}$  for all  $1 \leq i \leq n$ . If  $c(w^\ominus) \neq c(w^+)$ , then all the  $\mathcal{D}$ -edges corresponding to the clauses of  $\Phi$  are properly colored already by the precoloring and thus assigning all the vertices  $v_i^T$  the color  $c(w^\oplus)$  and all the vertices  $v_i^F$  the color  $c(w^\ominus)$  yields a proper extension of  $c$ .

Let us assume in the rest of the proof that  $c(w^\ominus) = c(w^+)$ . The color  $c(w^\oplus)$  represents true and the color  $c(w^\ominus)$  represents false in our construction. The presence of  $\mathcal{D}$ -edges  $\{v_i^T, v_i^F\}$  assures that each variable and its negation have opposite values (recall that the value of  $x_i$  is represented by the color of  $v_i^T$ ). The  $\mathcal{D}$ -edges  $\{w^\ominus, w^+, v_i^X, v_j^Y, v_k^Z\}$  force that each clause contains at least one true literal (a vertex colored by the color  $c(w^\oplus)$ ). Hence,  $c$  can be extended to  $H_\Phi$  iff there is a satisfying assignment of  $\Phi$ . ■

We now combine Lemmas 1 and 3 to get the following theorem:

**Theorem 4** *Let  $A_2$  be a finite non-empty subset of  $\{2, 3, \dots\}$  and  $A_1$  a proper (possibly empty) subset of  $A_2$ . It is NP-hard to decide whether the feasible set of a given mixed hypergraph  $H$  is equal to  $A_2$  even if it is promised that  $\mathcal{F}(H)$  is either  $A_1$  or  $A_2$ .*

**Proof:** We present a reduction from the well-known NP-complete problem 3SAT. Let  $\Phi$  be a given formula with  $n$  variables and  $H_\Phi$  the mixed hypergraph from Lemma 3. Consider the mixed hypergraph  $H^*$  from Lemma 1 for the set  $\mathcal{F} = A_2 = \{c_1, \dots, c_k\}$ . Let  $A_2 \setminus A_1 = \{c_{i_1}, \dots, c_{i_{k'}}\}$ . We create  $|A_2| - |A_1| = k' \geq 1$  copies of  $H_\Phi$  and we identify the vertices  $w^\ominus, w^+, w^\oplus$  of the  $j$ -th copy with the vertices  $w_{c_{i_j}}^\ominus, w_{c_{i_j}}^+, w_{c_{i_j}}^\oplus$  of  $H^*$ . Let  $H$  be the obtained mixed hypergraph.

It is easy to verify that  $H$  can be constructed in time polynomial in the number of variables and clauses of the formula  $\Phi$ . In particular, the number of vertices of  $H$  is at most  $3 \cdot \max A_2 + 2 \cdot |A_2 \setminus A_1| \cdot n$  and the number of its edges is cubic in the number of its vertices.

The mixed hypergraph  $H$  has a strict  $\ell$ -coloring for  $\ell \in A_1$  since any strict  $\ell$ -coloring of  $H^*$  can be extended to the copies of  $H_\Phi$  due to Lemma 3. Recall that  $c(w_{c_{i_j}}^\ominus) \neq c(w_{c_{i_j}}^+) = c(w_{c_{i_j}}^\oplus)$  for  $1 \leq j \leq k'$  for every strict  $\ell$ -coloring of  $H^*$  with  $\ell \in A_1$ . On the other hand,  $H$  has a strict  $\ell$ -coloring for  $\ell \in A_2 \setminus A_1$  iff  $\Phi$  is satisfiable: Since it holds that  $c(w_{c_{i_j}}^\ominus) = c(w_{c_{i_j}}^+) \neq c(w_{c_{i_j}}^\oplus)$  for every strict  $c_{i_j}$ -coloring  $c$  of  $H^*$ , the coloring  $c$  can be extended to the  $j$ -th copy of  $H_\Phi$  iff  $\Phi$  is satisfiable by Lemma 3. ■

Several interesting computational complexity corollaries follow almost immediately. These results are new except for Corollary 4 which was proved in a weaker form in [11]:

**Corollary 2** *It is NP-complete to decide whether a given mixed hypergraph  $H$  is colorable.*

**Proof:** This problem clearly belongs to the class NP. It is enough to set  $A_1 = \emptyset$  and  $A_2 = \{2\}$  in Theorem 4 to get the result. ■

**Corollary 3** *Let  $A$  be a fixed finite subset of  $\{2, 3, \dots\}$ . It is coNP-hard to decide whether the feasible set of a given mixed hypergraph  $H$  is equal to  $A$ . If  $A \neq \emptyset$ , then this problem is NP-hard, too.*

**Proof:** The coNP-hardness is established by setting  $A_1 = A$  and  $A_2$  to a proper finite superset of  $A$  omitting 1 in Theorem 4. The NP-hardness is established by setting  $A_2$  to  $A$  and  $A_1$  to a proper subset of  $A$ . ■

**Corollary 4** *It is both NP-hard and coNP-hard to decide whether the feasible set of a given mixed hypergraph  $H$  is gap-free even for a mixed hypergraph  $H$  with  $\bar{\chi}(H) = 4$ .*

**Proof:** The NP-hardness is established by setting  $A_1 = \{2, 4\}$  and  $A_2 = \{2, 3, 4\}$  in Theorem 4. The coNP-hardness is established by setting  $A_1 = \{4\}$  and  $A_2 = \{2, 4\}$ . ■

**Corollary 5** *There is no polynomial-time  $o(n)$ -approximation algorithm for the lower or the upper chromatic number of a mixed hypergraph where  $n$  is the number of its vertices unless  $P = NP$ .*

**Proof:** Suppose that there is a polynomial-time  $f(n)$ -approximation algorithm for the lower chromatic number where  $f(n) \in o(n)$  and  $n$  is the number of vertices of a given mixed hypergraph. Let  $\Phi$  be a given formula with clauses of size three with  $N$  variables. Choose  $m$  such that  $m > 2 \cdot f(3m + 2N)$ . It is not hard to see that there is such an integer  $m \in O(N)$  since  $f(n) \in o(n)$ . Let  $H$  be the mixed hypergraph from the construction of Theorem 4 for  $A_1 = \{m\}$  and  $A_2 = \{2, m\}$ . Note that the number of vertices of  $H$  is at most  $3k + 2N$ . The approximation algorithm for the lower chromatic number outputs a number which is less than  $m$  iff the feasible set of the input mixed hypergraph is  $A_2$ . Recall that  $\mathcal{F}(H) = A_2$  iff  $\Phi$  is satisfiable. Hence, the existence of the polynomial-time  $o(n)$ -approximation algorithm implies that  $3SAT \in P$  and consequently  $P = NP$ . The non-existence (unless  $P=NP$ ) of a polynomial-time  $o(n)$ -approximation algorithm for the upper chromatic number can be proved similarly. ■

## 5 Conclusion

There exists a mixed hypergraph whose feasible set is  $\mathcal{F}$  for any set  $\mathcal{F}$  of positive integers with  $1 \notin \mathcal{F}$ . We proved that there exists a mixed hypergraph whose spectrum is  $(r_1, \dots, r_m)$  for any vector  $(r_1, \dots, r_m)$  of non-negative integers such that  $r_1 = 0$ . The number of vertices of the smallest mixed hypergraph which is a realization of a given set  $\mathcal{F}$  has been substantially decreased from exponential to linear in  $\max \mathcal{F}$ . But the following question has not been answered: What is the number of vertices of the smallest mixed hypergraph whose feasible set is equal to a given set  $\mathcal{F}$ ? Or even, what is the number of vertices of the smallest mixed hypergraph whose spectrum is equal to a given spectrum  $(r_1, \dots, r_m)$ ? The answer to any of these questions probably requires some very fine analysis.

We have not dealt with mixed hypergraphs containing only  $\mathcal{C}$ -edges in this paper. It is clear that if  $r_1 \neq 0$  (this is equivalent to the fact that a mixed hypergraph contains only  $\mathcal{C}$ -edges), then  $r_1 = 1$ . Furthermore,  $r_2 = (2^n - 2)/2$  for some  $n$  since  $\mathcal{C}$ -edges of size two can be contracted without affecting the spectrum and any two-coloring of a mixed hypergraph on  $n$ -vertices with no  $\mathcal{D}$ -edges and with no  $\mathcal{C}$ -edges of size two is proper. This leads to the following problem: What are necessary and sufficient conditions for a vector  $(r_1, \dots, r_m)$  with  $r_1 = 1$  to be the spectrum of a mixed hypergraph?

## Acknowledgement

The author would like to thank Jiří Sgall for suggesting an easy way to prove Lemma 2, Jan Kratochvíl for helpful discussions, Zdeněk Dvořák for a careful reading of a preliminary version of this paper and the anonymous referee for helpful comments on the presentation of the results and pointing out several related references.

## References

- [1] C. Arbib, M. Flammini: On the upper chromatic number of  $(v_3, b_2)$ -configurations, *Discrete Appl. Math.* **120** (2002), 3–12.
- [2] C. Colbourn, J. Dinitz, A. Rosa: Bicoloring Triple Systems, *Electronic J. Combin.* **6(1)** (1999), #R25.
- [3] Ch. J. Colbourn, A. Rosa: Triple Systems, Clarendon Press, Oxford (1999), sect. 18.6. Strict colouring and the upper chromatic number, 340–341.
- [4] Z. Dvořák, D. Král': On Planar Mixed Hypergraphs, *Electronic J. Combin.* **8 (1)** (2001), #R35.
- [5] M. M. Halldórsson: A still better performance guarantee for approximate graph coloring, *Inform. Process. Lett.* **45** (1993), 19–23.

- [6] T. Jiang, D. Mubayi, Zs. Tuza, V. Voloshin, D. B. West: The chromatic spectrum of mixed hypergraphs, *Graphs and Combin.* **18(2)** (2002), 309–318.
- [7] D. Kobler, A. Kündgen: Gaps in the chromatic spectrum of face-constrained plane graphs, *Electronic J. Combin.* **8 (1)** (2001), #N3.
- [8] D. Král': A counter-example to Voloshin's hypergraph co-perfectness conjecture, *Australasian Journal on Combinatorics* **27** (2003), 253–262.
- [9] D. Král': Mixed hypergraphs and other graph coloring problems, submitted. Available as ITI report 2003-160.
- [10] D. Král', J. Kratochvíl, A. Proskurowski, H.-J. Voss: Coloring mixed hypertrees, to appear in *Discrete Appl. Math.*.
- [11] D. Král', J. Kratochvíl, H.-J. Voss: Complexity Note on Mixed Hypergraphs, in: Proc. 26th International Symposium, Mathematical Foundations of Computer Science 2001 (MFCS'01), LNCS vol. 2136 (2001), Springer-Verlag, 474–486.
- [12] D. Král', J. Kratochvíl, H.-J. Voss: Mixed Hypergraphs with Bounded Degree: Edge-Coloring of Mixed Multigraphs, *Theoretical Comp. Science* **295** (2003), 263–278.
- [13] D. Král', J. Kratochvíl, H.-J. Voss: Mixed hypercacti, to appear in *Discrete Math.*.
- [14] J. Kratochvíl, Z. Tuza, M. Voigt: New trends in the theory of graph colorings: Choosability and list coloring, in: Contemporary Trends in Discrete Mathematics (from DIMACS and DIMATIA to the future), DIMACS Series in Discrete Mathematics and Theoretical Computer Science, vol. 49, American Mathematical Society, Providence, RI (1999), 183–197.
- [15] A. Kündgen, E. Mendelsohn, V. Voloshin: Colouring planar mixed hypergraphs, *Electronic J. Combin.* **7** (2000), #R60.
- [16] H. Lefmann, V. Rödl, R. Thomas: Monochromatic vs. multicolored paths, *Graphs Combin.* **8** (1992), 323–332.
- [17] L. Milazzo, Zs. Tuza: Upper chromatic number of Steiner triple and quadruple systems, *Discrete Math.* **174** (1997), 247–259.
- [18] L. Milazzo, Zs. Tuza: Strict colorings for classes of Steiner triple systems, *Discrete Math.* **182** (1998), 233–243.
- [19] R. Scapellato, V. Voloshin: Some results on coloring of mixed hypergraphs, *Rend. Semin. Mat. Messina Ser. II 20* **5** (1999), 149–165.
- [20] Zs. Tuza, V. Voloshin, H. Zhou: Uniquely colorable mixed hypergraphs, *Discrete Math.* **248** (2002), 221–236.

- [21] V. Voloshin: Coloring mixed hypergraphs: Theory, algorithms and applications, AMS, Providence, 2002.
- [22] V. Voloshin: On the upper chromatic number of a mixed hypergraph, *Australasian Journal of Combin.* **11** (1995), 25–45.
- [23] V. Voloshin: The mixed hypergraphs, *Computer Science Journal of Moldova* **1** (1993), 45–52.