

Distinguishing infinite graphs

Wilfried Imrich

Montanuniversität Leoben, A-8700 Leoben, Austria
wilfried.imrich@mu-leoben.at

Sandi Klavžar*

Department of Mathematics and Computer Science
FNM, University of Maribor
Gosposvetska cesta 84, 2000 Maribor, Slovenia
sandi.klavzar@uni-mb.si

Vladimir Trofimov[†]

Institute of Mathematics and Mechanics
Russian Academy of Sciences
S. Kovalevskoy 16, 620219 Ekaterinburg, Russia
trofimov@imm.uran.ru

Submitted: Dec 9, 2006; Accepted: May 2, 2007; Published: May 11, 2007

Mathematics Subject Classifications: 05C25, 05C80, 03E10

Abstract

The distinguishing number $D(G)$ of a graph G is the least cardinal number \aleph such that G has a labeling with \aleph labels that is only preserved by the trivial automorphism. We show that the distinguishing number of the countable random graph is two, that tree-like graphs with not more than continuum many vertices have distinguishing number two, and determine the distinguishing number of many classes of infinite Cartesian products. For instance, $D(Q_{\aleph}) = 2$, where Q_{\aleph} is the infinite hypercube of dimension \aleph .

1 Introduction

The distinguishing number is a symmetry related graph invariant that was introduced by Albertson and Collins [2] and extensively studied afterward. In the last couple of

*Supported in part by the Ministry of Science of Slovenia under the grants P1-0297 and BI-AT/07-08-011.

[†]Supported in part by the Russian Foundation for Basic Research under the grant 06-01-00378. The work was done in part during the visit of Montanuniversität Leoben, Leoben, Austria in May, 2006.

years an amazing number of papers have been written on the topic. Let us mention just those that are directly related to our paper: It was proved independently in [6] and [12] that for a finite connected graph G , $D(G) \leq \Delta + 1$, where Δ is the largest degree of G , with equality if and only if G is a complete graph, a regular complete bipartite graph, or C_5 . For the computation of the distinguishing numbers of finite trees see [5]. Also, the distinguishing number of Cartesian products of finite graphs has been widely studied; see [1, 3, 8, 10, 11, 13].

In this paper we extend the study of the distinguishing number to infinite graphs. The starting point is the observation in [10] that the distinguishing number of the Cartesian product of two countable complete graphs is 2. The proof is surprisingly simple, just as one can easily show that the distinguishing number of the hypercube of countable dimension is 2 (cf. the first part of the proof of Theorem 5.4).

It turns out that many of the results for finite graphs can easily be generalized to countably infinite ones and that, with some additional effort, one can consider graphs with cardinality \mathfrak{c} , that is, the cardinality of the continuum. For higher cardinalities things are more complicated and are mainly treated in the last section. We wish to add that although we use “naive” set theory, it can be easily seen that our proofs remain in force in ZFC set theory.

In this paper we prove the following results for infinite graphs G : We prove that the distinguishing number of the (countable) random graph is 2, and prove that tree-like graphs (comprising all trees without pendant vertices and many other graphs) on at most \mathfrak{c} vertices have distinguishing number at most 2. For higher cardinalities we treat products of complete graphs and show that $D(Q_{\mathfrak{n}}) = 2$ where $Q_{\mathfrak{n}}$ is the hypercube of arbitrary infinite dimension.

2 Preliminaries

Let G be a graph and X a set. Then an X -labeling of G is just a mapping $V(G) \rightarrow X$. In most cases X will be a set of numbers, but we will also use the set $\{\text{black}, \text{white}\}$ to label graphs. When X will be clear from the context we will simply speak of *labelings* of G . Let g be an automorphism of G and l a labeling of G . Then we say that l is *preserved* by g if $l(v) = l(g(v))$ for any $v \in V(G)$. A labeling l of a graph G is *distinguishing* if l is only preserved by the trivial automorphism of G . The *distinguishing number* of a graph G is the least cardinal number \aleph such that there exists a distinguishing X -labeling of G with $|X| = \aleph$. The distinguishing number of a graph G is denoted by $D(G)$. An X -labeling l_1 of a graph G_1 and an X -labeling l_2 of a graph G_2 are *isomorphic* if there exists an isomorphism $\varphi : V(G_1) \rightarrow V(G_2)$ of G_1 and G_2 such that $l_1(v) = l_2(\varphi(v))$ for any $v \in V(G_1)$ (in this case we also say that X -labeled graphs G_1 and G_2 are isomorphic).

As usual, \aleph_0 stands for the cardinality of a countable set and \mathfrak{c} or 2^{\aleph_0} for the cardinality of reals. We will use \mathfrak{n} to denote an arbitrary finite or infinite cardinal.

The *Cartesian product* $G \square H$ of two finite or infinite graphs G and H is a graph with the vertex set $V(G) \times V(H)$, the vertices (g, h) and (g', h') being adjacent if $g = g'$ and $hh' \in E(H)$, or $h = h'$ and $gg' \in E(G)$. Let g be a vertex of G . Then the set of vertices

$\{(g, h) \mid h \in V(H)\}$ induces a subgraph of $G \square H$ isomorphic to H . It is called an H -fiber and denoted ${}^g H$. Analogously, for $h \in V(H)$, the G -fiber G^h is the subgraph of $G \square H$ induced by $\{(g, h) \mid g \in V(H)\}$.

Let $G_\iota, \iota \in I$, be an arbitrary family of graphs. Then the Cartesian product $\square_{\iota \in I} G_\iota$ is defined on the set of all functions $x : \iota \mapsto x_\iota, x_\iota \in G_\iota$, where two vertices (functions) are adjacent if there exists an index $\kappa \in I$ such that $x_\kappa y_\kappa \in E(G_\kappa)$ and $x_\iota = y_\iota$ for all $\iota \neq \kappa$.

We wish to point out that the Cartesian product of infinitely many nontrivial graphs is disconnected. Therefore, in this case one considers connected components of the Cartesian product and calls them *weak Cartesian products*. If all factors have transitive group any two components are isomorphic. Since we are only interested in connected graphs, for us a *Cartesian product graph* will always mean a weak Cartesian product graph.

For instance, the infinite hypercube $Q_{\mathfrak{n}}$ of dimension \mathfrak{n} is defined as the Cartesian product of \mathfrak{n} copies of K_2 . Its vertex set is the set of all functions of a set S of cardinality \mathfrak{n} into $\{0, 1\}$, where the preimage of 1 is finite, and where two vertices f, g are adjacent if there exists one and only one element x of S such that $f(x) \neq g(x)$. Clearly $Q_{\mathfrak{n}}$ is idempotent, that is $Q_{\mathfrak{n}} \cong Q_{\mathfrak{n}} \square Q_{\mathfrak{n}}$.

Let x be a vertex of a graph G . Then the neighborhood of x in G will be denoted $N_G(x)$ or $N(x)$ if G is clear from the context. In addition, $B_x(d)$ is the set of vertices of G at distance $\leq d$ from x .

By a *tree-like graph* we mean a connected graph G that contains a vertex x with the following property: for any vertex y there exists an up-vertex z with respect to x (that is, a vertex with $d(x, z) = d(x, y) + 1$), such that y is the only vertex from $N(z)$ that lies in an x, z -geodesic. In other words, for any vertex y there exists a vertex z such that $\{y\} = N(z) \cap B_x(d(x, z) - 1)$.

For terms not defined here, in particular for the Cartesian product of graphs and its properties, we refer to [9].

We have already mentioned that for a finite connected graph G , $D(G) \leq \Delta + 1$, where Δ is the largest degree of G . We conclude the section with the following analogue for infinite graphs.

Theorem 2.1 *Let \mathfrak{n} be a cardinal number, and G a connected, infinite graph such that the degree of any vertex of G is not greater than \mathfrak{n} . Then $D(G) \leq \mathfrak{n}$.*

Proof. Suppose first that \mathfrak{n} is a finite cardinal. Then G is an infinite graph of bounded, finite degree and G contains a one-way infinite isometric path R . Let u_0 be the first vertex of R , and label all vertices of R with Δ , where Δ is the largest degree of G . No other vertex will receive the label Δ . Since u_0 is labeled Δ and adjacent to exactly one vertex labeled Δ , while any other vertex of R is adjacent to two vertices labeled Δ , u_0 is fixed by every automorphism of G . Consequently, R is fixed pointwise by every automorphism. Construct a BFS-tree of G with root u_0 and label the vertices of G in BFS order as follows: Let x be a vertex in this order, then label its neighbors that are not yet labeled with different labels from $\{1, 2, \dots, \Delta - 1\}$. The BFS order implies that the labeling is well-defined and that every newly labeled vertex is fixed by every automorphism.

Assume next that the degree of G is not bounded. The vertex set of G is the union of the vertices in the BFS-levels of G . Since the number of vertices in every such level is at most \mathbf{n} we have that $|G| \leq \mathbf{n}$. \square

We note that an infinite locally finite graph can have infinite distinguishing number. For a simple example consider the graph that is obtained from the one-way infinite path v_0, v_1, v_2, \dots by attaching n pendant vertices to v_n for any $n \geq 1$.

3 Distinguishing the random graph

In this section we determine the distinguishing number of the countable random graph. So, let R be the countable random graph defined in [7]. Then (see, for example, Propositions 4.1, 5.1, 8.1 and 3.1 in [4]) R contains any countable graph as an induced subgraph, the automorphism group $\text{Aut}(R)$ of R is a vertex-transitive group of order 2^{\aleph_0} , and R has the following property:

- (*) For any finite disjoint subsets X and Y of $V(R)$, there are infinitely many vertices z of R such that $zx \in E(R)$ for all $x \in X$ and $zy \notin E(R)$ for all $y \in Y$.

In particular, R is a connected graph of diameter 2 and of countable degree.

Theorem 3.1 *Let R be the countable random graph. Then $D(R) = 2$.*

Proof. Since $\text{Aut}(R) \neq 1$, we have $D(R) > 1$. Let u be a vertex of R , and let $N(u) = \{v_1, v_2, \dots\}$ be the set of neighbors of u in R . Fix a sequence n_1, n_2, \dots of positive integers such that $n_i > n_1 + \dots + n_{i-1}$ for all $i > 1$. Define a labeling $l : V(R) \rightarrow \{0, 1\}$ in the following way.

Put $l(u) = 0$ and $l(v_i) = 0$ for all $i \geq 1$. By (*), there exist (distinct) vertices $w_{1,1}, \dots, w_{1,n_1}$ in $N(v_1) \setminus (\{u\} \cup N(u))$. Put $l(w_{1,j}) = 0$ for all $j, 1 \leq j \leq n_1$, and $l(w) = 1$ for all other vertices w from $N(v_1) \setminus (\{u\} \cup N(u))$. Suppose now that, for some positive integer k , l is defined on $\cup_{i=1}^k (N(v_i) \setminus (\{u\} \cup N(u)))$ in such a way that, for each $i \leq k$, there are exactly n_i vertices in $N(v_i) \setminus (\{u\} \cup N(u))$ labeled 0. Then the number n of vertices in $(N(v_{k+1}) \setminus (\{u\} \cup N(u))) \cap (\cup_{i=1}^k (N(v_i) \setminus (\{u\} \cup N(u))))$ labeled 0 (by assumption, l is defined on $\cup_{i=1}^k (N(v_i) \setminus (\{u\} \cup N(u)))$) is $\leq n_1 + \dots + n_k < n_{k+1}$. By (*), there exist (distinct) vertices $w_{k+1,1}, \dots, w_{k+1,n_{k+1}-n}$ in $N(v_{k+1}) \setminus ((\{u\} \cup N(u)) \cup (\cup_{i=1}^k N(v_i)))$. Put $l(w_{k+1,j}) = 0$ for all $j, 1 \leq j \leq n_{k+1} - n$, and $l(w) = 1$ for all other vertices w from $N(v_{k+1}) \setminus ((\{u\} \cup N(u)) \cup (\cup_{i=1}^k N(v_i)))$. As a result, for any positive integer $i \leq k + 1$, l is defined on $N(v_i) \setminus (\{u\} \cup N(u))$ and there are exactly n_i vertices in $N(v_i) \setminus (\{u\} \cup N(u))$ labeled 0. Since the diameter of R is 2, continuing in this way we get a labeling $l : V(R) \rightarrow \{0, 1\}$.

By the definition of l , all vertices in $\{u\} \cup N(u) = \{u, v_1, v_2, \dots\}$ are labeled 0, and, for each positive integer i , the number of vertices in $N(v_i) \setminus (\{u\} \cup N(u))$ labeled 0 is n_i . Furthermore, u is the only vertex of R for which all neighbors are labeled 0. In fact, suppose u' is another vertex of R for which all neighbors are labeled 0. By (*), there exists a vertex w of R such that wu', ww_1 and $w_{1,1}, \dots, w_{1,n_1}$ (where $w_{1,1}, \dots, w_{1,n_1}$ are

all vertices in $N(v_1) \setminus (\{u\} \cup N(u))$ labeled 0) are edges of R , while $wu \notin E(R)$. But $wv_1 \in E(R)$ and $wu \notin E(R)$ imply $w \in N(v_1) \setminus (\{u\} \cup N(u))$. At the same time, for each j , $1 \leq j \leq n_1$, we have $w \neq w_{1,j}$ by $ww_{1,j} \in E(R)$. Thus $l(w) = 1$ contrary to $wu' \in E(R)$.

Let g be an automorphism of R preserving the labeling l . By the above, $g(u) = u$ and hence $g(v_i) = v_i$ for all positive integers i . Suppose $g(w) \neq w$ for some vertex w of R . By $(*)$, there exists a vertex v of R such that vu and vw are edges of R , while $vg(w) \notin E(R)$. Since $vu \in E(R)$, we have $v = v_i$ for some positive integer i , and hence $g(v) = v$. Now $vw \in E(R)$ while $g(v)g(w) = vg(w) \notin E(R)$, a contradiction.

Thus the labeling l is distinguishing. □

4 Tree-like graphs

We now move to uncountable graphs. The main result of this section asserts that the distinguishing number of tree-like graphs of cardinality not greater than \mathfrak{c} is either one or two, but the result is no longer true for larger cardinalities. We need the following somewhat technical result.

Lemma 4.1 *Let T be a tree with $u \in V(T)$ such that $1 \leq \deg(u) \leq 2^{\aleph_0}$ and $1 < \deg(v) \leq 2^{\aleph_0}$ for all $v \in V(T) \setminus \{u\}$. Then there exist 2^{\aleph_0} pairwise nonisomorphic distinguishing $\{0, 1\}$ -labelings l_α^\pm , $\alpha \in A$ (where $|A| = 2^{\aleph_0}$), of T such that, for each $\alpha \in A$, $l_\alpha^-(u) = 0$, $l_\alpha^+(u) = 1$ and with respect to each of l_α^- and l_α^+ , u is the only vertex of T with all neighbors in T labeled 0.*

Proof. Let N_α , $\alpha \in A$ where $|A| = 2^{\aleph_0}$, be distinct subsets of the set of integers > 2 with $|N_\alpha| > 1$ for all α . For each $\alpha \in A$, let S_α be the set of (infinite) sequences (i_1, i_2, \dots) such that $\{i_1, i_2, \dots\} = N_\alpha$. Of course, $|S_\alpha| = 2^{\aleph_0}$ for all $\alpha \in A$. For each $\alpha \in A$ and each $s = (i_1, i_2, \dots) \in S_\alpha$, let \bar{s} be the $\{0, 1\}$ -sequence $(\epsilon_0, \epsilon_1, \dots)$ with $\epsilon_j = 0$ if and only if $j = i_1, i_1 + i_2, i_1 + i_2 + i_3, \dots$. Put $\bar{S}_\alpha = \{\bar{s} : s \in S_\alpha\}$. Of course, $|\bar{S}_\alpha| = |S_\alpha| = 2^{\aleph_0}$ for all $\alpha \in A$.

Now let T be a tree with $u \in V(T)$ such that $\deg(v) > 1$ for all $v \in V(T) \setminus \{u\}$ and $\deg(v) \leq 2^{\aleph_0}$ for all $v \in V(T)$. For each $\alpha \in A$, we define a distinguishing $\{0, 1\}$ -labeling l_α^- such that u and all neighbors of u in T are labeled 0, and u is the only vertex of T with all neighbors in T labeled 0.

Since $|V(T)| \leq 2^{\aleph_0}$, there is an injection $v \mapsto s_v$ from $V(T)$ into \bar{S}_α . Define l_α^- inductively by Steps.

As Step 1, put $l_\alpha^-(u) = 0$ and $l_\alpha^-(v) = 0$ for all neighbors v of u in T .

Suppose now that after Step n , n a positive integer, we have a labeling of some vertices of T such that

- (i) _{n} all vertices at distance $\leq n$ from u in T are labeled;
- (ii) _{n} if a vertex v from $V(T) \setminus \{u\}$ is labeled and $v' \in V(T) \setminus \{v\}$ with $d_T(u, v') + d_T(v', v) = d_T(u, v)$ then v' and all neighbors of v' in T are labeled;

(iii_n) all neighbors of any vertex of T labeled 1 are labeled.

If not all vertices of T are labeled after Step n (otherwise the definition of l_α^- is completed), Step $n + 1$ is defined as follows:

Let $\{v_\beta : \beta \in B\}$ be the set of all vertices of T which are unlabeled but have labeled neighbors (at least one; note that by (iii_n) all these neighbors are labeled 0). For each $\beta \in B$, choose an infinite path $P_{v_\beta} : v_{\beta,0} = v_\beta, v_{\beta,1}, \dots$ of T such that $d_T(u, v_{\beta,j}) - d_T(u, v_{\beta,0}) = j$ for all non-negative integers j . By (ii_n), the vertices of P_{v_β} are unlabeled. We label vertices of P_{v_β} using the sequence \bar{s}_{v_β} . Let $X_{v_\beta}^0$ be the set of vertices v of T such that, for some non-negative integer j , v is a neighbor of $v_{\beta,j}$ different from $v_{\beta,j+1}$, $d_T(u, v) = d_T(u, v_{\beta,j}) + 1$ and $v_{\beta,j}$ is labeled 0. Analogously, let $X_{v_\beta}^1$ be the set of vertices v of T such that, for some non-negative integer j , v is a neighbor of $v_{\beta,j}$ different from $v_{\beta,j+1}$, $d_T(u, v) = d_T(u, v_{\beta,j}) + 1$ and $v_{\beta,j}$ is labeled 1. Let Y_{v_β} be the set of vertices v' of T such that v' is a neighbor of some $v \in X_{v_\beta}^1$ with $d_T(u, v') = d_T(u, v) + 1$. Note that, by (ii_n), no vertex in $X_{v_\beta}^0$, $X_{v_\beta}^1$ or Y_{v_β} is labeled. Put $l_\alpha^-(v) = 0$ for all $v \in X_{v_\beta}^0 \cup X_{v_\beta}^1 \cup Y_{v_\beta}$. Using the same procedure for all v_β , where $\beta \in B$, we complete Step $n + 1$. It is easy to see that after Step $n + 1$ we get a partial labeling of T which has properties (i_{n+1}) – (iii_{n+1}).

We now show that the resulting labeling $l_\alpha^- : V(T) \rightarrow \{0, 1\}$ is distinguishing. Since all integers in N_α are > 2 , it follows from the definition of l_α^- that u is the only vertex of T with all neighbors in T labeled 0. Hence any automorphism of T preserving l_α^- stabilizes u . Note now that infinite paths x_0, x_1, \dots of T with the following properties a) – d):

- a) $d_T(u, x_j) - d_T(u, x_0) = j$ for all non-negative integers j ,
- b) if $l_\alpha^-(x_j) = 0$ for some non-negative integer j , then $l_\alpha^-(x_{j+1}) = 1$,
- c) $l_\alpha^-(x_0) = 1$ and $d_T(u, x_0) \geq 2$,
- d) if w is the neighbor of x_0 with $d_T(u, w) = d_T(u, x_0) - 1$, and w' is the neighbor of w with $d_T(u, w') = d_T(u, w) - 1$, then $l_\alpha^-(w) = l_\alpha^-(w') = 0$,

are exactly the paths P_{v_β} used in one of the Steps. Indeed, any path $P_{v_\beta} = x_0, x_1, \dots$ obviously has properties a) – d). On the other hand, if an infinite path x_0, x_1, \dots of T has properties a) – d), then c) implies that x_0 is a vertex of some path of the form P_{v_β} . Furthermore, by d) we have $x_0 = v_\beta$. Now a) and b) imply that x_0, x_1, \dots coincides with P_{v_β} .

By the definition of l_α^- , distinct such paths have distinct labelings, thus any automorphism of T preserving l_α^- stabilizes each such path. Since for each vertex v of T , there exists a vertex v' with $d_T(u, v) + d_T(v, v') = d_T(u, v')$ which is a vertex of one such path, l_α^- is a distinguishing labeling.

Define l_α^+ by $l_\alpha^+(u) = 1$ and $l_\alpha^+(v) = l_\alpha^-(v)$ for all $v \in V(T) \setminus \{u\}$. Then, with respect to l_α^+ , u is the only vertex of T with all neighbors in T labeled 0. Since l_α^- is a distinguishing labeling, it follows that l_α^+ is a distinguishing labeling as well.

The labels of each path of the form P_{v_β} from the definition of l_α^- make it possible to reconstruct the corresponding set N_α . Therefore the labelings l_α^- of T for distinct $\alpha \in A$ are (pairwise) nonisomorphic. Hence labelings l_α^+ of T for distinct $\alpha \in A$ are (pairwise) nonisomorphic as well. Besides, for $\alpha, \alpha' \in A$, l_α^- and $l_{\alpha'}^+$ are nonisomorphic labelings of T , since, with respect to each of l_α^- and $l_{\alpha'}^+$, u is the only vertex of T with all neighbors in

T labeled 0 while $l_\alpha^-(u) \neq l_{\alpha'}^+(u)$. □

Theorem 4.2 *Let G be a tree-like graph with $\Delta \leq 2^{\aleph_0}$. Then $D(G) \leq 2$.*

Proof. Recall that every tree-like graph G contains a vertex x with the following property: for any $y \in V(G)$ there exists a $z \in V(G)$ such that $\{y\} = N(z) \cap B_x(d(x, z) - 1)$.

Let T be a spanning subgraph of G with the property that $x'x'' \in E(T)$ if and only if $\{x'\} = N(x'') \cap B_x(d(x, x'') - 1)$ or $\{x''\} = N(x') \cap B_x(d(x, x') - 1)$.

Clearly T is a forest with no finite component, and uniquely defined. Furthermore, for any connected component C of T , there exists a unique vertex u_C of C at a smallest distance from x , and the valency in T of any vertex of C , different from u_C , is > 1 . Of course $x = u_{C'}$ for the connected component C' of T containing x , and the set of neighbors of x in T coincides with the set of neighbors of x in G .

Define a labeling $l : V(G) \rightarrow \{0, 1\}$ as follows. By Lemma 4.1, there exists a distinguishing labeling l^- of C' such that $l^-(x) = 0$ and $l^-(v) = 0$ for all neighbors v of x in T (or, equivalently, in G). Put $l(w) = l^-(w)$ for all vertices w of C' . Let $\{C_i : i \in I\}$ be the set of connected components of T different from C' . Then $|I| \leq 2^{\aleph_0}$. Hence, by Lemma 4.1, there exist distinguishing labelings l_i^+ of C_i for all $i \in I$ such that $l_i^+(u_{C_i}) = 1$, u_{C_i} is the only vertex of C_i for which all neighbors in C_i are l_i^+ -labeled 0, and, for any distinct $i', i'' \in I$, the $l_{i'}^+$ -labeled tree $C_{i'}$ and the $l_{i''}^+$ -labeled tree $C_{i''}$ are not isomorphic. For each $i \in I$ and each $w \in C_i$, put $l(w) = l_i^+(w)$. This completes the definition of l .

Let g be an arbitrary automorphism of G preserving l . By the definition of l , x is the only vertex of G labeled 0 for which all neighbors in G are also labeled 0. Therefore $g(x) = x$ and hence $g \in \text{Aut}(T)$. Since the restrictions of l to distinct connected components of T give pairwise nonisomorphic labeled graphs, g stabilizes each connected component of T . As the restrictions of l to any connected component C of T is a distinguishing labeling of C , we infer that $g = 1$.

Thus l is a distinguishing labeling. □

Let \mathfrak{n} be a cardinal $> 2^{\aleph_0}$ and T a tree with $x \in V(T)$ such that $\deg(x) = \mathfrak{n}$ and $\deg(y) = 2$ for all $y \in V(T) \setminus \{x\}$. Of course, $D(T) \leq |V(T)| = \mathfrak{n}$. On the other side, for any set M with $|M| < \mathfrak{n}$, there are $< \mathfrak{n}$ distinct sequences of elements of M . Thus, for any M -labeling l of T , there exist two distinct infinite paths x'_1, x'_2, \dots and x''_1, x''_2, \dots of T with $d_T(x, x'_i) = d_T(x, x''_i) = i$ for all positive integers i such that $l(x'_i) = l(x''_i)$ for all positive integers i . The automorphism g of T defined by $g(x'_i) = x''_i$ and $g(x''_i) = x'_i$ for all positive integers i , and $g(y) = y$ for all $y \in V(T) \setminus \{x'_1, x'_2, \dots, x''_1, x''_2, \dots\}$, is a non-trivial automorphism preserving l . Hence $D(T) = \mathfrak{n}$. Thus the condition $\deg(v) \leq 2^{\aleph_0}$ in the hypothesis of Theorem 4.2 cannot be weakened (even for trees and even for a single vertex).

We close the section by noting that countable trees have also independently been treated by Tucker [15].

5 Cartesian products of arbitrary cardinality

In this section we generalize results about the distinguishing number of Cartesian products of complete graphs [8, 10] to the infinite case and find many similarities. We also determine the distinguishing number of infinite hypercubes of arbitrary dimension.

To this end we take recourse to elementary results about infinite cardinals and ordinals, transfinite induction, and the well-ordering theorem; see e.g. [14].

We also apply several basic facts about prime factorizations and automorphisms of connected infinite graphs with respect to the weak Cartesian product: Every connected, infinite graph can be uniquely represented as the weak Cartesian product of prime graphs, that is, graphs that are not the product of two nontrivial graphs; see [9]. Moreover, given a product

$$G = \square_{i \in I} G_i$$

of prime graphs, every automorphism of G is induced by transpositions of isomorphic factors and automorphisms of the factors themselves. That is, if $\psi \in \text{Aut}(G)$ then there exists a permutation π of I together with isomorphisms $\psi_i : G_i \rightarrow G_{\pi i}$ such that

$$(\psi v)_i = \psi_{\pi^{-1} i} v_{\pi^{-1} i}.$$

Also, complete graphs are prime with respect to the Cartesian product.

Theorem 5.1 *Let \mathfrak{n} be an infinite cardinal number. Then $D(K_{\mathfrak{n}} \square K_{\mathfrak{n}}) = 2$.*

Proof. Let $G = G_1 \square G_2$, where both G_1 and G_2 are isomorphic to $K_{\mathfrak{n}}$. Let $V = V(G_1) = V(G_2)$ and \prec a well-ordering of V with the first element a_0 .

We label every vertex $(x, y) \in G_1 \square G_2$ white if $x \prec y$, otherwise we label it black.

This labelling is distinguishing. To see this we first note that every G_1 -fiber has infinitely many black vertices, but that the G_2 -fiber ${}^{a_0}G_2$ has only one black vertex, namely (a_0, a_0) . Thus, every automorphism of G is generated by automorphisms of the factors. In other words, every automorphism maps G_1 -fibers into G_1 -fibers and G_2 -fibers into G_2 -fibers.

Since $G_1^{a_0}$ is the only G_1 -fiber with no white vertices it is stabilized by all automorphisms. Similarly, ${}^{a_0}G_2$ is stabilized by all automorphisms of G because it is the only G_2 -fiber with only one black vertex. Thus $(a_0, a_0) = G_1^{a_0} \cap {}^{a_0}G_2$ is fixed by all $\alpha \in \text{Aut}(G)$.

If all diagonal elements (x, x) are fixed, then G_1^x and xG_2 are stabilized for every x . Thus $G_1^x \cap {}^yG_2 = (x, y)$ is fixed for every $(x, y) \in V(G)$ and the labeling is distinguishing.

Hence, if the labeling is not distinguishing there is a smallest ordinal c in V such that (c, c) is not fixed by the label preserving automorphisms of G , but the vertices $(x, y) \in V(G)$ with $x, y \prec c$ are. Let L be the union of the G_1 - and G_2 -fibers through these vertices. Clearly L is stabilized by the all label preserving automorphisms. Moreover, G_1^c is the only G_1 -fiber with no white vertex outside of L and cG_2 the only G_2 -fiber with just one black vertex outside of L . Thus (c, c) is also fixed. \square

Theorem 5.2 *Let \mathfrak{n} be an infinite cardinal. Then $D(K_{\mathfrak{n}} \square K_{2^{\mathfrak{n}}}) = 2$.*

Proof. We consider the complete graphs $G_2 = K_{\mathfrak{n}}$ with vertex set W and $H = K_{2^{\mathfrak{n}}}$ with vertex set V . Of course, W is a set of cardinality \mathfrak{n} and V a set of cardinality $2^{\mathfrak{n}}$. Write V as a disjoint union $V' \cup V''$, where V' is a set of cardinality \mathfrak{n} and V'' a set of cardinality $2^{\mathfrak{n}}$. Let G_1 be the subgraph of H induced by V' and G_0 the subgraph of H induced by V'' . Then $|G_2| = |G_1| = \mathfrak{n}$ and $|G_0| = |H| = 2^{\mathfrak{n}}$.

Consider the $2^{\mathfrak{n}}$ different black and white labelings of G_2 and assign them to the G_2 -fibers of $G_2 \square H$, yielding a labeling l on $G_2 \square H$. The labeling l induces labelings l' of $G_2 \square G_1$ and l'' of $G_2 \square G_0$. By Theorem 5.1 we can choose l such that the labeling l' is distinguishing.

If l'' is not distinguishing, there must be a nontrivial l'' -preserving automorphism γ of $G_2 \square G_0$. This mapping is of the form

$$\gamma(x, y) = (\alpha x, \beta y),$$

where $\alpha \in \text{Aut}(G_2)$ and $\beta \in \text{Aut}(G_0)$. If α is the identity mapping ι , then $\beta = \iota$ too, because any two G_2 -fibers have different labelings. We can thus assume that $\alpha \neq \iota$.

We now consider the mapping

$$\delta : (x, y) \mapsto (\alpha x, y)$$

of $G_2 \square H$ into itself. This mapping permutes the labelings of the G_2 -fibers of $G_2 \square H$. In other words, there is permutation β' of $V(H)$ such that

$$(x, y) \mapsto (\alpha x, \beta' y)$$

preserves l . By the definition of γ we infer that $\beta'|_{V(G_0)} = \beta$. But then $\beta'|_{V(G_1)}$ is a permutation of $V(G_0)$. But this is impossible, because l' is distinguishing. \square

If the cardinality of the second factor is bigger than $2^{\mathfrak{n}}$ the situation is strikingly different.

Proposition 5.3 *Let $2^{\mathfrak{n}} < \mathfrak{m}$. Then $D(K_{\mathfrak{n}} \square K_{\mathfrak{m}}) > \mathfrak{n}$.*

Proof. The number of labelings of $K_{\mathfrak{n}}$ with $\mathfrak{a} \leq \mathfrak{n}$ labels is $\mathfrak{a}^{\mathfrak{n}} \leq 2^{\mathfrak{n}} < \mathfrak{m}$. \square

If the generalized continuum hypothesis does not hold, there are cardinals \mathfrak{n} and \mathfrak{m} such that $\mathfrak{n} < \mathfrak{m} < 2^{\mathfrak{n}}$. We do not know whether $D(K_{\mathfrak{n}} \square K_{\mathfrak{m}}) = 2$ in this case.

Theorem 5.4 *Let \mathfrak{n} be an infinite cardinal. Then $D(Q_{\mathfrak{n}}) = 2$.*

Proof. The proof is by transfinite induction. Let $\mathfrak{n} = \aleph_0$. The vertices of Q_{\aleph_0} are all 01-sequences with finitely many ones. Let v_i be the vertex whose first i coordinates are 1, and the others 0. We consider the one sided infinite path

$$P = v_0 v_1 \dots$$

and label its vertices black, all the other vertices white. Clearly P is fixed by every automorphism of Q_{\aleph_0} . Also, P meets every Θ -class of Q_{\aleph_0} in exactly one edge. Since this edge is fixed, all Θ -classes are stabilized and its edges cannot be inverted. Thus the labeling is distinguishing.

Furthermore, take all vertices with an even number of ones, throw out those that are in $N(P)$, and call this set A . Then A is an infinite independent set of $Q_{\aleph_0} \setminus N(P)$. We now create 2^{\aleph_0} labelings of Q_{\aleph_0} in which the vertices of P are labeled black, the vertices of A in all possible ways, and the remaining vertices white. Since no two vertices of A are adjacent and no vertex of A is a neighbor of a vertex in P , each such labeling contains exactly one black vertex (the vertex with all zeros) with exactly one black neighbor. Note also that given two copies of Q_{\aleph_0} with distinct labelings created as above, there is no label preserving isomorphism between them.

We show next how to find a distinguishing labeling of Q_{\aleph_1} . To this end we choose \aleph_1 hypercubes Q_{\aleph_0} and label them with distinct nonisomorphic distinguishing labelings. This is possible, because $\aleph_1 \leq 2^{\aleph_0}$. Let these hypercubes, together with their labelings, be G_ι , $\iota \in I$, $|I| = \aleph_1$. Note that every G_ι has exactly one black vertex that has exactly one black neighbor, we call it the root of G_ι . We now consider the weak Cartesian product

$$G = \square_{\iota \in I} G_\iota$$

with respect to the vertex v_0 whose coordinates are the roots of the G_ι . We also label G such that every unit fiber, that is, the G_ι -fibers through the root vertex, inherits the labeling of its factors. All other vertices are labeled white.

Clearly, G is isomorphic to Q_{\aleph_1} . Also, since v_0 is the only vertex of G that is labeled black and has \aleph_1 black neighbors, it is fixed by all automorphisms. Clearly this implies that the unit fibers are also fixed and the labeling is distinguishing.

Note that the subgraph induced by the black vertices in G has one connected component and isolated vertices. Moreover, the black component has exactly one vertex of degree \aleph_1 , all other vertices have smaller degree.

Again we can find an independent set A of cardinality \aleph_1 where no vertex of A is adjacent to a unit fiber vertex. This we can use to find 2^{\aleph_1} nonisomorphic distinguishing labelings of G . Every such labeling consists of isolated black vertices and one connected subgraph of black vertices that has a vertex, the root of G , of maximum degree \aleph_1 .

Suppose \mathfrak{n} is a cardinal with the following properties:

- (i) There are $2^{\mathfrak{n}}$ nonisomorphic distinguishing labelings of $Q_{\mathfrak{n}}$.
- (ii) In every one of these labelings the black vertices induce a subgraph consisting of isolated vertices and a large connected component with exactly one vertex of degree \mathfrak{n} which we call the root.

Then the above arguments imply that the successor cardinal \mathfrak{n}' of \mathfrak{n} also satisfies (i) and (ii).

In order to complete the proof it remains to be shown that a limit cardinal \mathfrak{n} satisfies (i) and (ii) if every cardinal $\mathfrak{m} < \mathfrak{n}$ does.

For every \mathfrak{m} we select a graph $G_{\mathfrak{m}} \cong Q_{\mathfrak{m}}$ together with a distinguishing labeling satisfying (i) and (ii).

We can assume that the distinguishing labelings of the $G_{\mathfrak{m}}$ have the property that to any two K_2 -factors of $G_{\mathfrak{m}}$ there is a black (isolated) vertex that projects into the vertex of the K_2 -factor that is different from the root. We now consider the Cartesian product $G_{\mathfrak{m}} \square G_{\mathfrak{m}'}$, $\mathfrak{m} \neq \mathfrak{m}'$, where the unit fibers inherit the labeling from $G_{\mathfrak{m}}$ and $G_{\mathfrak{m}'}$, and where all other vertices are white. This means that no automorphism can map a K_2 -factor of $G_{\mathfrak{m}}$ into one of $G_{\mathfrak{m}'}$. Again we consider the weak Cartesian product

$$G = \square G_{\mathfrak{m}}$$

where the components of the root vertex are the roots of the $G_{\mathfrak{m}}$. Again we let the unit fibers inherit the labelings of the respective factors and label everything else white. Clearly this yields a distinguishing labeling of G . As before we can find an independent set A of cardinality \mathfrak{n} that contains no vertex that is adjacent to a unit fiber. Labeling A in $2^{\mathfrak{n}}$ distinct ways in black and white yields $2^{\mathfrak{n}}$ nonisomorphic distinguishing labelings of G . Clearly

$$G = Q_{\sum_{\mathfrak{m} < \mathfrak{n}} \mathfrak{m}} = Q_{\mathfrak{n}}$$

and satisfies (i) and (ii). □

We wish to remark that this actually means that we can distinguish the Cartesian and not only the weak Cartesian product of \mathfrak{n} copies of K_2 by two colors.

Concluding remarks

Very recently (Apr 4, 2007) a closely related paper [16] was published. We note that Theorem 3.1 of [16] is a special case of our Theorem 4.2.

We thank a referee for carefully reading the manuscript and numerous helpful remarks.

References

- [1] M. O. Albertson, Distinguishing Cartesian powers of graphs, *Electron. J. Combin.* 12 (2005) #N17.
- [2] M. O. Albertson and K. L. Collins, Symmetry breaking in graphs, *Electron. J. Combin.* 3 (1996) #R18.
- [3] B. Bogstad and L. Cowen, The distinguishing number of hypercubes, *Discrete Math.* 283 (2004) 29–35.
- [4] P. Cameron, The random graph, pp. 333–351 in *The Mathematics of Paul Erdős, II* (ed. R. L. Graham and J. Nešetřil), *Algorithms and Combinatorics, 14*, Springer, Berlin, 1997.
- [5] C. T. Cheng, On computing the distinguishing numbers of trees and forests, *Electron. J. Combin.* 13 (2006) #R11.

- [6] K. L. Collins and A. N. Trenk, The distinguishing chromatic number, *Electron. J. Combin.* 13 (2006) #R16.
- [7] P. Erdős and A. Rényi, Asymmetric graphs, *Acta Math. Acad. Sci. Hungar.* 14 (1963) 295–315.
- [8] M. J. Fisher, G. Isaak, Distinguishing colorings of Cartesian products of complete graphs, *Discrete Math.*, to appear.
- [9] W. Imrich and S. Klavžar, *Product Graphs: Structure and Recognition*, John Wiley & Sons, New York, 2000.
- [10] W. Imrich, J. Jerebic and S. Klavžar, Distinguishing Cartesian product graphs, *European J. Combin.*, to appear.
- [11] W. Imrich and S. Klavžar, Distinguishing Cartesian powers of graphs, *J. Graph Theory* 53 (2006) 250–260.
- [12] S. Klavžar, T.-L. Wong and X. Zhu, Distinguishing labelings of group action on vector spaces and graphs, *J. Algebra* 303 (2006) 626–641.
- [13] S. Klavžar and X. Zhu, Cartesian powers of graphs can be distinguished by two labels, *European J. Combin.* 28 (2007) 303–310.
- [14] K. Kuratowski and A. Mostowski, *Set Theory*, North-Holland, Amsterdam, 1968.
- [15] T. Tucker, Distinguishability of maps, preprint, 2005.
- [16] M. E. Watkins and X. Zhou, Distinguishability of locally finite trees, *Electron. J. Combin.* 14 (2007) #R29.