

The valuations of the near polygon \mathbb{G}_n

Bart De Bruyn*

Department of Pure Mathematics and Computer Algebra
Ghent University, Gent, Belgium
bdb@cage.ugent.be

Submitted: Aug 7, 2009; Accepted: Nov 4, 2009; Published: Nov 13, 2009

Mathematics Subject Classifications: 51A50, 05B25, 51A45, 51E12

Abstract

We show that every valuation of the near $2n$ -gon \mathbb{G}_n , $n \geq 2$, is induced by a unique classical valuation of the dual polar space $DH(2n - 1, 4)$ into which \mathbb{G}_n is isometrically embeddable.

1 Basic definitions and main results

A *near polygon* is a connected partial linear space $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \text{I})$, $\text{I} \subseteq \mathcal{P} \times \mathcal{L}$, with the property that for every point x and every line L , there exists a unique point on L nearest to x . Here, distances $d(\cdot, \cdot)$ are measured in the collinearity graph Γ of \mathcal{S} . If d is the diameter of Γ , then the near polygon is called a *near $2d$ -gon*. A near 0-gon is a point and a near 2-gon is a line. Near quadrangles are usually called generalized quadrangles. If X_1 and X_2 are two nonempty sets of points of \mathcal{S} , then $d(X_1, X_2)$ denotes the smallest distance between a point of X_1 and a point of X_2 . If X_1 is a singleton $\{x\}$, then we will also write $d(x, X_2)$ instead of $d(\{x\}, X_2)$. For every $i \in \mathbb{N}$ and every nonempty set X of points of \mathcal{S} , $\Gamma_i(X)$ denotes the set of all points $x \in X$ for which $d(x, X) = i$. If X is a singleton $\{x\}$, then we will also write $\Gamma_i(x)$ instead of $\Gamma_i(\{x\})$.

Let \mathcal{S} be a near polygon. A set X of points of \mathcal{S} is called a *subspace* if every line of \mathcal{S} having two of its points in X has all its points in X . If X is a subspace, then we denote by \tilde{X} the subgeometry of \mathcal{S} induced on the point set X by those lines of \mathcal{S} which have all their points in X . A set X of points of \mathcal{S} is called *convex* if every point on a shortest path between two points of X is also contained in X . If X is a non-empty convex subspace of \mathcal{S} , then \tilde{X} is also a near polygon. Clearly, the intersection of any number of (convex) subspaces is again a (convex) subspace. If $*_1, *_2, \dots, *_k$ are $k \geq 1$ objects (i.e., points or nonempty sets of points) of \mathcal{S} , then $\langle *_1, *_2, \dots, *_k \rangle$ denotes the smallest convex subspace

*Postdoctoral Fellow of the Research Foundation - Flanders

of \mathcal{S} containing $*_1, *_2, \dots, *_k$. The set $\langle *_1, *_2, \dots, *_k \rangle$ is well-defined since it equals the intersection of all convex subspaces containing $*_1, *_2, \dots, *_k$.

A near polygon \mathcal{S} is called *dense* if every line is incident with at least three points and if every two points at distance 2 have at least two common neighbors. If x and y are two points of a dense near polygon \mathcal{S} at distance δ from each other, then by Brouwer and Wilbrink [6, Theorem 4], $\langle x, y \rangle$ is the unique convex subspace of diameter δ containing x and y . The convex subspace $\langle x, y \rangle$ is called a *quad* if $\delta = 2$, a *hex* if $\delta = 3$ and a *max* if $\delta = n - 1$. We will now describe two classes of dense near polygons.

(I) Let $n \geq 2$, let \mathbb{K}' be a field with involutory automorphism ψ and let \mathbb{K} denote the fixed field of ψ . Let V be a $2n$ -dimensional vector space over \mathbb{K}' equipped with a nondegenerate skew- ψ -Hermitian form f_V of maximal Witt index n . The subspaces of V which are totally isotropic with respect to f_V define a Hermitian polar space $H(2n - 1, \mathbb{K}'/\mathbb{K})$. We denote the corresponding Hermitian dual polar space by $DH(2n - 1, \mathbb{K}'/\mathbb{K})$. So, $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ is the point-line geometry whose points, respectively lines, are the n -dimensional, respectively $(n - 1)$ -dimensional, subspaces of V which are totally isotropic with respect to f_V , with incidence being reverse containment. The dual polar space $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ is a dense near $2n$ -gon. In the finite case, we have $\mathbb{K} \cong \mathbb{F}_q$ and $\mathbb{K}' \cong \mathbb{F}_{q^2}$ for some prime power q . In this case, we will denote $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ also by $DH(2n - 1, q^2)$. The dual polar space $DH(3, q^2)$ is isomorphic to the generalized quadrangle $Q^-(5, q)$ described in Payne and Thas [24, Section 3.1].

(II) Let $n \geq 2$, let V be a $2n$ -dimensional vector space over \mathbb{F}_4 with basis $B = \{\bar{e}_1, \bar{e}_2, \dots, \bar{e}_{2n}\}$. The *support* of a vector $\bar{x} = \sum_{i=1}^{2n} \lambda_i \bar{e}_i$ of V is the set of all $i \in \{1, \dots, 2n\}$ satisfying $\lambda_i \neq 0$; the cardinality of the support of \bar{x} is called the *weight* of \bar{x} . Now, we can define the following point-line geometry $\mathbb{G}_n(V, B)$. The points of $\mathbb{G}_n(V, B)$ are the n -dimensional subspaces of V which are generated by n vectors of weight 2 whose supports are two by two disjoint. The lines of $\mathbb{G}_n(V, B)$ are of two types:

(a) *Special lines*: these are $(n - 1)$ -dimensional subspaces of V which are generated by $n - 1$ vectors of weight 2 whose supports are two by two disjoint.

(b) *Ordinary lines*: these are $(n - 1)$ -dimensional subspaces of V which are generated by $n - 2$ vectors of weight 2 and 1 vector of weight 4 such that the $n - 1$ supports associated with these vectors are mutually disjoint.

Incidence is reverse containment. By De Bruyn [10] (see also [11, Section 6.3]), the geometry $\mathbb{G}_n(V, B)$ is a dense near $2n$ -gon with three points on each line. The isomorphism class of the geometry $\mathbb{G}_n(V, B)$ is independent from the vector space V and the basis B of V . We will denote by \mathbb{G}_n any suitable element of this isomorphism class. The near polygon \mathbb{G}_2 is isomorphic to the generalized quadrangle $Q^-(5, 2)$.

Now, endow the vector space V with the (skew-)Hermitian form f_V which is linear in the first argument, semi-linear in the second argument and which satisfies $f_V(\bar{e}_i, \bar{e}_j) = \delta_{ij}$ for all $i, j \in \{1, \dots, 2n\}$. With the pair (V, f_V) , there is associated a Hermitian dual polar space $DH(V, B) \cong DH(2n - 1, 4)$, and every point of $\mathbb{G}_n(V, B)$ is also a point of $DH(V, B)$. By [10] or [11, Section 6.3], the set X of points of $\mathbb{G}_n(V, B)$ is a subspace of $DH(V, B)$ and the following two properties hold:

(1) $\tilde{X} = \mathbb{G}_n(V, B)$;

(2) If x and y are two points of X , then the distance between x and y in \tilde{X} equals the distance between x and y in $DH(V, B)$.

Properties (1) and (2) imply that the near polygon \mathbb{G}_n admits a full and isometric embedding into the dual polar space $DH(2n - 1, 4)$. It can be shown that there exists up to isomorphism a unique such isometric embedding, see De Bruyn [16].

Suppose $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \mathbb{I})$ is a dense near polygon. A function $f : \mathcal{P} \rightarrow \mathbb{N}$ is called a *valuation* of \mathcal{S} if it satisfies the following properties:

(V1) $f^{-1}(0) \neq \emptyset$.

(V2) Every line L contains a unique point x_L with smallest f -value and $f(x) = f(x_L) + 1$ for every point $x \in L \setminus \{x_L\}$.

(V3) Through every point x of \mathcal{S} , there exists a (necessarily unique) convex subspace F_x such that the following holds for any point y of F_x : (i) $f(y) \leq f(x)$; (ii) if z is a point collinear with y such that $f(z) = f(y) - 1$, then $z \in F_x$.

Valuations of dense near polygons were introduced in De Bruyn and Vandecasteele [18] and are a very important tool for classifying dense near polygons. For several classes of dense near polygons, see De Bruyn [14, Corollary 1.4], it can be shown that Property (V3) is a consequence of Property (V2). This is also the case for the Hermitian dual polar space $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ and the dense near polygon \mathbb{G}_n ($n \geq 2$). We now describe two classes of valuations of a dense near polygon $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \mathbb{I})$ which were also mentioned in [18].

(1) For every point x of \mathcal{S} , the map $\mathcal{P} \rightarrow \mathbb{N}; y \mapsto d(x, y)$ is a valuation of \mathcal{S} . This valuation is called the *classical valuation of \mathcal{S} with center x* .

(2) Suppose F is a (not necessarily convex) subspace of \mathcal{S} satisfying the following properties: (i) \tilde{F} is a dense near polygon; (ii) if x and y are two points of F , then the distance between x and y in \tilde{F} equals the distance between x and y in \mathcal{S} . If f is a valuation of \mathcal{S} and if $m = \min\{f(y) \mid y \in F\}$, then the map $F \rightarrow \mathbb{N}; x \mapsto f(x) - m$ is a valuation of \tilde{F} . This valuation is called the *valuation of \tilde{F} induced by f* .

By Theorem 6.8 of De Bruyn [11], every valuation of the dual polar space $DH(2n - 1, 4)$, $n \geq 2$, is classical. What about valuations of the near polygon \mathbb{G}_n ? If we regard \mathbb{G}_n as a subgeometry of $DH(2n - 1, 4)$ which is isometrically embedded into $DH(2n - 1, 4)$, then we know by the above discussion that every (classical) valuation of $DH(2n - 1, 4)$ will induce a valuation of \mathbb{G}_n . Is the converse also true: is every valuation of \mathbb{G}_n induced by some valuation of $DH(2n - 1, 4)$? The main result of this paper gives a positive answer to this question.

Theorem 1.1 *Regard \mathbb{G}_n , $n \geq 2$, as a subgeometry of $DH(2n - 1, 4)$ which is isometrically embedded into $DH(2n - 1, 4)$. Then every valuation of \mathbb{G}_n is induced by a unique (classical) valuation of $DH(2n - 1, 4)$.*

We will prove Theorem 1.1 by induction on n . The case $n = 2$ is trivial since $\mathbb{G}_2 \cong Q^-(5, 2) \cong DH(3, 4)$. The cases $n = 3$ and $n = 4$ were respectively treated in De Bruyn & Vandecasteele [19, Proposition 7.7] and [21, Proposition 6.13]. We will make use of the results of [21] to obtain a proof of Theorem 1.1 for any $n \geq 5$.

Definition. Two valuations f_1 and f_2 of a dense near polygon \mathcal{S} are called *neighboring valuations* if there exists an $\epsilon \in \mathbb{Z}$ such that $|f_1(x) - f_2(x) + \epsilon| \leq 1$ for every point x of \mathcal{S} . If this condition holds, then we necessarily have $\epsilon \in \{-1, 0, 1\}$, see Proposition 2.6.

We will also prove the following.

Theorem 1.2 *Regard \mathbb{G}_n , $n \geq 2$, as a subgeometry of $DH(2n - 1, 4)$ which is isometrically embedded into $DH(2n - 1, 4)$. Let f_1 and f_2 be two distinct valuations of \mathbb{G}_n and let x_i , $i \in \{1, 2\}$, denote the unique point of $DH(2n - 1, 4)$ such that the valuation f_i of \mathbb{G}_n is induced by the classical valuation of $DH(2n - 1, 4)$ with center x_i . Then the following are equivalent:*

- (1) f_1 and f_2 are neighboring valuations of \mathbb{G}_n ;
- (2) x_1 and x_2 are collinear.

2 (Semi-)Valuations

2.1 Semi-valuations of general point-line geometries

Throughout this subsection, we suppose that $\mathcal{S} = (\mathcal{P}, \mathcal{L}, I)$ is a connected partial linear space.

Definitions. (1) A semi-valuation of \mathcal{S} is a map $f : \mathcal{P} \rightarrow \mathbb{Z}$ such that for every line L of \mathcal{S} , there exists a unique point x_L on L such that $f(x) = f(x_L) + 1$ for every point x of L distinct from x_L .

(2) It is possible to define an equivalence relation on the set of all semi-valuations of \mathcal{S} : two semi-valuations f_1, f_2 of \mathcal{S} are called *equivalent* if there exists an $\epsilon \in \mathbb{Z}$ such that $f_2(x) = f_1(x) + \epsilon$ for every point x of \mathcal{S} . The equivalence class containing the semi-valuation f of \mathcal{S} will be denoted by $[f]$.

(3) A hyperplane of \mathcal{S} is a proper subspace meeting each line of \mathcal{S} . If f is a semi-valuation of \mathcal{S} attaining a maximal value, then the set of points of \mathcal{S} with non-maximal f -value is a hyperplane H_f of \mathcal{S} . If f_1 and f_2 are two equivalent semi-valuations of \mathcal{S} attaining a maximal value, then $H_{f_1} = H_{f_2}$.

(4) Two semi-valuations f_1 and f_2 of \mathcal{S} are called *neighboring semi-valuations* if there exists an $\epsilon \in \mathbb{Z}$ such that $|f_1(x) - f_2(x) + \epsilon| \leq 1$ for every point x of \mathcal{S} .

Lemma 2.1 *Suppose f_1 and f_2 are two neighboring semi-valuations of \mathcal{S} and let $\epsilon \in \mathbb{Z}$ such that $|f_1(x) - f_2(x) + \epsilon| \leq 1$ for every point x of \mathcal{S} . Then the following holds:*

- (1) *If the set $\{f_1(x) \mid x \in \mathcal{P}\}$ has a minimal element m_1 , then the set $\{f_2(x) \mid x \in \mathcal{P}\}$ has a minimal element m_2 and $|m_1 - m_2 + \epsilon| \leq 1$.*

(2) If the set $\{f_1(x) \mid x \in \mathcal{P}\}$ has a maximal element M_1 , then the set $\{f_2(x) \mid x \in \mathcal{P}\}$ has a maximal element M_2 and $|M_1 - M_2 + \epsilon| \leq 1$.

(3) If L is a line of \mathcal{S} such that the unique point x_1 of L with smallest f_1 -value is distinct from the unique point x_2 of L with smallest f_2 -value, then $\epsilon = f_2(x_2) - f_1(x_1)$.

Proof. Clearly, $f_1(x) + \epsilon - 1 \leq f_2(x) \leq f_1(x) + \epsilon + 1$ for every point x of \mathcal{S} . So, if the set $\{f_1(x) \mid x \in \mathcal{P}\}$ has a minimal (respectively maximal) element, then also the set $\{f_2(x) \mid x \in \mathcal{P}\}$ has a minimal (respectively maximal) element.

(1) If $m_1 - m_2 + \epsilon \leq -2$, then for every point x with f_1 -value m_1 , we have $f_1(x) - f_2(x) + \epsilon = m_1 - f_2(x) + \epsilon \leq m_1 - m_2 + \epsilon \leq -2$, a contradiction. If $m_1 - m_2 + \epsilon \geq 2$, then for every point x with f_2 -value m_2 , we have $f_1(x) - f_2(x) + \epsilon = f_1(x) - m_2 + \epsilon \geq m_1 - m_2 + \epsilon \geq 2$, a contradiction. Hence, $|m_1 - m_2 + \epsilon| \leq 1$.

(2) If $M_1 - M_2 + \epsilon \geq 2$, then for every point x with f_1 -value M_1 , we have $f_1(x) - f_2(x) + \epsilon = M_1 - f_2(x) + \epsilon \geq M_1 - M_2 + \epsilon \geq 2$, a contradiction. If $M_1 - M_2 + \epsilon \leq -2$, then for every point x with f_2 -value M_2 , we have $f_1(x) - f_2(x) + \epsilon = f_1(x) - M_2 + \epsilon \leq M_1 - M_2 + \epsilon \leq -2$, a contradiction. Hence, $|M_1 - M_2 + \epsilon| \leq 1$.

(3) Since $f_1(x_1) - f_2(x_1) = f_1(x_1) - f_2(x_2) - 1$ and $f_1(x_2) - f_2(x_2) = f_1(x_1) - f_2(x_2) + 1$, we necessarily have that $\epsilon = f_2(x_2) - f_1(x_1)$. ■

Lemma 2.2 *Let f_1 and f_2 be two semi-valuations of \mathcal{S} satisfying the following property:*

(*) *For every line L of \mathcal{S} , the unique point of L with smallest f_1 -value coincides with the unique point of L with smallest f_2 -value.*

Then f_1 and f_2 are equivalent.

Proof. Let x^* be an arbitrary point of \mathcal{S} and put $\epsilon := f_2(x^*) - f_1(x^*)$. We prove by induction on the distance $d(x^*, x)$ that $f_2(x) = f_1(x) + \epsilon$ for every point x of \mathcal{S} . Obviously, this holds if $x = x^*$. So, suppose $d(x^*, x) \geq 1$ and let y be a point collinear with x at distance $d(x^*, x) - 1$ from x^* . By the induction hypothesis, $f_2(y) = f_1(y) + \epsilon$. Applying property (*) to the line xy , we find that $f_2(x) = f_1(x) + \epsilon$. ■

The following is an immediate corollary of Lemma 2.1(3) and Lemma 2.2.

Corollary 2.3 *The following holds for two neighboring semi-valuations f_1 and f_2 of \mathcal{S} .*

(1) *If f_1 and f_2 are equivalent, then there exist precisely three $\epsilon \in \mathbb{Z}$ such that $|f_1(x) - f_2(x) + \epsilon| \leq 1$ for every point x of \mathcal{S} . These three possible values of ϵ are consecutive integers.*

(2) *Suppose f_1 and f_2 are not equivalent. Then there exists a unique $\epsilon \in \mathbb{Z}$ such that $|f_1(x) - f_2(x) + \epsilon| \leq 1$ for every point x of \mathcal{S} . There also exists a line L of \mathcal{S} such that the unique point x_1 of L with smallest f_1 -value is distinct from the unique point x_2 of L with smallest f_2 -value. Moreover, $\epsilon = f_2(x_2) - f_1(x_1)$.*

For the remainder of this subsection, we suppose that every line of $\mathcal{S} = (\mathcal{P}, \mathcal{L}, I)$ is incident with precisely 3 points.

Definition. Suppose $f_1 : \mathcal{P} \rightarrow \mathbb{Z}$ and $f_2 : \mathcal{P} \rightarrow \mathbb{Z}$ are two maps such that $|f_1(x) - f_2(x)| \leq 1$ for every point $x \in \mathcal{P}$. If $f_1(x) = f_2(x)$, then we define $f_1 \diamond f_2(x) := f_1(x) - 1 = f_2(x) - 1$. If $|f_1(x) - f_2(x)| = 1$, then we define $f_1 \diamond f_2(x) := \max\{f_1(x), f_2(x)\}$. Clearly, $f_2 \diamond f_1 = f_1 \diamond f_2$. Notice also that $|f_1(x) - f_1 \diamond f_2(x)|, |f_2(x) - f_1 \diamond f_2(x)| \leq 1$ for every point x of \mathcal{S} . Moreover $(f_1 \diamond f_2) \diamond f_1 = f_2$ and $(f_1 \diamond f_2) \diamond f_2 = f_1$.

Proposition 2.4 *If f_1 and f_2 are two semi-valuations of \mathcal{S} such that $|f_1(u) - f_2(u)| \leq 1$ for every point u of \mathcal{S} , then also $f_3 := f_1 \diamond f_2$ is a semi-valuation of \mathcal{S} . If two semi-valuations of the set $\{f_1, f_2, f_3\}$ are equivalent, then all of them are equivalent. If this occurs, then two of them, say f_{i_1} and f_{i_2} , are equal and the third one f_{i_3} satisfies $f_{i_3}(x) = f_{i_1}(x) - 1 = f_{i_2}(x) - 1$ for every point x of \mathcal{S} .*

Proof. Let $L = \{x, y, z\}$ be an arbitrary line of \mathcal{S} . Without loss of generality, we may suppose that one of the following cases occurs:

(1) x is the unique point of L with smallest f_1 -value and smallest f_2 -value. If $f_1(x) = f_2(x)$, then $f_3(x) = f_1(x) - 1$ and $f_3(y) = f_3(z) = f_1(x)$. If $f_1(x) \neq f_2(x)$, then $f_3(x) = \max\{f_1(x), f_2(x)\}$ and $f_3(y) = f_3(z) = \max\{f_1(x) + 1, f_2(x) + 1\} = f_3(x) + 1$.

(2) x is the unique point of L with smallest f_1 -value and y is the unique point of L with smallest f_2 -value. The fact that $|f_1(u) - f_2(u)| \leq 1$ for every $u \in L$ implies that $f_1(x) = f_2(y)$. Since $f_2(x) = f_2(y) + 1 = f_1(x) + 1$, we have $f_3(x) = f_1(x) + 1$. Since $f_1(y) = f_1(x) + 1$ and $f_2(y) = f_1(x)$, we have $f_3(y) = f_1(x) + 1$. Since $f_1(z) = f_1(x) + 1$ and $f_2(z) = f_2(y) + 1 = f_1(x) + 1$, we have $f_3(z) = f_1(x)$.

In both cases, L contains a unique point with smallest f_3 -value. So, f_3 is a semi-valuation. From the definition of the map $f_1 \diamond f_2$, it follows that if f_1 and f_2 are equivalent, then $f_3 = f_1 \diamond f_2$ is equivalent with f_1 and f_2 . So, if f_1 and f_3 are equivalent, then $f_3 \diamond f_1 = (f_1 \diamond f_2) \diamond f_1 = f_2$ is equivalent with f_1 and f_3 , and if f_2 and f_3 are equivalent, then $f_3 \diamond f_2 = (f_1 \diamond f_2) \diamond f_2 = f_1$ is equivalent with f_2 and f_3 . ■

Definition. Suppose f_1 and f_2 are two neighboring semi-valuations of \mathcal{S} . Then we define $[f_1] * [f_2] := [g_1 \diamond g_2]$ where $g_1 \in [f_1]$ and $g_2 \in [f_2]$ are chosen such that $|g_1(x) - g_2(x)| \leq 1$ for every point x of \mathcal{S} . Using Corollary 2.3, it is straightforward to verify that $[g_1 \diamond g_2]$ is independent from the chosen $g_1 \in [f_1]$ and $g_2 \in [f_2]$ satisfying $|g_1(x) - g_2(x)| \leq 1, \forall x \in \mathcal{P}$. Notice also that f_1, f_2 and $g_1 \diamond g_2$ are three mutually neighboring semi-valuations of \mathcal{S} . For every semi-valuation f of \mathcal{S} , we have $[f] * [f] = [f]$.

Notice that if H_1 and H_2 are two distinct hyperplanes of \mathcal{S} , then the complement of the symmetric difference of H_1 and H_2 is again a hyperplane of \mathcal{S} .

Proposition 2.5 *Suppose f_1, f_2 and f_3 are three mutually neighboring semi-valuations of \mathcal{S} such that $[f_3] = [f_1] * [f_2]$. Suppose also that at least one (and hence all) of f_1, f_2, f_3 attains a maximal value. Then precisely one of the following cases occurs:*

- (1) $H_{f_1} \neq H_{f_2}$ and H_{f_3} is the complement of the symmetric difference $H_{f_1} \Delta H_{f_2}$ of H_{f_1} and H_{f_2} .
- (2) One of H_{f_1}, H_{f_2} is properly contained in the other, and H_{f_3} is the larger of the two.
- (3) H_{f_3} is (properly or improperly) contained in $H_{f_1} = H_{f_2}$.

Proof. Without loss of generality, we may suppose that $|f_1(x) - f_2(x)| \leq 1$ for every point x of \mathcal{S} and $f_3 = f_1 \diamond f_2$. Let M_i , $i \in \{1, 2, 3\}$, denote the maximal value attained by f_i . By Lemma 2.1(2), $|M_1 - M_2| \leq 1$. Without loss of generality, we may suppose that $M_2 \geq M_1$.

(a) Suppose that $M_1 = M_2$. If $x \in H_{f_1} \cap H_{f_2}$, then since $f_1(x), f_2(x) \leq M_1 - 1$, we have $f_3(x) \leq M_1 - 1$. If $x \in H_{f_1} \setminus H_{f_2}$, then since $f_1(x) \leq M_1 - 1$ and $f_2(x) = M_1$, we have $f_1(x) = M_1 - 1$ and $f_3(x) = M_1$. Similarly, if $x \in H_{f_2} \setminus H_{f_1}$, then $f_3(x) = M_1$. Finally, if $x \notin H_{f_1} \cup H_{f_2}$, then since $f_1(x) = f_2(x) = M_1$, we have $f_3(x) = M_1 - 1$. If $H_{f_1} \neq H_{f_2}$, then $M_3 = M_1$ and H_{f_3} is the complement of the symmetric difference of H_{f_1} and H_{f_2} . If $H_{f_1} = H_{f_2}$, then $M_3 = M_1 - 1$ and H_{f_3} is contained in $H_{f_1} = H_{f_2}$.

(b) Suppose that $M_2 = M_1 + 1$. Then $H_{f_1} \subseteq H_{f_2}$ since every point of H_{f_1} has f_1 -value at most $M_1 - 1$ and hence f_2 -value at most $M_1 < M_2$. If $x \in H_{f_2}$, then since $f_1(x), f_2(x) \leq M_1$, we have $f_3(x) \leq M_1$. If $x \notin H_{f_2}$, then since $f_1(x) = M_1$ and $f_2(x) = M_2 = M_1 + 1$, we have $f_3(x) = M_1 + 1$. So, $M_3 = M_1 + 1$ and $H_{f_3} = H_{f_2}$. If $H_{f_1} \neq H_{f_2}$, then case (2) of the proposition occurs. If $H_{f_1} = H_{f_2}$, then case (3) occurs. ■

2.2 Valuations of dense near polygons

In this section, we suppose that $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \mathbf{I})$ is a dense near $2n$ -gon. Since every valuation of \mathcal{S} is also a semi-valuation, the definitions and results of Section 2.1 also apply to valuations of \mathcal{S} .

Proposition 2.6 *If f_1 and f_2 are two neighboring valuations of \mathcal{S} and if $\epsilon \in \mathbb{Z}$ such that $|f_1(x) - f_2(x) + \epsilon| \leq 1$ for every point x of \mathcal{S} , then $\epsilon \in \{-1, 0, 1\}$.*

Proof. This is a special case of Lemma 2.1(1). ■

Proposition 2.7 *If f_1 and f_2 are two valuations of \mathcal{S} , then $f_1 = f_2$ if and only if $H_{f_1} = H_{f_2}$.*

Proof. Obviously, $H_{f_1} = H_{f_2}$ if $f_1 = f_2$. We will now also prove that $f_1 = f_2$ if $H_{f_1} = H_{f_2}$.

Let $i \in \{1, 2\}$. Let M_i denote the maximal value attained by f_i . Then the complement $\overline{H_{f_i}}$ of H_{f_i} consists of those points of \mathcal{S} with f_i -value M_i . By Property (V2), $d(x, \overline{H_{f_i}}) \geq M_i - f_i(x)$ for every point x of \mathcal{S} (consider a shortest path between x and $\overline{H_{f_i}}$). We will now prove by induction on $M_i - f_i(x)$ that $d(x, \overline{H_{f_i}}) = M_i - f_i(x)$ for every point x of \mathcal{S} . Obviously, this holds if $M_i - f_i(x) = 0$ since $x \in \overline{H_{f_i}}$ in this case. So, suppose that $M_i - f_i(x) > 0$. Let F_x denote the convex subspace through x as mentioned in Property (V3). Then $f_i(y) \leq f_i(x) \leq M_i - 1$ for every point y of F_x . So, $F_x \neq \mathcal{S}$ and there exists a line L through x not contained in F_x . By Property (V3), L contains a point x' with f_i -value $f_i(x) + 1$. By the induction hypothesis, $d(x', \overline{H_{f_i}}) = M_i - f_i(x') = M_i - f_i(x) - 1$. Hence, $d(x, \overline{H_{f_i}}) \leq M_i - f_i(x)$. Together with $d(x, \overline{H_{f_i}}) \geq M_i - f_i(x)$, this implies that $d(x, \overline{H_{f_i}}) = M_i - f_i(x)$.

Now, suppose $H_{f_1} = H_{f_2}$. Then $M_1 = \max\{d(y, \overline{H_{f_1}}) \mid y \in \mathcal{P}\} = \max\{d(y, \overline{H_{f_2}}) \mid y \in \mathcal{P}\} = M_2$ and $f_1(x) = M_1 - d(x, \overline{H_{f_1}}) = M_2 - d(x, \overline{H_{f_2}}) = f_2(x)$ for every point x of \mathcal{S} . ■

The proof of the following proposition is straightforward.

Proposition 2.8 *Let F be a subspace of \mathcal{S} , isometrically embedded in \mathcal{S} , such that \widetilde{F} is a dense near polygon. Let f_1 and f_2 be two neighboring valuations of \mathcal{S} and let $f'_i, i \in \{1, 2\}$, denote the valuation of \widetilde{F} induced by f_i . Then f'_1 and f'_2 are neighboring valuations of \widetilde{F} .*

Definitions. (1) If F is a convex subspace of \mathcal{S} , then for every point x of \mathcal{S} satisfying $d(x, F) \leq 1$, there exists a unique point in F nearest to x . We will denote this point by $\pi_F(x)$. By Theorem 1.5 of [11], if $d(x, F) \leq 1$, then $d(x, y) = d(x, \pi_F(x)) + d(\pi_F(x), y)$ for every point $y \in F$.

(2) Two convex subspaces F_1 and F_2 of \mathcal{S} are called *parallel* if for every $i \in \{1, 2\}$ and every point $x \in F_i$, there exists a unique point $x' \in F_{3-i}$ at distance $d(F_1, F_2)$ from x and $d(x, y) = d(x, x') + d(x', y) = d(F_1, F_2) + d(x', y)$ for every point y of F_{3-i} . The following proposition is precisely Theorem 1.10 of De Bruyn [11].

Proposition 2.9 *Let F_1 and F_2 be two parallel convex subspaces of \mathcal{S} . Then the map $\pi_{i,3-i} : F_i \rightarrow F_{3-i}, i \in \{1, 2\}$, which maps a point x of F_i to the unique point of F_{3-i} nearest to x , is an isomorphism from \widetilde{F}_i to \widetilde{F}_{3-i} . Moreover, $\pi_{2,1} = \pi_{1,2}^{-1}$.*

Proposition 2.10 *Let f be a valuation of \mathcal{S} , let F_1 and F_2 be two parallel convex subspaces at distance 1 from each other, and let $f_i, i \in \{1, 2\}$, denote the valuation of \widetilde{F}_i induced by f . For every point x of F_1 , put $f'_1(x) := f_2(\pi_{F_2}(x))$. Then f_1 and f'_1 are neighboring valuations of \widetilde{F}_1 .*

Proof. Observe first that f'_1 is a valuation of \widetilde{F}_1 by Proposition 2.9. Let $\delta_i, i \in \{1, 2\}$, be the unique element of \mathbb{N} such that $f(x) = f_i(x) + \delta_i$ for every $x \in F_i$. For every point x of F_1 , we have $|f_1(x) - f'_1(x) + \delta_1 - \delta_2| = |f(x) - f_2(\pi_{F_2}(x)) - \delta_2| = |f(x) - f(\pi_{F_2}(x))| \leq 1$. So, f_1 and f'_1 are neighboring valuations of \widetilde{F}_1 . ■

Definition. (1) Let O be an *ovoid* of \mathcal{S} , i.e. a set of points of \mathcal{S} intersecting each line of \mathcal{S} in a singleton. For a point x of \mathcal{S} , define $f(x) := 0$ if $x \in O$ and $f(x) := 1$ if $x \notin O$. Then f is a so-called *ovoidal valuation* of \mathcal{S} .

(2) Let $\delta \in \{0, \dots, n-1\}$, let x be a point of \mathcal{S} and let O be a set of points of \mathcal{S} at distance at least $\delta + 2$ from x such that every line at distance at least $\delta + 1$ from x has a unique point in common with O . For a point y of \mathcal{S} , we define

$$\begin{cases} f(y) := d(x, y) & \text{if } d(x, y) \leq \delta + 1; \\ f(y) := \delta + 1 & \text{if } d(x, y) \geq \delta + 2 \text{ and } y \notin O; \\ f(y) := \delta & \text{if } d(x, y) \geq \delta + 2 \text{ and } y \in O. \end{cases}$$

By [18, Section 3.1] or [11, Section 5.6.1], f is a (so-called *hybrid*) valuation of \mathcal{S} . We denote f also by $f_{x,\delta,O}$. If $\delta = 0$, then f is an ovoidal valuation of \mathcal{S} with associated ovoid $O \cup \{x\}$. If $\delta = n-1$, then f is a classical valuation of \mathcal{S} . If $\delta = n-2$, then f is called a *semi-classical* valuation of \mathcal{S} .

Proposition 2.11 *Let $\delta \in \{0, \dots, n-1\}$, let L be a line of \mathcal{S} , let x_1 and x_2 be two (not necessarily distinct) points of L and let O_i , $i \in \{1, 2\}$, be a set of points of \mathcal{S} at distance at least $\delta + 2$ from x_i such that every line at distance at least $\delta + 1$ from x_i has a unique point in common with O_i . Then $f_1 := f_{x_1, \delta, O_1}$ and $f_2 := f_{x_2, \delta, O_2}$ are neighboring valuations of \mathcal{S} .*

Proof. Let y be an arbitrary point of \mathcal{S} .

If $d(y, L) \leq \delta$, then $d(x_1, y), d(x_2, y) \leq \delta + 1$ and $|f_1(y) - f_2(y)| = |d(x_1, y) - d(x_2, y)| \leq d(x_1, x_2) \leq 1$ by the triangle inequality.

Suppose $d(y, L) \geq \delta + 1$. Then $d(y, x_1), d(y, x_2) \geq \delta + 1$. It follows that $f_1(y), f_2(y) \in \{\delta, \delta + 1\}$ and $|f_1(y) - f_2(y)| \leq 1$. ■

In the following corollary, we collect two special cases of Proposition 2.11.

Corollary 2.12 (1) *Every two ovoidal valuations of \mathcal{S} are neighboring valuations.*

(2) *If f_1 and f_2 are two classical valuations whose centers lie at distance at most 1 from each other, then f_1 and f_2 are neighboring valuations.*

Definition. Suppose that every line of \mathcal{S} is incident with precisely three points. If f_1 and f_2 are two neighboring valuations of \mathcal{S} , then we denote by $f_1 * f_2$ the unique element of $[f_1] * [f_2]$ whose minimal value is equal to 0. By Proposition 2.4, we know that $f_1 * f_2$ is a semi-valuation of \mathcal{S} .

Proposition 2.13 *Suppose every line of \mathcal{S} is incident with precisely three points. Let F_1 and F_2 be two parallel convex subspaces at distance 1 from each other and let F_3 denote the set of all points of \mathcal{S} not contained in $F_1 \cup F_2$ which are contained in a line joining a point of F_1 with a point of F_2 . Suppose moreover that F_3 is also a convex subspace of \mathcal{S} . Let f be a valuation of \mathcal{S} and let f_i , $i \in \{1, 2, 3\}$, denote the valuation of \widetilde{F}_i induced by f . For every point x of F_1 , we define $f'_1(x) = f_2(\pi_{F_2}(x))$ and $f''_1(x) = f_3(\pi_{F_3}(x))$. Then $f''_1 = f_1 * f'_1$.*

Proof. Notice first that f_1 and f'_1 are neighboring valuations of \widetilde{F}_1 by Proposition 2.10. For every point x of F_1 , we put $g_1(x) := f(x)$, $g_2(x) := f(\pi_{F_2}(x))$ and $g_3(x) := f(\pi_{F_3}(x))$. Then g_1 , g_2 and g_3 are semi-valuations of \widetilde{F}_1 . Since every line meeting F_1 , F_2 and F_3 contains a unique point with smallest f -value (recall (V2)), we necessarily have $g_3 = g_1 \diamond g_2$. It follows that $f''_1 = f_1 * f'_1$. ■

Proposition 2.14 *Suppose that every line of \mathcal{S} is incident with precisely three points. If f_1 and f_2 are distinct neighboring valuations of \mathcal{S} , then $H_{f_1 * f_2}$ is the complement of the symmetric difference of H_{f_1} and H_{f_2} .*

Proof. By Proposition 2.7, $H_{f_1} \neq H_{f_2}$. By Blok and Brouwer [1, Theorem 7.3] or Shult [26, Lemma 6.1], every hyperplane of a dense near polygon is also a maximal subspace. In particular, H_{f_1} , H_{f_2} and $H_{f_1 * f_2}$ are maximal subspaces of \mathcal{S} . It is now clear that case (1)

of Proposition 2.5 must occur. So, $H_{f_1 * f_2}$ is the complement of the symmetric difference of H_{f_1} and H_{f_2} . ■

Suppose again that every line of \mathcal{S} is incident with precisely three points. If f_1 and f_2 are distinct neighboring valuations of \mathcal{S} , then $f_1 * f_2$ satisfies properties (V1) and (V2) in the definition of valuation. The following question can now be considered: does $f_1 * f_2$ also satisfy Property (V3)? If this is the case, then $f_1 * f_2$ is a valuation of \mathcal{S} . We will demonstrate below that the claim that $f_1 * f_2$ is a valuation is false in general, but true for a large class of dense near polygons. We will construct counter examples with the aid of the following lemma. Recall that by Corollary 2.12(1) any two ovoidal valuations of a given dense near polygon are neighboring valuations.

Lemma 2.15 *Suppose every line of \mathcal{S} is incident with precisely three points and that f_1 and f_2 are two distinct ovoidal valuations of \mathcal{S} for which $|H_{f_1} \cap H_{f_2}| \geq 2$ (so, $n \geq 3$). If $f_1 * f_2$ is a valuation of \mathcal{S} , then $f_1 * f_2$ is neither classical nor ovoidal.*

Proof. Since H_{f_1} and H_{f_2} are two distinct maximal subspaces of \mathcal{S} , $H_{f_1} \setminus H_{f_2} \neq \emptyset \neq H_{f_2} \setminus H_{f_1}$. So, $H_{f_1} \Delta H_{f_2} \neq \emptyset$.

Put $f_3 := f_1 \diamond f_2$. If $x \in H_{f_1} \cap H_{f_2}$, then $f_3(x) = -1$. If $x \in H_{f_1} \Delta H_{f_2}$, then $f_3(x) = 1$. If $x \notin H_{f_1} \cup H_{f_2}$, then $f_3(x) = 0$. So, $f_1 * f_2(x)$ is equal to 0 if $x \in H_{f_1} \cap H_{f_2}$, equal to 2 if $x \in H_{f_1} \Delta H_{f_2}$ and equal to 1 if $x \notin H_{f_1} \cup H_{f_2}$. Since $|H_{f_1} \cap H_{f_2}| \geq 2$, $f_1 * f_2$ is not a classical valuation of \mathcal{S} . Since $f_1 * f_2$ can take the value 2, it cannot be an ovoidal valuation of \mathcal{S} . ■

We will now apply Lemma 2.15 to two particular cases.

Example 1. By Brouwer [2], there exists up to isomorphism a unique dense near hexagon \mathcal{S} which satisfies the following properties: (1) every line of \mathcal{S} is incident with precisely 3 points; (2) every point of \mathcal{S} is incident with precisely 12 lines; (3) every quad of \mathcal{S} is a (3×3) -grid. This near hexagon is related to the extended ternary Golay code, see Shult and Yanushka [27, p. 30]. Using the notation of [11] we will denote this near hexagon by \mathbb{E}_1 . The ovoids of the near hexagon \mathbb{E}_1 have been classified in De Bruyn [9, Theorem 4.2]. There are 36 distinct ovoids (all of size 243) and any two distinct ovoids intersect in either 0 or 81 points. The valuations of the near hexagon \mathbb{E}_1 have been classified in De Bruyn and Vandecasteele [20]. Every valuation of \mathbb{E}_1 is either classical or ovoidal. Now, suppose f_1 and f_2 are two ovoidal valuations of \mathbb{E}_1 for which $|H_{f_1} \cap H_{f_2}| = 81$. Then Lemma 2.15 implies that $f_1 * f_2$ is not a valuation of \mathbb{E}_1 . So, the map $f_1 * f_2$ satisfies properties (V1) and (V2), but not (V3). Such maps (for \mathbb{E}_1) were already constructed in De Bruyn [14, Section 4.1].

Example 2. By Brouwer [3], there exists up to isomorphism a unique dense near hexagon \mathcal{S} which satisfies the following properties: (1) every line of \mathcal{S} is incident with precisely 3 points; (2) every point of \mathcal{S} is incident with precisely 15 lines; (3) every quad of \mathcal{S} is isomorphic to the symplectic generalized quadrangle $W(2)$. This near hexagon is related to the Steiner system $S(5, 8, 24)$, see Shult and Yanushka [27, p. 40]. Using the notation of [11] we will denote this near hexagon by \mathbb{E}_2 . The ovoids of the near hexagon \mathbb{E}_2 have

been classified by Brouwer and Lambeck [5, p. 105], see also De Bruyn [11, Section 6.6.2] for an alternative proof. There are 24 distinct ovoids (all of size 253) and any two distinct ovoids intersect in precisely 77 points. The valuations of the near hexagon \mathbb{E}_2 have been classified in De Bruyn and Vandecasteele [20]. Every valuation of \mathbb{E}_2 is either classical or ovoidal. Now, suppose f_1 and f_2 are two distinct ovoidal valuations of \mathbb{E}_2 . Then Lemma 2.15 implies that $f_1 * f_2$ is not a valuation of \mathbb{E}_2 . So, the map $f_1 * f_2$ satisfies properties (V1) and (V2), but not (V3). Such maps (for \mathbb{E}_2) were already constructed in De Bruyn [14, Section 4.2].

The above two examples allow us to draw the following conclusion.

If f_1 and f_2 are two distinct neighboring valuations of a general dense near polygon \mathcal{S} with three points per line, then $f_1 * f_2$ is not necessarily a valuation of \mathcal{S} .

Definition. For every point x of \mathcal{S} , the following point-line geometry $\mathcal{L}(\mathcal{S}, x)$ can be defined. The points of $\mathcal{L}(\mathcal{S}, x)$ are the lines of \mathcal{S} through x , the lines of $\mathcal{L}(\mathcal{S}, x)$ are the quads of \mathcal{S} through x , and incidence is containment. The point-line geometry $\mathcal{L}(\mathcal{S}, x)$ is a linear space and is called the *local space at x* . If F is a convex subspace through x , then the set of all lines of F through x is a subspace of $\mathcal{L}(\mathcal{S}, x)$. The local space $\mathcal{L}(\mathcal{S}, x)$ is called *regular* if every subspace of $\mathcal{L}(\mathcal{S}, x)$ arises from a convex subspace through x in the above-described way.

In De Bruyn [14, Theorem 1.3 + Corollary 1.4], we proved the following:

Proposition 2.16 (1) *If $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \mathbf{I})$ is a dense near polygon, every local space of which is regular, then every map $f : \mathcal{P} \rightarrow \mathbb{N}$ which satisfies properties (V1) and (V2) also satisfies property (V3).*

(2) *If \mathcal{S} is a thick dual polar space, then every local space of \mathcal{S} is regular.*

(3) *If \mathcal{S} is a known dense near polygon without hexes isomorphic to \mathbb{E}_1 or \mathbb{E}_2 , then every local space of \mathcal{S} is regular.*

By Propositions 2.4 and 2.16, we have

Corollary 2.17 *Let \mathcal{S} be a dense near polygon with three points on each line, every local space of which is regular. If f_1 and f_2 are two neighboring valuations of \mathcal{S} , then $f_1 * f_2$ is also a valuation of \mathcal{S} . In particular, this holds if \mathcal{S} is a known dense near hexagon with three points on each line which does not contain hexes isomorphic to \mathbb{E}_1 or \mathbb{E}_2 .*

The following special case of Corollary 2.17 will be of importance in this paper. (The regularity of the local spaces of \mathbb{G}_n was demonstrated in [14, Section 3 (IV)]; also, no hex of \mathbb{G}_n is isomorphic to \mathbb{E}_1 or \mathbb{E}_2 , see [11, Section 6.3.2]).

Corollary 2.18 *If f_1 and f_2 are two neighboring valuations of the near polygon \mathbb{G}_n , $n \geq 2$, then also $f_1 * f_2$ is a valuation of \mathbb{G}_n .*

3 Projective embeddings

3.1 Embeddings of general point-line geometries

A *full (projective) embedding* of a partial linear space $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \mathbb{I})$ into a projective space Σ is an injective mapping e from \mathcal{P} to the point-set of Σ satisfying: (i) $\langle e(\mathcal{P}) \rangle_\Sigma = \Sigma$; (ii) $e(L) := \{e(x) \mid x \in L\}$ is a line of Σ for every line L of \mathcal{S} . The dimensions $\dim(\Sigma)$ and $\dim(\Sigma) + 1$ are respectively called the *projective dimension* and the *vector dimension* of e . If $e : \mathcal{S} \rightarrow \Sigma$ is a full embedding of \mathcal{S} into the projective space Σ , then for every hyperplane α of Σ , $H(\alpha) := e^{-1}(\alpha \cap e(\mathcal{P}))$ is a hyperplane of \mathcal{S} . We say that the hyperplane $H(\alpha)$ of \mathcal{S} *arises from the embedding* e . If H is a hyperplane of \mathcal{S} which is also a maximal subspace of \mathcal{S} (as it is always the case if \mathcal{S} is a dense near polygon), then $\langle e(H) \rangle_\Sigma$ is either Σ or a hyperplane of Σ . Moreover, if $\langle e(H) \rangle_\Sigma$ is a hyperplane of Σ , then $H = e^{-1}(\langle e(H) \rangle_\Sigma \cap e(\mathcal{P}))$, i.e. H arises from the embedding e .

Two full embeddings $e_1 : \mathcal{S} \rightarrow \Sigma_1$ and $e_2 : \mathcal{S} \rightarrow \Sigma_2$ of \mathcal{S} are called *isomorphic* ($e_1 \cong e_2$) if there exists an isomorphism $f : \Sigma_1 \rightarrow \Sigma_2$ such that $e_2 = f \circ e_1$. If $e : \mathcal{S} \rightarrow \Sigma$ is a full embedding of \mathcal{S} and if U is a subspace of Σ satisfying (C1): $\langle U, e(p) \rangle_\Sigma \neq U$ for every point p of \mathcal{S} , (C2): $\langle U, e(p_1) \rangle_\Sigma \neq \langle U, e(p_2) \rangle_\Sigma$ for any two distinct points p_1 and p_2 of \mathcal{S} , then there exists a full embedding e/U of \mathcal{S} into the quotient space Σ/U mapping each point p of \mathcal{S} to $\langle U, e(p) \rangle_\Sigma$. If $e_1 : \mathcal{S} \rightarrow \Sigma_1$ and $e_2 : \mathcal{S} \rightarrow \Sigma_2$ are two full embeddings of \mathcal{S} , then we say that $e_1 \geq e_2$ if there exists a subspace U in Σ_1 satisfying (C1), (C2) and $e_1/U \cong e_2$. If $e : \mathcal{S} \rightarrow \Sigma$ is a full embedding of \mathcal{S} , then by Ronan [25], there exists (up to isomorphism) a unique full embedding $\tilde{e} : \mathcal{S} \rightarrow \tilde{\Sigma}$ satisfying (i) $\tilde{e} \geq e$, (ii) if $e' \geq e$ for some embedding e' of \mathcal{S} , then $\tilde{e} \geq e'$. We say that \tilde{e} is *universal relative to* e . If $\tilde{e} \cong e$ for some full embedding e of \mathcal{S} , then we say that e is *relatively universal*. A full embedding e of \mathcal{S} is called *absolutely universal* if it is universal relative to any full embedding of \mathcal{S} defined over the same division ring as e .

Suppose $\mathcal{S} = (\mathcal{P}, \mathcal{L}, \mathbb{I})$ is a fully embeddable point-line geometry with three points on each line. Then by Ronan [25], \mathcal{S} admits the absolutely universal embedding and every hyperplane of \mathcal{S} arises from this embedding. We now give a description of the absolutely universal embedding of \mathcal{S} . Let V be a vector space over the field \mathbb{F}_2 with a basis B whose vectors are indexed by the elements of \mathcal{P} , e.g. $B = \{\bar{v}_p \mid p \in \mathcal{P}\}$. Let W denote the subspace of V generated by all vectors $\bar{v}_{p_1} + \bar{v}_{p_2} + \bar{v}_{p_3}$ where $\{p_1, p_2, p_3\}$ is a line of \mathcal{S} . Then the map $p \in \mathcal{P} \mapsto \{\bar{v}_p + W, W\}$ defines a full embedding of \mathcal{S} into the projective space $\text{PG}(V/W)$ which is isomorphic to the absolutely universal embedding of \mathcal{S} .

3.2 The Grassmann embedding of the Hermitian dual polar space $DH(2n - 1, \mathbb{K}'/\mathbb{K})$

Let $n \geq 2$, let \mathbb{K}' be a field with involutory automorphism ψ and let \mathbb{K} denote the fixed field of ψ . Then \mathbb{K}' can be regarded as a two-dimensional vector space over \mathbb{K} . Let V be a $2n$ -dimensional vector space over \mathbb{K}' equipped with a nondegenerate skew- ψ -Hermitian form f_V of maximal Witt index n . With the pair (V, f_V) , there is associated a Hermitian

dual polar space $DH(2n - 1, \mathbb{K}'/\mathbb{K})$.

For every point $p = \langle \bar{f}_1, \bar{f}_2, \dots, \bar{f}_n \rangle$ of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$, let $e_1(p)$ denote the point $\langle \bar{f}_1 \wedge \bar{f}_2 \wedge \dots \wedge \bar{f}_n \rangle$ of $\text{PG}(\wedge^n V)$. By Cooperstein [8] and De Bruyn [13], there exists a (necessarily unique) Baer- \mathbb{K} -subgeometry Σ of $\text{PG}(\wedge^n V)$ containing the image of e_1 . Moreover, e_1 defines a full embedding of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ into Σ . This embedding is called the *Grassmann embedding* of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$. By results of Cooperstein [8], De Bruyn & Pasini [17], Kasikova & Shult [22] and Tits [28], we know that the Grassmann embedding of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ is absolutely universal if $n = 2$ or $|\mathbb{K}'| > 4$. The same conclusion cannot be drawn in the case $n \geq 3$, $\mathbb{K} \cong \mathbb{F}_2$ and $\mathbb{K}' \cong \mathbb{F}_4$. Li [23] proved that the absolutely universal embedding of $DH(2n - 1, 2)$ has vector dimension $\frac{4^n+2}{3}$ (which is bigger than $\binom{2n}{n}$ if $n \geq 3$).

Now, let B be a set of $\binom{2n}{n}$ vectors of $\wedge^n V$ such that $\Sigma = \text{PG}(W)$, where W is the $\binom{2n}{n}$ -dimensional vector space over \mathbb{K} whose vectors consist of all \mathbb{K} -linear combinations of the elements of B . By De Bruyn [15, Section 4], there exists a nondegenerate bilinear form f_W on W satisfying the following properties:

(1) f_W is symplectic (or alternating) if either n is odd or $\text{char}(\mathbb{K}) = 2$ and orthogonal if n is even and $\text{char}(\mathbb{K}) \neq 2$.

(2) If ζ is the polarity of $\Sigma = \text{PG}(W)$ associated with f_W , then for every point x of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$, $e_1(x)^\zeta = \langle e_1(H_x) \rangle_\Sigma$, where H_x is the hyperplane of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ consisting of all points of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$ at distance at most $n - 1$ from x .

Lemma 3.1 *Let $\{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n\}$ be a set of n linearly independent vectors of V . Let A denote the set of all vectors $\bar{v} \in V$ for which $\bar{v}_1 \wedge \bar{v}_2 \wedge \dots \wedge \bar{v}_n \wedge \bar{v} = 0$. Then $A = \langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_n \rangle$. As a consequence, A has dimension n .*

Proof. Clearly, $\bar{v}_1 \wedge \bar{v}_2 \wedge \dots \wedge \bar{v}_n \wedge \bar{v} = 0$ if and only if $\{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n, \bar{v}\}$ is linearly dependent, i.e. if and only if $\bar{v} \in \langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_n \rangle$. ■

Lemma 3.2 *Let $\{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n\}$ and $\{\bar{v}'_1, \bar{v}'_2, \dots, \bar{v}'_n\}$ be two sets of n linearly independent vectors of V such that $\langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_n \rangle \neq \langle \bar{v}'_1, \bar{v}'_2, \dots, \bar{v}'_n \rangle$. Let $\delta \in \mathbb{K}' \setminus \{0\}$ and put $\chi := \bar{v}_1 \wedge \bar{v}_2 \wedge \dots \wedge \bar{v}_n + \delta \cdot \bar{v}'_1 \wedge \bar{v}'_2 \wedge \dots \wedge \bar{v}'_n$. Let A denote the set of all $\bar{v} \in V$ for which $\chi \wedge \bar{v} = 0$. Then A is an n -dimensional subspace of V if and only if the subspace $I := \langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_n \rangle \cap \langle \bar{v}'_1, \bar{v}'_2, \dots, \bar{v}'_n \rangle$ has dimension $n - 1$. Moreover, if $\dim(I) = n - 1$, then $\chi = \bar{v}''_1 \wedge \bar{v}''_2 \wedge \dots \wedge \bar{v}''_n$ where $\bar{v}''_1, \bar{v}''_2, \dots, \bar{v}''_n$ are n linearly independent vectors of V such that $\langle \bar{v}''_1, \bar{v}''_2, \dots, \bar{v}''_n \rangle$ is an n -dimensional subspace of V through I distinct from $\langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_n \rangle$ and $\langle \bar{v}'_1, \bar{v}'_2, \dots, \bar{v}'_n \rangle$.*

Proof. Put $k := \dim(I)$. Without loss of generality, we may suppose that $\bar{v}_i = \bar{v}'_i$ for every $i \in \{1, \dots, k\}$. Extend $\{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n\}$ to a basis $\{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_{2n}\}$ of V such that $\bar{v}_{n+i} = \bar{v}'_{k+i}$ for every $i \in \{1, \dots, n - k\}$. Let $\lambda_1, \lambda_2, \dots, \lambda_{2n} \in \mathbb{K}$. Then $\chi \wedge (\lambda_1 \bar{v}_1 + \lambda_2 \bar{v}_2 + \dots + \lambda_{2n} \bar{v}_{2n})$ is equal to

$$\begin{aligned} & (\lambda_{n+1} \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \dots \wedge \bar{v}_n \wedge \bar{v}_{n+1}) + (\lambda_{n+2} \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \dots \wedge \bar{v}_n \wedge \bar{v}_{n+2}) + \dots + \\ & (\lambda_{2n} \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \dots \wedge \bar{v}_n \wedge \bar{v}_{2n}) + ((-1)^{n-k} \lambda_{k+1} \delta \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \dots \wedge \bar{v}_k \wedge \bar{v}_{k+1} \wedge \bar{v}_{n+1} \wedge \bar{v}_{n+2} \wedge \dots \end{aligned}$$

$$\begin{aligned}
& \wedge \bar{v}_{2n-k}) + ((-1)^{n-k} \lambda_{k+2} \delta \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \cdots \wedge \bar{v}_k \wedge \bar{v}_{k+2} \wedge \bar{v}_{n+1} \wedge \bar{v}_{n+2} \wedge \cdots \wedge \bar{v}_{2n-k}) \\
& + \cdots + ((-1)^{n-k} \lambda_n \delta \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \cdots \wedge \bar{v}_k \wedge \bar{v}_n \wedge \bar{v}_{n+1} \wedge \bar{v}_{n+2} \wedge \cdots \wedge \bar{v}_{2n-k}) \\
& + (\lambda_{2n-k+1} \delta \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \cdots \wedge \bar{v}_k \wedge \bar{v}_{n+1} \wedge \bar{v}_{n+2} \wedge \cdots \wedge \bar{v}_{2n-k} \wedge \bar{v}_{2n-k+1}) \\
& + (\lambda_{2n-k+2} \delta \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \cdots \wedge \bar{v}_k \wedge \bar{v}_{n+1} \wedge \bar{v}_{n+2} \wedge \cdots \wedge \bar{v}_{2n-k} \wedge \bar{v}_{2n-k+2}) \\
& + \cdots + (\lambda_{2n} \delta \cdot \bar{v}_1 \wedge \bar{v}_2 \wedge \cdots \wedge \bar{v}_k \wedge \bar{v}_{n+1} \wedge \bar{v}_{n+2} \wedge \cdots \wedge \bar{v}_{2n-k} \wedge \bar{v}_{2n}).
\end{aligned}$$

If $k \leq n - 2$, then the $2n$ vectors of the form $\bar{v}_{i_1} \wedge \bar{v}_{i_2} \wedge \cdots \wedge \bar{v}_{i_{n+1}}$ occurring in the above sum are distinct and linearly independent. So, in this case $\chi \wedge (\lambda_1 \bar{v}_1 + \cdots + \lambda_{2n} \bar{v}_{2n}) = 0$ if and only if $\lambda_{k+1} = \lambda_{k+2} = \cdots = \lambda_{2n} = 0$. It follows that $\dim(A) = k < n$.

If $k = n - 1$, then $\chi = \bar{v}_1 \wedge \bar{v}_2 \wedge \cdots \wedge \bar{v}_{n-1} \wedge (\bar{v}_n + \delta \bar{v}'_n)$. By Lemma 3.1, it then follows that $\dim(A) = n$. Notice also that $\langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_{n-1}, \bar{v}_n + \delta \bar{v}'_n \rangle$ is an n -dimensional subspace of V through $I = \langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_{n-1} \rangle$ distinct from $\langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_{n-1}, \bar{v}_n \rangle$ and $\langle \bar{v}_1, \bar{v}_2, \dots, \bar{v}_{n-1}, \bar{v}'_n \rangle = \langle \bar{v}'_1, \bar{v}'_2, \dots, \bar{v}'_{n-1}, \bar{v}'_n \rangle$. ■

The following corollary to Lemmas 3.1 and 3.2 will be useful later.

Corollary 3.3 *Any line of Σ containing at least three points of the image of e_1 is of the form $e_1(L)$ for some line L of $DH(2n - 1, \mathbb{K}'/\mathbb{K})$.*

3.3 Embeddings of the dense near $2n$ -gon \mathbb{G}_n

Let $n \geq 2$. In Section 1, we mentioned that there exists a subspace X of $DH(2n - 1, 4)$ satisfying: (i) $\tilde{X} \cong \mathbb{G}_n$; (ii) if $x_1, x_2 \in X$, then the distance between x_1 and x_2 in the geometry \tilde{X} is equal to the distance between x_1 and x_2 in the dual polar space $DH(2n - 1, 4)$. It can be proved, see De Bruyn [16], that there exists up to isomorphism a unique set of points of $DH(2n - 1, 4)$ satisfying (i) and (ii).

Since X is a subspace in $DH(2n - 1, 4)$, the Grassmann embedding $e_1 : DH(2n - 1, 4) \rightarrow \Sigma$ of $DH(2n - 1, 4)$ will induce an embedding e_2 of $\tilde{X} \cong \mathbb{G}_n$ into a subspace Σ' of Σ . In De Bruyn [12], we proved that $\Sigma' = \Sigma$ and that e_2 is the absolutely universal embedding of $\tilde{X} \cong \mathbb{G}_n$. The latter implies (recall Ronan [25]) that every hyperplane of \tilde{X} arises from the embedding e_2 . Since every hyperplane of $\tilde{X} \cong \mathbb{G}_n$ is also a maximal subspace of \tilde{X} , we can say more: if H is a hyperplane of \tilde{X} , then $\Pi = \langle e_2(H) \rangle_\Sigma$ is a hyperplane of Σ and $H = e_2^{-1}(e_2(X) \cap \Pi)$. The embedding e_1 is not absolutely universal if $n \geq 3$ ¹. However, since every hyperplane of $DH(2n - 1, 4)$ is a maximal subspace of $DH(2n - 1, 4)$, a similar property as above holds: if H is a hyperplane of $DH(2n - 1, 4)$ arising from e_1 , then $\Pi = \langle e_1(H) \rangle_\Sigma$ is a hyperplane of Σ and $H = e_1^{-1}(e_1(\mathcal{P}) \cap \Pi)$, where \mathcal{P} denotes the point-set of $DH(2n - 1, 4)$.

Let ζ denote the polarity of Σ as defined in Section 3.2. Recall that for every point $x \in \mathcal{P}$, H_x denotes the hyperplane of $DH(2n - 1, 4)$ consisting of all points of $DH(2n - 1, 4)$

¹The reader might be puzzled by the fact that e_2 is absolutely universal, while e_1 is not. This happens because, when you lift e_1 to the absolutely universal embedding \tilde{e}_1 of $DH(2n - 1, 4)$, the image of \mathbb{G}_n lifts to a set of points that spans a complement of the kernel of the projection of \tilde{e}_1 onto e_1 .

at distance at most $n - 1$ from x . For every point x of $DH(2n - 1, 4)$, we have $\langle e_1(H_x) \rangle_\Sigma = e_1(x)^\zeta$.

Lemma 3.4 *Let x be a point of $DH(2n - 1, 4)$ and let f denote the valuation of $\tilde{X} \cong \mathbb{G}_n$ induced by the classical valuation of $DH(2n - 1, 4)$ with center x . Then $\langle e_2(H_f) \rangle_\Sigma = \langle e_1(H_x) \rangle_\Sigma$. Hence, $e_1(x) = \langle e_1(H_x) \rangle_\Sigma^\zeta = \langle e_2(H_f) \rangle_\Sigma^\zeta$.*

Proof. Since both $\langle e_2(H_f) \rangle_\Sigma = \langle e_1(H_f) \rangle_\Sigma$ and $\langle e_1(H_x) \rangle_\Sigma = e_1(x)^\zeta$ are hyperplanes of Σ and $H_f \subseteq H_x$, we necessarily have $\langle e_2(H_f) \rangle_\Sigma = \langle e_1(H_x) \rangle_\Sigma$. ■

The last claim of Lemma 3.4 says that the point x is uniquely determined by the hyperplane H_f of \mathbb{G}_n . So, we have:

Corollary 3.5 *For every valuation f of $\tilde{X} \cong \mathbb{G}_n$, there exists at most one point x of $DH(2n - 1, 4)$ such that f is induced by the classical valuation of $DH(2n - 1, 4)$ with center x .*

Lemma 3.6 *Let f_1 and f_2 be two distinct neighboring valuations of $\tilde{X} \cong \mathbb{G}_n$ and let f_3 be the valuation $f_1 * f_2$ of \tilde{X} . Suppose that for every $i \in \{1, 2, 3\}$, there exists a (necessarily unique) point x_i of $DH(2n - 1, 4)$ such that the valuation f_i of \tilde{X} is induced by the classical valuation of $DH(2n - 1, 4)$ with center x_i . Then $\{x_1, x_2, x_3\}$ is a line of $DH(2n - 1, 4)$.*

Proof. By Proposition 2.14, H_{f_3} is the complement of the symmetric difference of H_{f_1} and H_{f_2} . This implies that $\langle e_2(H_{f_1}) \rangle_\Sigma$, $\langle e_2(H_{f_2}) \rangle_\Sigma$ and $\langle e_2(H_{f_3}) \rangle_\Sigma$ are the three hyperplanes of Σ through a given subspace of Σ of co-dimension 2. It follows that $e_1(x_1) = \langle e_2(H_{f_1}) \rangle_\Sigma^\zeta$, $e_1(x_2) = \langle e_2(H_{f_2}) \rangle_\Sigma^\zeta$ and $e_1(x_3) = \langle e_2(H_{f_3}) \rangle_\Sigma^\zeta$ determine a line of Σ . By Corollary 3.3, $\{x_1, x_2, x_3\}$ is a line of $DH(2n - 1, 4)$. ■

4 Several useful lemmas

A max M of a dense near polygon \mathcal{S} is called *big* if every point of \mathcal{S} has distance at most 1 from M . If M is a big max of \mathcal{S} , then by Theorem 2.30 of [11], every quad of \mathcal{S} which meets M is either contained in M or intersects M in a line.

If M_1 and M_2 are two disjoint big maxes of a dense near polygon \mathcal{S} , then M_1 and M_2 are parallel convex subspaces at distance 1 from each other. Proposition 2.9 tells us that there exist a natural isomorphism between \widetilde{M}_1 and \widetilde{M}_2 . If F is a convex subspace of diameter δ of M_1 , then $\langle F, \pi_{M_2}(F) \rangle$ is a convex subspace of diameter $\delta + 1$ of \mathcal{S} .

Suppose \mathcal{S} is a dense near polygon with three points on each line and that M is a big max of \mathcal{S} . For every point x of M , we define $\mathcal{R}_M(x) := x$. For every point x of \mathcal{S} not contained in M , let $\mathcal{R}_M(x)$ denote the unique point of the line $x\pi_M(x)$ distinct from x and $\pi_M(x)$. By Theorem 1.11 of [11], \mathcal{R}_M is an automorphism of \mathcal{S} . So, if M' is a (big) max of \mathcal{S} , then $\mathcal{R}_M(M')$ is also a (big) max of \mathcal{S} .

Every max of the dual polar space $DH(2n - 1, 4)$, $n \geq 2$, is big. If F is a convex subspace of the dual polar space $DH(2n - 1, 4)$, $n \geq 2$, then for every point x of $DH(2n -$

1, 4), there exists a unique point $\pi_F(x) \in F$ nearest to x . Moreover, $d(x, y) = d(x, \pi_F(x)) + d(\pi_F(x), y)$ for every $y \in F$. If F has diameter $\delta \in \{2, \dots, n\}$, then $\tilde{F} \cong DH(2\delta - 1, 4)$.

Let V be a $2n$ -dimensional vector space ($n \geq 2$) with basis B . We will now collect several properties of the near polygon $\mathbb{G}_n := \mathbb{G}_n(V, B)$. We refer to [11, Section 6.3] for proofs.

If \bar{x} is a vector of weight 2 of V , then the set of all points of \mathbb{G}_n which, regarded as n -dimensional subspaces of V , contain the vector \bar{x} is a big max of \mathbb{G}_n . In the sequel, we will say that M is the big max of \mathbb{G}_n corresponding to \bar{x} . If $n \geq 3$, then every big max of \mathbb{G}_n arises from a vector of weight 2 of V . If M is a big max of \mathbb{G}_n , $n \geq 3$, then $\overline{M} \cong \mathbb{G}_{n-1}$. Suppose M is a big max of \mathbb{G}_n corresponding to a vector \bar{x} of weight 2 of V . The set of points of $DH(V, B) \cong DH(2n - 1, 4)$ which, regarded as n -dimensional subspaces of V , contain the vector \bar{x} is a max \overline{M} of $DH(V, B)$. \overline{M} is the unique max of $DH(V, B)$ containing M .

Let \bar{x}_1 and \bar{x}_2 be two linearly independent vectors of weight 2 of V and let M_i , $i \in \{1, 2\}$, denote the big max of \mathbb{G}_n corresponding to \bar{x}_i . If \bar{x}_1 and \bar{x}_2 have disjoint supports, then M_1 and M_2 meet. If the supports of \bar{x}_1 and \bar{x}_2 are not disjoint, then M_1 and M_2 are disjoint.

Suppose the supports of \bar{x}_1 and \bar{x}_2 are not disjoint. Then the two-space $\langle \bar{x}_1, \bar{x}_2 \rangle$ contains a unique vector \bar{x}_3 of weight 2 distinct from \bar{x}_1 and \bar{x}_2 , and we denote by M_3 the big max of \mathbb{G}_n corresponding to \bar{x}_3 . We have $M_3 = \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$. If \overline{M}_i , $i \in \{1, 2, 3\}$, denotes the unique max of $DH(V, B)$ containing M_i , then $\overline{M}_3 = \mathcal{R}_{\overline{M}_1}(\overline{M}_2) = \mathcal{R}_{\overline{M}_2}(\overline{M}_1)$. So, every line meeting M_1 (\overline{M}_1) and M_2 (\overline{M}_2) also meets M_3 (\overline{M}_3). If the supports of \bar{x}_1 and \bar{x}_2 are equal, then every line meeting M_1 , M_2 (and M_3) is special. If the supports of \bar{x}_1 and \bar{x}_2 intersect in a singleton, then every line meeting M_1 , M_2 (and M_3) is an ordinary line.

Every quad of \mathbb{G}_n , $n \geq 3$, is isomorphic to either the (3×3) -grid, the generalized quadrangle $W(2)$ or the generalized quadrangle $Q^-(5, 2)$.

If $n \geq 3$, then the automorphism group of \mathbb{G}_n has two orbits on the set of lines of \mathbb{G}_n , namely the set of ordinary lines and the set of special lines. A line of \mathbb{G}_n , $n \geq 3$, is an ordinary line if and only if it is contained in a $W(2)$ -quad. An ordinary line of \mathbb{G}_n , $n \geq 3$, is contained in a unique $Q^-(5, 2)$ -quad. The automorphism group of \mathbb{G}_n , $n \geq 3$, acts transitively on the set of $W(2)$ -quads of \mathbb{G}_n and the set of $Q^-(5, 2)$ -quads of \mathbb{G}_n . A grid-quad of \mathbb{G}_n , $n \geq 3$, is said to be of *Type I* if it contains a special line, otherwise it is called a *grid-quad of Type II*. Every grid-quad of \mathbb{G}_3 has Type I and the automorphism group of \mathbb{G}_3 acts transitively on the set of its grid-quads. The automorphism group of \mathbb{G}_n , $n \geq 4$, has two orbits on the set of grid-quads of \mathbb{G}_n , namely the set of grid-quads of Type I and the set of grid-quads of Type II.

Every point of \mathbb{G}_n , $n \geq 3$, is contained in precisely n special lines. If L_1, \dots, L_k are $k \geq 2$ special lines through a given point of \mathbb{G}_n , then $\langle L_1, \dots, L_k \rangle \cong \mathbb{G}_k$. Conversely, if F is a convex subspace of \mathbb{G}_n , $n \geq 3$, such that $\tilde{F} \cong \mathbb{G}_k$ for some $k \geq 2$, then through every point of F , there are precisely k special lines of \mathbb{G}_n which are contained in F . If F is a convex subspace of \mathbb{G}_n , $n \geq 3$, such that $\tilde{F} \cong \mathbb{G}_k$ for some $k \geq 3$, then a line contained in F is a special line of F if and only if it is a special line of \mathbb{G}_n .

The following lemma was proved in De Bruyn [11, Section 6.3.3].

Lemma 4.1 *Let Q be a quad of \mathbb{G}_n , $n \geq 3$, containing a special line. Then there are two possibilities:*

(1) *Q is a grid-quad of Type I. Then Q contains precisely three special lines. These three lines partition the point-set of Q .*

(2) *Q is a $Q^-(5, 2)$ -quad of \mathbb{G}_n . Then Q can be partitioned into three subgrids G_1, G_2, G_3 . A line of Q is special if and only if it is contained in one of the grids G_1, G_2 and G_3 .*

Lemma 4.2 (1) *Every grid-quad Q of Type I of \mathbb{G}_n , $n \geq 3$, is contained in a unique hex isomorphic to \mathbb{G}_3 .*

(2) *Every $Q^-(5, 2)$ -quad Q of \mathbb{G}_n , $n \geq 3$, is contained in precisely $n - 2$ hexes isomorphic to \mathbb{G}_3 .*

(3) *Let M_1 and M_2 be two disjoint maxes of \mathbb{G}_n , $n \geq 3$, such that every line meeting M_1 and M_2 is special. Let F be a convex subspace of M_1 such that $\tilde{F} \cong \mathbb{G}_k$ for some $k \geq 2$. Then $\langle F, \pi_{M_2}(F) \rangle \cong \mathbb{G}_{k+1}$.*

Proof. (1) Let x be an arbitrary point of Q , let L_1 denote the unique special line of Q through x and let M denote the unique ordinary line of Q through x . Then M is contained in a unique $Q^-(5, 2)$ -quad R of \mathbb{G}_n . Let L_2 and L_3 denote the unique special lines of R through x . Then $\langle L_1, L_2, L_3 \rangle = \langle Q, R \rangle$ is a \mathbb{G}_3 -hex containing Q . Conversely, if F is a \mathbb{G}_3 -hex through Q , then there exists a $Q^-(5, 2)$ -quad of \tilde{F} containing the line M . This $Q^-(5, 2)$ -quad necessarily coincides with R . So, $F = \langle Q, R \rangle$.

(2) Let x be an arbitrary point of Q , and let L_1 and L_2 be the two special lines of Q through x . If F is a \mathbb{G}_3 -hex through Q , then there exists a unique special line $L_3 \notin \{L_1, L_2\}$ through x contained in F . Conversely, if L_3 is one of the $n - 2$ special lines of \mathbb{G}_n through x distinct from L_1 and L_2 , then $\langle L_1, L_2, L_3 \rangle$ is a \mathbb{G}_3 -hex containing Q . It follows that there are precisely $n - 2$ \mathbb{G}_3 -hexes containing Q .

(3) Recall that since F has diameter δ , the convex subspace $\langle F, \pi_{M_2}(F) \rangle$ has diameter $\delta + 1$. Let x be an arbitrary point of F and let L_1, \dots, L_k denote the k special lines of F through x . Let $\underline{L_{k+1}}$ denote the unique line through x meeting M_2 . Then L_{k+1} is a special line. So, $\langle F, \pi_{M_2}(F) \rangle = \langle F, L_{k+1} \rangle = \langle L_1, L_2, \dots, L_{k+1} \rangle \cong \mathbb{G}_{k+1}$. ■

Lemma 4.1 implies the following.

Corollary 4.3 *Let L_1 and L_2 be two disjoint special lines of \mathbb{G}_n , $n \geq 3$, which are contained in a quad Q , let G denote the unique (3×3) -subgrid of Q containing L_1, L_2 and let L_3 denote the unique line of G disjoint from L_1 and L_2 . Then also L_3 is a special line of \mathbb{G}_n .*

Let \mathcal{S}_n , $n \geq 3$, be the following point-line geometry:

- The points of \mathcal{S}_n are the special lines of \mathbb{G}_n ;
- The lines of \mathcal{S}_n are all the triples $\{L_1, L_2, L_3\}$, where L_1, L_2 and L_3 are three mutually disjoint special lines which are contained in some (3×3) -subgrid of \mathbb{G}_n .

- Incidence is containment.

Lemma 4.4 *The complement of a proper subspace of \mathcal{S}_n , $n \geq 3$, is connected.*

Proof. Let S be a subspace of \mathcal{S}_n and let L_1, L_2 be two distinct special lines contained in the complement of S . We will prove by induction on $d(L_1, L_2)$ that L_1 and L_2 are connected by a path which entirely consists of points of \mathcal{S}_n not contained in S . Here, $d(L_1, L_2)$ denotes the distance between L_1 and L_2 in the near polygon \mathbb{G}_n .

First, suppose that $d(L_1, L_2) = 0$. Then L_1 and L_2 are contained in a unique $Q^-(5, 2)$ -quad Q . By Lemma 4.1(2), there exist special lines L'_1 and L''_1 of Q such that: (i) $\{L_1, L'_1, L''_1\}$ is a line of \mathcal{S}_n ; (ii) the unique (3×3) -subgrid of Q containing L_1, L'_1 and L''_1 does not contain L_2 . Since $L_1 \notin S$, at least one of L'_1, L''_1 does not belong to S . Hence, L_1, L'_1, L_2 or L_1, L''_1, L_2 is a path of \mathcal{S}_n contained in the complement of S .

Suppose now that $d(L_1, L_2) > 0$. Let $x_1 \in L_1$ and $x_2 \in L_2$ be points such that $d(x_1, x_2) = d(L_1, L_2)$. Let M_1 denote a line through x_1 containing a unique point y_1 at distance $d(x_1, x_2) - 1$ from x_2 and let M_2 denote a line of $\langle x_1, x_2 \rangle$ through x_2 which is not contained in $\langle x_2, y_1 \rangle$. Then M_2 contains a unique point y_2 at distance $d(x_1, x_2) - 1$ from x_1 . Let $z_i, i \in \{1, 2\}$, denote the unique point of M_i distinct from x_i and y_i .

Since $d(x_2, y_1) = d(x_1, x_2) - 1$, we have $d(x_2, z_1) = d(x_1, x_2)$. Since the line x_2y_2 is not contained in $\langle x_2, y_1 \rangle$, we have $d(y_2, y_1) = d(y_2, x_2) + d(x_2, y_1) = d(x_1, x_2)$. Together with $d(y_2, x_1) = d(x_1, x_2) - 1$, this implies that $d(y_2, z_1) = d(x_1, x_2)$. Since the line x_2z_2 is not contained in $\langle x_2, y_1 \rangle$, we have $d(z_2, y_1) = d(z_2, x_2) + d(x_2, y_1) = d(x_1, x_2)$. Since $d(x_1, y_2) = d(x_1, x_2) - 1$, we have $d(z_2, x_1) = d(x_1, x_2)$. Finally, since $d(z_2, x_1) = d(z_2, y_1) = d(x_1, x_2)$, we have $d(z_2, z_1) = d(x_1, x_2) - 1$. We can conclude that for every point u_i of $M_i, i \in \{1, 2\}$, there exists a unique point of M_{3-i} at distance $d(x_1, x_2) - 1$ from u_i .

Notice that $L_1 \neq M_1$ and $L_2 \neq M_2$. So, $\langle L_1, M_1 \rangle$ and $\langle L_2, M_2 \rangle$ are quads. We will now define a special line L'_i of $\langle L_i, M_i \rangle$ through y_i disjoint from L_i ($i \in \{1, 2\}$). Since L_1 and L_2 are special lines, we can distinguish two cases by Lemma 4.1.

(i) Suppose $\langle L_i, M_i \rangle$ is a grid-quad of Type I. Then let L'_i denote the unique line of $\langle L_i, M_i \rangle$ through y_i disjoint from L_i . Then L'_i is special.

(ii) Suppose $\langle L_i, M_i \rangle$ is a $Q^-(5, 2)$ -quad. Then there are precisely two special lines of $\langle L_i, M_i \rangle$ through y_i . Let L'_i denote any special line of $\langle L_i, M_i \rangle$ through y_i not meeting L_i .


The lines L_i and L'_i are contained in a unique (3×3) -subgrid of $\langle L_i, M_i \rangle$. We denote by L''_i the unique line of this subgrid which is disjoint from L_i and L'_i . Then also L''_i is special and $z_i \in L''_i$. Since $L_i \notin S$, at most one of L_i, L'_i, L''_i belongs to S . Since $|M_1| = |M_2| = 3$, there exist points $u_1 \in M_1$ and $u_2 \in M_2$ such that (i) $d(u_1, u_2) = d(x_1, x_2) - 1 = d(L_1, L_2) - 1$; (ii) for every $i \in \{1, 2\}$, the unique line $U_i \in \{L_i, L'_i, L''_i\}$ containing u_i does not belong to S .

By the induction hypothesis, U_1 and U_2 are connected by a path which entirely consists of points of \mathcal{S}_n which are contained in the complement of S . Hence, also L_1 and L_2 are connected by a path of \mathcal{S}_n which entirely consists of points of \mathcal{S}_n which are contained in the complement of S . ■

Lemma 4.5 Let M_1 and M_2 be two disjoint big maxes of \mathbb{G}_n , $n \geq 4$, such that every line meeting M_1 and M_2 is special. Let $\mathcal{S}_{n-1}(M_1)$ denote the geometry isomorphic to \mathcal{S}_{n-1} defined on the set of special lines of M_1 (recall $\widetilde{M}_1 \cong \mathbb{G}_{n-1}$). Let f be a semi-valuation of \mathbb{G}_n and let S denote the set of special lines L of M_1 such that the unique points of the lines L and $\pi_{M_2}(L)$ with smallest f -values are collinear. Then S is a subspace of $\mathcal{S}_{n-1}(M_1)$.

Proof. Let $\{K_1, L_1, N_1\}$ be an arbitrary line of $\mathcal{S}_{n-1}(M_1)$ such that $K_1, L_1 \in S$. We need to prove that $N_1 \in S$. Put $K_2 = \pi_{M_2}(K_1)$, $L_2 = \pi_{M_2}(L_1)$ and $N_2 = \pi_{M_2}(N_1)$.

Case I. Suppose the unique point u of K_1 with smallest f -value is collinear with the unique point v of L_1 with smallest f -value. In the following picture we sketch this situation and indicate the values of the points of K_1 and L_1 .




If $\gamma = \delta$, then using the fact that every line meeting K_1 and L_1 contains a unique point with smallest value, we obtain that $f(x) = \gamma - 1$, $f(y) = \gamma$ and $f(z) = \gamma$. So, x is the unique point of N_1 with smallest f -value.

If $\gamma \neq \delta$, then using the fact that every line meeting K_1 and L_1 contains a unique point with smallest value, we obtain $f(x) = \max\{\gamma, \delta\}$ and $f(y) = f(z) = \max\{\gamma + 1, \delta + 1\} = f(x) + 1$. So, again x is the unique point of N_1 with smallest f -value.

Now, since $K_1, L_1 \in S$, $\pi_{M_2}(u)$ is the unique point of K_2 with smallest f -value and $\pi_{M_2}(v)$ is the unique point of L_2 with smallest f -value. Since u and v are collinear, also $\pi_{M_2}(u)$ and $\pi_{M_2}(v)$ are collinear. Repeating the above reasoning for the lines K_2, L_2, N_2 instead of K_1, L_1, N_1 , we find that $\pi_{M_2}(x)$ is the unique point of N_2 with smallest f -value. Since x is collinear with $\pi_{M_2}(x)$, we have $N_1 \in S$ as we needed to prove.

Case II. The unique point u of K_1 with smallest f -value is not collinear with the unique point v of L_1 with smallest f -value. This situation is sketched in the following picture, where the values of the points of K_1 and L_1 are mentioned.



Since the f -values of two collinear points differ by at most 1, we have $|(\gamma + 1) - \delta| \leq 1$ and $|\gamma - (\delta + 1)| \leq 1$. It follows that $\gamma = \delta$. Since every line meeting K_1 and L_1 contains a unique point with smallest f -value, we have $f(x) = \gamma + 1$, $f(y) = \gamma + 1$ and $f(z) = \gamma$. So, z is the unique point of N_1 with smallest f -value.

Now, since $K_1, L_1 \in S$, $\pi_{M_2}(u)$ is the unique point of K_2 with smallest f -value and $\pi_{M_2}(v)$ is the unique point of L_2 with smallest f -value. Since u and v are not collinear, also $\pi_{M_2}(u)$ and $\pi_{M_2}(v)$ are not collinear. Repeating the above reasoning for the lines K_2, L_2, N_2 instead of K_1, L_1, N_1 , we find that $\pi_{M_2}(z)$ is the unique point of N_2 with smallest f -value. Since z is collinear with $\pi_{M_2}(z)$, we have $N_1 \in S$ as we needed to prove.

■

Lemma 4.6 (1) *Let M_1 and M_2 be two disjoint big maxes of \mathbb{G}_n , $n \geq 3$, such that every line meeting M_1 and M_2 is special. Put $M_3 := \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$. Then every quad meeting M_1, M_2 (and M_3) is either a grid-quad of Type I or a $Q^-(5, 2)$ -quad.*

(2) *Every point x of \mathbb{G}_n not contained in $M_1 \cup M_2 \cup M_3$ is contained in a unique quad Q_x which intersect M_1, M_2 (and M_3) in lines. This quad Q_x is a $Q^-(5, 2)$ -quad.*

(3) *Let L be a line of M_1 . Then $\langle L, \pi_{M_2}(L) \rangle$ is a grid-quad of Type I if L is an ordinary line and a $Q^-(5, 2)$ -quad if L is a special line.*

Proof. (1) Suppose Q is a quad meeting M_1 in a line L_1 and M_2 in a line L_2 . Let $x \in L_1$. Since Q contains the points x and $\pi_{M_2}(x) \in L_2$, it contains the special line $x\pi_{M_2}(x)$. Hence, Q is either a grid-quad of Type I or a $Q^-(5, 2)$ -quad by Lemma 4.1.

(2) Suppose x is a point of \mathbb{G}_n not contained in $M_1 \cup M_2 \cup M_3$. If Q is a quad through x meeting M_1 and M_2 in lines, then Q necessarily contains the points $\pi_{M_1}(x)$ and $\pi_{M_2}(x)$. If $x\pi_{M_1}(x) = x\pi_{M_2}(x)$, then $\{x, \pi_{M_1}(x), \pi_{M_2}(x)\}$ is a line meeting M_1 and M_2 , a contradiction, since $x \notin M_3$. Hence, $x\pi_{M_1}(x) \neq x\pi_{M_2}(x)$ and Q necessarily coincides with the quad $Q_x := \langle x\pi_{M_1}(x), x\pi_{M_2}(x) \rangle$. Since Q_x meets M_1 and M_2 in lines it is either a grid-quad or a $Q^-(5, 2)$ -quad by part (1). Since $Q_x \cap (M_1 \cup M_2 \cup M_3)$ is a subgrid of Q_x and $x \notin M_1 \cup M_2 \cup M_3$, Q_x necessarily is a $Q^-(5, 2)$ -quad.

(3) Let $x \in L$ and let L' denote the unique line through x meeting M_2 . Then $\langle L, \pi_{M_2}(L) \rangle = \langle L, L' \rangle$. If L is special, then $\langle L, L' \rangle$ is a $Q^-(5, 2)$ -quad since L and L' are two distinct special lines through x .

Conversely, suppose that $\langle L, L' \rangle$ is a $Q^-(5, 2)$ -quad. There are precisely n special lines through x , two of these special lines are contained in $\langle L, L' \rangle$ and $n - 1$ of these special lines are contained in M_1 (recall $\widetilde{M}_1 \cong \mathbb{G}_{n-1}$). It follows that $L = \langle L, L' \rangle \cap M_1$ is a special line. \blacksquare

Lemma 4.7 *Let M_1 and M_2 be two disjoint (big) maxes of $DH(2n - 1, 4)$, $n \geq 2$, and put $M_3 := \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$. Then every point x of $DH(2n - 1, 4)$ not contained in $M_1 \cup M_2 \cup M_3$ is contained in a unique quad Q_x which intersects M_1 , M_2 (and M_3) in lines.*

Proof. Similarly as in the proof of Lemma 4.6(2), we have that $x\pi_{M_1}(x) \neq x\pi_{M_2}(x)$ and that Q_x is the unique quad of $DH(2n - 1, 4)$ containing the lines $x\pi_{M_1}(x)$ and $x\pi_{M_2}(x)$. \blacksquare

As already mentioned in Section 1, the following lemma was proved in [19, Proposition 7.7] in the case $n = 3$ and in [21, Proposition 6.13] in the case $n = 4$.

Lemma 4.8 *Regard \mathbb{G}_n , $n \in \{3, 4\}$, as a subgeometry of $DH(2n - 1, 4)$ which is isometrically embedded into $DH(2n - 1, 4)$. Then every valuation of \mathbb{G}_n is induced by a unique (classical) valuation of $DH(2n - 1, 4)$.*

Lemma 4.9 *Let M_1 and M_2 be two disjoint big maxes of the near polygon \mathbb{G}_3 such that every line meeting M_1 and M_2 is special. Let f be a valuation of \mathbb{G}_3 having the property that there exists a line K of M_1 such that the unique point of K with smallest f -value is not collinear with the unique point of $\pi_{M_2}(K)$ with smallest f -value. Then there exists a special line L of M_1 such that the unique point of L with smallest f -value is not collinear with the unique point of $\pi_{M_2}(L)$ with smallest f -value.*

Proof. We regard \mathbb{G}_3 as a subgeometry of $DH(5, 4)$ which is isometrically embedded into $DH(5, 4)$. Then by Lemma 4.8, there exists a unique point x of $DH(5, 4)$ such that f is induced by the classical valuation of $DH(5, 4)$ with center x . Put $M_3 := \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$. We have $\widetilde{M}_1 \cong \widetilde{M}_2 \cong \widetilde{M}_3 \cong Q^-(5, 2)$. So, M_1 , M_2 and M_3 are quads of both \mathbb{G}_3 and $DH(5, 4)$.

We prove that $x \notin M_1 \cup M_2 \cup M_3$. Suppose to the contrary that $x \in M_i$ for a certain $i \in \{1, 2, 3\}$. Let u_j , $j \in \{1, 2, 3\}$, denote the unique point of $\pi_{M_j}(K)$ nearest to x . Then u_j is the unique point of $\pi_{M_j}(K)$ with smallest f -value. Since $d(x, y) = d(x, \pi_{M_i}(y)) + d(\pi_{M_i}(y), y)$ for every $j \in \{1, 2, 3\}$ and every point $y \in \pi_{M_j}(K)$, we have $u_j = \pi_{M_j}(u_i)$. So, $\{u_1, u_2, u_3\}$ is a line meeting M_1 , M_2 and M_3 . This contradicts the fact that the unique point of K with smallest f -value is not collinear with the unique point of $\pi_{M_2}(K)$ with smallest f -value.

So, $x \notin M_1 \cup M_2 \cup M_3$. By Lemma 4.7, there exists a unique quad Q_x of $DH(5, 4)$ through x intersecting M_1 , M_2 and M_3 in lines. By Lemma 4.1(2), there exists a special line L in M_1 disjoint from the line $Q_x \cap M_1$. Let x_i , $i \in \{1, 2\}$, denote the unique point of M_i collinear with x and let y_i denote the unique point of $\pi_{M_i}(L)$ collinear with x_i . Since $x \notin M_1 \cup M_2 \cup M_3$, x_1 and x_2 are not collinear. Hence, also y_1 and y_2 are not collinear.

Now, for every $i \in \{1, 2\}$ and every point z of $\pi_{M_i}(L)$, we have $d(x, z) = d(x, x_i) + d(x_i, z)$. So, y_i , $i \in \{1, 2\}$, is the unique point of $\pi_{M_i}(L)$ nearest to x , or equivalently, the unique point of $\pi_{M_i}(L)$ with smallest f -value.

Summarizing, we have that the unique point of the special line L with smallest f -value is not collinear with the unique point of $\pi_{M_2}(L)$ with smallest f -value. ■

Lemma 4.10 *Let f be a semi-valuation of the near polygon \mathbb{G}_n , $n \geq 2$, and let Q be a $Q^-(5, 2)$ -quad of \mathbb{G}_n . Then Q contains a unique point x^* with smallest f -value and $f(x) = f(x^*) + d(x^*, x)$ for every point x of Q .*

Proof. It is easy to show (see e.g. De Bruyn [14, Lemma 2.2]) that every semi-valuation of a thick generalized quadrangle is equivalent to either a classical valuation or an ovoidal valuation. Since the generalized quadrangle $Q^-(5, 2)$ has no ovoids (see e.g. Payne and Thas [24, 3.4.1]), f is equivalent with a classical valuation of $Q^-(5, 2)$. The lemma follows. ■

If K and L are two lines of a near polygon, then by Theorem 1.3 of [11] precisely one of the following two cases occurs: (a) there exists a unique point $k^* \in K$ and a unique point $l^* \in L$ such that $d(k, l) = d(k, k^*) + d(k^*, l^*) + d(l^*, l)$ for every point $k \in K$ and every point $l \in L$; (b) for every point k in K , there exists a unique point $l \in L$ such that $d(k, l) = d(K, L)$. If case (b) occurs, then K and L are parallel.

Lemma 4.11 *Let M_1 and M_2 be two disjoint maxes of the dual polar space $DH(2n-1, 4)$, $n \geq 3$, and let $M_3 = \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$. Let Q and R be two quads of $DH(2n-1, 4)$ which intersect M_1 and M_2 in lines. If x is a point of R such that the unique points of $Q \cap M_1$ and $Q \cap M_2$ nearest to x are not collinear, then*

- (1) $K := Q \cap M_1$ and $L := R \cap M_1$ are parallel lines;
- (2) Q and R are parallel quads;
- (3) $x \in R \setminus (M_1 \cup M_2 \cup M_3)$.

Proof. (1) Suppose to the contrary that K and L are not parallel and let $k^* \in K$ and $l^* \in L$ denote the unique points such that $d(k, l) = d(k, k^*) + d(k^*, l^*) + d(l^*, l)$ for all $k \in K$ and all $l \in L$. Since the map $M_1 \rightarrow M_2; x \mapsto \pi_{M_2}(x)$ is an isomorphism between \widetilde{M}_1 and \widetilde{M}_2 , the lines $\pi_{M_2}(K)$ and $\pi_{M_2}(L)$ are not parallel and $d(k, l) = d(k, \pi_{M_2}(k^*)) + d(\pi_{M_2}(k^*), \pi_{M_2}(l^*)) + d(\pi_{M_2}(l^*), l)$ for all $k \in \pi_{M_2}(K)$ and $l \in \pi_{M_2}(L)$. For every $i \in \{1, 2\}$ and every $y \in Q \cap M_i = \pi_{M_i}(K)$, we have $d(x, y) = d(x, \pi_{M_i}(x)) + d(\pi_{M_i}(x), y) = d(x, \pi_{M_i}(x)) + d(\pi_{M_i}(x), \pi_{M_i}(l^*)) + d(\pi_{M_i}(l^*), \pi_{M_i}(k^*)) + d(\pi_{M_i}(k^*), y)$. So, k^* is the unique point of $K = Q \cap M_1$ nearest to x and $\pi_{M_2}(k^*)$ is the unique point of $\pi_{M_2}(K) = Q \cap M_2$ nearest to x . This contradicts the fact that the unique points of $Q \cap M_1$ and $Q \cap M_2$ nearest to x are not collinear.

(2) By part (1), K and L are parallel. Put $\delta := d(K, L)$. For every point u of R , there exists a unique point $\pi_Q(u) \in Q$ nearest to u and $d(u, v) = d(u, \pi_Q(u)) + d(\pi_Q(u), v)$ for every $v \in Q$. We prove that $\pi_Q(u)$ has distance δ from u . It suffices to prove the following things:

(a) If $u \in R \cap (M_1 \cup M_2 \cup M_3)$, then $\{d(u, v) \mid v \in Q \cap (M_1 \cup M_2 \cup M_3)\} = \{\delta, \delta + 1, \delta + 2\}$.

(b) If $u \in R \setminus (M_1 \cup M_2 \cup M_3)$, then $\{d(u, v) \mid v \in Q \cap (M_1 \cup M_2 \cup M_3)\} = \{\delta + 1, \delta + 2\}$.

Moreover, there is more than one $v \in Q \cap (M_1 \cup M_2 \cup M_3)$ for which $d(u, v) = \delta + 1$.

(a) Suppose $u \in R \cap M_i$ for some $i \in \{1, 2, 3\}$. Let u' denote the unique point of $Q \cap M_i$ nearest to u . Then $d(u, u') = \delta$ and $d(u, v) = \delta + 1$ for every $v \in (Q \cap M_i) \setminus \{u'\}$. Now, let $j \in \{1, 2, 3\} \setminus \{i\}$. Then $d(u, \pi_{M_j}(u')) = d(u, u') + d(u', \pi_{M_j}(u')) = \delta + 1$. If $v \in (Q \cap M_j) \setminus \{u'\}$, then $d(u, \pi_{M_j}(v)) = d(u, v) + d(v, \pi_{M_j}(v)) = \delta + 2$. This proves (a).

(b) Suppose $u \in R \setminus (M_1 \cup M_2 \cup M_3)$. Let $u_i, i \in \{1, 2, 3\}$, denote the unique point of $M_i \cap R$ collinear with u and let u'_i denote the unique point of $M_i \cap Q$ nearest to u . Then $d(u, u'_i) = d(u, u_i) + d(u_i, u'_i) = \delta + 1$ and for every $v \in (Q \cap M_i) \setminus \{u'_i\}$, we have $d(u, v) = d(u, u_i) + d(u_i, v) = \delta + 2$. This proves (b).

Similarly, for every point u of Q , there exists a unique point $\pi_R(u) \in R$ nearest to u and $d(u, v) = d(u, \pi_R(u)) + d(\pi_R(u), v)$ for every $v \in R$. With a similar reasoning as above, one can show that $\pi_R(u)$ has distance δ from u . It follows that Q and R are parallel quads.

(3) Suppose to the contrary that $x \in R \cap M_i$ for some $i \in \{1, 2, 3\}$. Let $x_j, j \in \{1, 2, 3\}$, denote the unique point of $Q \cap M_j$ nearest to x . For every $y \in Q \cap M_i$ and $j \in \{1, 2, 3\}$, we have $d(x, \pi_{M_j}(y)) = d(x, y) + d(y, \pi_{M_j}(y))$. So, $x_j = \pi_{M_j}(x_i)$ for every $j \in \{1, 2, 3\}$. This would imply that x_1 and x_2 are collinear, a contradiction. It follows that $x \in R \setminus (M_1 \cup M_2 \cup M_3)$. \blacksquare

Lemma 4.12 *Regard \mathbb{G}_4 as a subgeometry of $DH(7, 4)$ which is isometrically embedded into $DH(7, 4)$. Let f be a semi-valuation of \mathbb{G}_4 . Let M_1 and M_2 be two disjoint big maxes of \mathbb{G}_4 such that every line meeting M_1 and M_2 is special and let Q be a $Q^-(5, 2)$ -quad of \mathbb{G}_4 which intersect M_1 and M_2 in lines such that the unique points of $Q \cap M_1$ and $Q \cap M_2$ with smallest f -values are not collinear. Then f is uniquely determined by the values that it takes on the set $M_1 \cup M_2 \cup Q$.*

Proof. By Proposition 2.16, the semi-valuation f of \mathbb{G}_4 is equivalent with a unique valuation f' of \mathbb{G}_4 . By Lemma 4.8, the valuation f' of \mathbb{G}_4 is induced by a unique classical valuation f'' of $DH(7, 4)$. It suffices to prove that the center x of f'' is uniquely determined by the values that f takes on the set $M_1 \cup M_2 \cup Q$.

Put $M_3 = \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$ and let $\overline{M}_i, i \in \{1, 2, 3\}$, denote the unique max of $DH(7, 4)$ containing M_i . By Lemma 4.7, there exists a quad Q_x of $DH(7, 4)$ through x intersecting $\overline{M}_1, \overline{M}_2$ and \overline{M}_3 in lines. (Clearly, this is also valid if x would be contained in $\overline{M}_1 \cup \overline{M}_2 \cup \overline{M}_3$.) Now, the unique points of $Q \cap M_1$ and $Q \cap M_2$ with smallest f -value are not collinear, or equivalently, the unique points of $Q \cap M_1$ and $Q \cap M_2$ nearest to x are not collinear. By Lemma 4.11(2), Q and Q_x are parallel quads and $x \in Q_x \setminus (\overline{M}_1 \cup \overline{M}_2 \cup \overline{M}_3)$. Let $x_i, i \in \{1, 2, 3\}$, denote the unique point of \overline{M}_i collinear with x . Then x_1, x_2 and x_3 are mutually noncollinear. Since $d(x, y) = d(x, x_i) + d(x_i, y) = 1 + d(x_i, y)$ for every $i \in \{1, 2, 3\}$ and every $y \in \overline{M}_i$, the valuation of \overline{M}_i induced by f is also induced by the valuation of $\widetilde{\overline{M}_i}$ with center x_i . We also know that the valuation of Q induced by f is

classical (recall Lemma 4.10) and that the center of this classical valuation is the unique point of Q nearest to x .

The above discussion allows us to construct x from the values that f takes on the set $M_1 \cup M_2 \cup Q$. Let $f_i, i \in \{1, 2\}$, denote the valuation of \widetilde{M}_i induced by f . Then by Lemma 4.8, f_i is induced by a unique classical valuation of \widetilde{M}_i . We denote by x_i^* the center of this classical valuation of \widetilde{M}_i . By the above, x_1^* and x_2^* lie at distance 2 from each other. So, they determine a unique quad Q^* which is parallel with Q . If y^* denotes the unique point of Q with smallest f -value, then x necessarily is the unique point of Q^* nearest to y^* . ■

Lemma 4.13 *Regard $\mathbb{G}_n, n \geq 4$, as a subgeometry of $DH(2n-1, 4)$ which is isometrically embedded into $DH(2n-1, 4)$. Let f be a semi-valuation of \mathbb{G}_n . Let M_1 and M_2 be two disjoint big maxes of \mathbb{G}_n such that every line meeting M_1 and M_2 is special, and let Q be a $Q^-(5, 2)$ -quad of \mathbb{G}_n which intersects M_1 and M_2 in lines such that the unique points of $Q \cap M_1$ and $Q \cap M_2$ with smallest f -values are not collinear. Then f is uniquely determined by the values that it takes on the set $M_1 \cup M_2 \cup Q$.*

Proof. Notice first that contrary to the situation in the proof of Lemma 4.12, we do not know (yet) whether the valuation of \mathbb{G}_n which is equivalent with f is induced by a classical valuation of $DH(2n-1, 4)$. Put $M_3 = \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$ and let $\overline{M}_i, i \in \{1, 2, 3\}$, denote the unique max of $DH(2n-1, 4)$ containing M_i . Let x be an arbitrary point of \mathbb{G}_n not contained in $M_1 \cup M_2$.

Suppose first that $x \in M_3$. Then there exists a unique line L through x meeting M_1 in a point x_1 and M_2 in a point x_2 . If $f(x_1) = f(x_2)$, then $f(x) = f(x_1) - 1 = f(x_2) - 1$. If $f(x_1) \neq f(x_2)$, then $f(x) = \max\{f(x_1), f(x_2)\}$. So, $f(x)$ is uniquely determined by the values that f takes on the set $M_1 \cup M_2 \cup Q$.

By the previous paragraph, we may suppose that $x \notin M_1 \cup M_2 \cup M_3$. Then by Lemma 4.6(2) there exists a unique $Q^-(5, 2)$ -quad Q_x through x which intersect M_1, M_2 and M_3 in (special) lines.

Suppose first that the unique point u_1 of $M_1 \cap Q_x$ with smallest f -value is collinear with the unique point u_2 of $M_2 \cap Q_x$ with smallest f -value. Let u denote the point of the line u_1u_2 with smallest f -value. Now, f takes three values on the subgrid $Q_x \cap (M_1 \cup M_2 \cup M_3)$ of Q_x , namely $f(u), f(u) + 1$ and $f(u) + 2$. It follows that the valuation of Q_x induced by f is classical with center u (recall also Lemma 4.10). So, the f -values of the points of Q_x (in particular, of x) are uniquely determined by the values that f takes on the set $M_1 \cup M_2 \cup Q$.

Suppose next that the unique point of $M_1 \cap Q_x$ with smallest f -value is not collinear with the unique point of $M_2 \cap Q_x$ with smallest f -value. Let $\mathcal{S}_{n-1}(M_1)$ denote the geometry isomorphic to \mathcal{S}_{n-1} defined on the set of special lines of \widetilde{M}_1 . Let S denote the set of special lines L of \widetilde{M}_1 such that the unique points of L and $\pi_{M_2}(L)$ with smallest f -values are collinear. Then S is a subspace of $\mathcal{S}_{n-1}(M_1)$ by Lemma 4.5. It is a proper subspace since $Q \cap M_1 \notin S$. So, the complement of S is connected by Lemma 4.4. It follows that there exists a sequence $Q = Q_1, Q_2, \dots, Q_k = Q_x$ of $k \geq 1$ $Q^-(5, 2)$ -quads which intersect

M_1 and M_2 in lines and which satisfy: (1) for every $i \in \{1, \dots, k\}$, $Q_i \cap M_1$ is a special line not belonging to S ; (2) for every $i \in \{1, \dots, k-1\}$, $Q_i \cap M_1$ and $Q_{i+1} \cap M_1$ are collinear points of $\mathcal{S}_{n-1}(M_1)$. It suffices to prove that for every $i \in \{1, \dots, k-1\}$, the values $f(x)$, $x \in Q_{i+1}$, are uniquely determined by the values that f takes on the set $M_1 \cup M_2 \cup Q_i$. By Lemma 4.1 there are two possibilities for $\langle Q_i \cap M_1, Q_{i+1} \cap M_1 \rangle$. Either $\langle Q_i \cap M_1, Q_{i+1} \cap M_1 \rangle$ is a special grid-quad of Type I or a $Q^-(5, 2)$ -quad. In any case, $\langle Q_i \cap M_1, Q_{i+1} \cap M_1 \rangle$ is contained in a \mathbb{G}_3 -hex $F \subseteq M_1$ by Lemma 4.2(1)+(2). The convex sub-octagon $\langle F, \pi_{M_2}(F) \rangle$ contains $Q_i \cup Q_{i+1}$ and is isomorphic to \mathbb{G}_4 by Lemma 4.2(3). By Lemma 4.12, the values $f(x)$, $x \in Q_{i+1}$, are uniquely determined by the values that f takes on the set $F \cup \pi_{M_2}(F) \cup Q_i$ and hence (a fortiori) also by the values that f takes on the set $M_1 \cup M_2 \cup Q_i$. This was precisely what we needed to show. ■

5 Proof of Theorem 1.1

We regard \mathbb{G}_n , $n \geq 2$, as a subgeometry of $DH(2n-1, 4)$ which is isometrically embedded into $DH(2n-1, 4)$. Recall that by De Bruyn [16], there exists up to isomorphism a unique isometric embedding of \mathbb{G}_n into $DH(2n-1, 4)$.

Let f be a valuation of the near polygon \mathbb{G}_n . We will prove by induction on n that f is induced by a unique (classical) valuation of $DH(2n-1, 4)$. This trivially holds if $n = 2$. By Lemma 4.8, this claim also holds if $n \in \{3, 4\}$. So, in the sequel we will suppose that $n \geq 5$. Let M_1 and M_2 be two disjoint big maxes of \mathbb{G}_n such that every line meeting M_1 and M_2 is special. Recall that $\widetilde{M}_1 \cong \widetilde{M}_2 \cong \mathbb{G}_{n-1}$. Put $M_3 := \mathcal{R}_{M_1}(M_2) = \mathcal{R}_{M_2}(M_1)$. Let \widetilde{M}_i , $i \in \{1, 2, 3\}$, denote the unique max of $DH(2n-1, 4)$ containing M_i . Then $\widetilde{M}_3 := \mathcal{R}_{\widetilde{M}_1}(\widetilde{M}_2) = \mathcal{R}_{\widetilde{M}_2}(\widetilde{M}_1)$. By Proposition 2.9, there exists for any two distinct $i, j \in \{1, 2, 3\}$ a natural isomorphism between \widetilde{M}_i and \widetilde{M}_j . This isomorphism induces an isomorphism between \widetilde{M}_i and \widetilde{M}_j . Let f_i , $i \in \{1, 2, 3\}$, denote the valuation of \widetilde{M}_i induced by f . For every point x of M_1 , we define $f'_1(x) := f_2(\pi_{M_2}(x))$ and $f''_1(x) := f_3(\pi_{M_3}(x))$. By Propositions 2.10 and 2.13, f_1 , f'_1 and f''_1 are two by two neighboring valuations of \widetilde{M}_1 and $f''_1 = f_1 * f'_1$. We distinguish two cases.

Case I: f_1 and f'_1 are equal.

In this case, $f_1 = f'_1 = f''_1$. Let x^* denote a point of $M_1 \cup M_2 \cup M_3$ such that $f(x) \geq f(x^*)$ for every point $x \in M_1 \cup M_2 \cup M_3$. Let $i^* \in \{1, 2, 3\}$ such that $x^* \in M_{i^*}$. Considering the unique line through x^* meeting M_1 , M_2 and M_3 , we see that $f(\pi_{M_i}(x^*)) = f(x^*) + 1$ for every $i \in \{1, 2, 3\} \setminus \{i^*\}$. Since $f_1 = f'_1 = f''_1$, we necessarily have that $f(\pi_{M_i}(x)) = f(x) + 1$ for every point x of M_{i^*} and every $i \in \{1, 2, 3\} \setminus \{i^*\}$. For every $i \in \{1, 2, 3\}$ there exists by the induction hypothesis a unique point $x_i^* \in \widetilde{M}_i$ such that the valuation f_i of \widetilde{M}_i is induced by the classical valuation of \widetilde{M}_i with center x_i^* . Taking into account the natural isomorphisms between the near polygons \widetilde{M}_1 , \widetilde{M}_2 and \widetilde{M}_3 , we see that $\{x_1^*, x_2^*, x_3^*\}$ must be a line of $DH(2n-1, 4)$ meeting \widetilde{M}_1 , \widetilde{M}_2 and \widetilde{M}_3 . Put $y^* := x_{i^*}^*$ and let f^* be the

valuation of \mathbb{G}_n induced by the classical valuation of $DH(2n-1, 4)$ with center y^* . Since $d(y^*, x) = d(y^*, \pi_{M_i}(y^*)) + d(\pi_{M_i}(y^*), x) = d(y^*, x_i^*) + d(x_i^*, x)$ for every $i \in \{1, 2, 3\}$ and every point $x \in \overline{M_i}$, the valuation of $\widetilde{M_i}$ induced by f^* is equal to the valuation of $\widetilde{M_i}$ induced by the classical valuation of $\widetilde{M_i}$ with center x_i^* , i.e. is equal to f_i .

Claim. *We prove that if f' is a valuation of \mathbb{G}_n and $\epsilon \in \mathbb{Z}$ such that $f'(x) = f(x) + \epsilon$ for every point $x \in M_1 \cup M_2$, then $\epsilon = 0$ and $f' = f$.*

PROOF. (i) Let x_3 be an arbitrary point of M_3 and let L be the unique line through x_3 intersecting M_1 in a point x_1 and M_2 in a point x_2 . If $f(x_1) = f(x_2)$, then $f(x_3) = f(x_1) - 1 = f(x_2) - 1$. If $f(x_1) \neq f(x_2)$, then $f(x_3) = \max\{f(x_1), f(x_2)\}$. Similarly, if $f'(x_1) = f'(x_2)$, then $f'(x_3) = f'(x_1) - 1 = f'(x_2) - 1$ and if $f'(x_1) \neq f'(x_2)$, then $f'(x_3) = \max\{f'(x_1), f'(x_2)\}$. Since $f'(x_1) = f(x_1) + \epsilon$ and $f'(x_2) = f(x_2) + \epsilon$, we have $f'(x_3) = f(x_3) + \epsilon$.

(ii) Let x be an arbitrary point of \mathbb{G}_n not contained in $M_1 \cup M_2 \cup M_3$. Then by Lemma 4.6(2), there exists a unique $Q^-(5, 2)$ -quad Q_x through x which intersects M_1 , M_2 and M_3 in lines. So, $G := Q_x \cap (M_1 \cup M_2 \cup M_3)$ is a (3×3) -grid. Since $f_1 = f'_1 = f''_1$, the grid G is easily seen to contain a unique point u with smallest f -value. Moreover, $f(v) = f(u) + d(u, v)$ for every point $v \in G$. By (i) we then also know that $f'(v) = f'(u) + d(u, v)$ for every point $v \in G$. Hence, the valuations of Q_x induced by f and f' coincide with the classical valuation of Q_x with center u . This implies that $f'(y) = f(y) + \epsilon$ for every $y \in Q_x$. In particular, $f'(x) = f(x) + \epsilon$.

By (i) and (ii), f and f' differ by a constant ϵ . Since f and f' have minimal value 0, we have $\epsilon = 0$ and $f = f'$. (qed)

Since f_1 is the valuation of $\widetilde{M_1}$ induced by f and also the valuation of $\widetilde{M_1}$ induced by f^* , there exists an $\epsilon \in \mathbb{Z}$ such that $f^*(x) = f(x) + \epsilon$ for every $x \in M_1$.

If $y^* = x_1^*$, then $i^* = 1$ and $f(\pi_{M_2}(x)) = f(x) + 1$ for every $x \in M_1$. Since $d(y^*, \pi_{M_2}(x)) = d(y^*, x) + 1$, we also have $f^*(\pi_{M_2}(x)) = f^*(x) + 1$ for every $x \in M_1$.

If $y^* = x_2^*$, then $i^* = 2$ and $f(\pi_{M_2}(x)) = f(x) - 1$ for every $x \in M_1$. Since $d(y^*, \pi_{M_2}(x)) = d(y^*, x) - 1$, we also have $f^*(\pi_{M_2}(x)) = f^*(x) - 1$ for every $x \in M_1$.

If $y^* = x_3^*$, then $i^* = 3$ and $f(\pi_{M_2}(x)) = f(x)$ for every $x \in M_1$. Since $d(y^*, \pi_{M_2}(x)) = d(y^*, x)$, we also have $f^*(\pi_{M_2}(x)) = f^*(x)$ for every $x \in M_1$.

It follows that $f^*(x) = f(x) + \epsilon$ for every $x \in M_1 \cup M_2$. By the above Claim we then have that $f^* = f$. So, f is induced by a classical valuation of $DH(2n-1, 4)$. Corollary 3.5 then implies that f is induced by a unique classical valuation of $DH(2n-1, 4)$.

Case II: f_1 and f'_1 are not equal.

By the induction hypothesis, there exists for every $i \in \{1, 2, 3\}$ a unique point $x_i \in \overline{M_i}$ such that the valuation f_i of $\widetilde{M_i}$ is induced by the classical valuation of $\widetilde{M_i}$ with center x_i . Since the map $\overline{M_j} \rightarrow \overline{M_1}; x \mapsto \pi_{\overline{M_1}}(x)$, $j \in \{2, 3\}$, is an isomorphism between $\widetilde{M_j}$ and $\widetilde{M_1}$, the valuation f'_1 of $\widetilde{M_1}$ is induced by the classical valuation of $\widetilde{M_1}$ with center $\pi_{\overline{M_1}}(x_2)$ and

the valuation f''_1 of \widetilde{M}_1 is induced by the classical valuation of \widetilde{M}_1 with center $\pi_{\overline{M}_1}(x_3)$. By Lemma 3.6, $L := \{x_1, \pi_{\overline{M}_1}(x_2), \pi_{\overline{M}_1}(x_3)\}$ is a line of \overline{M}_1 . Now, let R denote the unique quad of $DH(2n - 1, 4)$ containing L and $\pi_{\overline{M}_2}(L)$. Then $G_R := R \cap (\overline{M}_1 \cup \overline{M}_2 \cup \overline{M}_3)$ is a (3×3) -subgrid of R and $\{x_1, x_2, x_3\}$ is an ovoid of G_R .

Since $f_1 \neq f'_1$, there exists by Corollary 2.3(2) a line K of M_1 such that the unique point of K with smallest f_1 -value is distinct from the unique point of K with smallest f'_1 -value, or equivalently, such that the unique point u_1 of K with smallest f -value is not collinear with the unique point u_2 of $\pi_{M_2}(K)$ with smallest f -value. Here, $u_i, i \in \{1, 2\}$, is the unique point of $\pi_{M_i}(K)$ nearest to x_i . Let u_3 denote the unique point of $\pi_{M_3}(K)$ nearest to x_3 .

Now, consider an arbitrary $Q^-(5, 2)$ -quad T of M_1 through the line K . By Lemma 4.2(3), $\langle T, \pi_{M_2}(T) \rangle$ is a \mathbb{G}_3 -hex. Applying Lemma 4.9 to the near polygon $\langle T, \pi_{M_2}(T) \rangle \cong \mathbb{G}_3$, the big maxes T and $\pi_{M_2}(T)$ of $\langle T, \pi_{M_2}(T) \rangle$ and the valuation of $\langle T, \pi_{M_2}(T) \rangle$ induced by f , we see that we may without loss of generality suppose that the line K which we introduced in the previous paragraph is a special line of \mathbb{G}_n .

Let Q be the quad $\langle K, \pi_{M_2}(K) \rangle$. Since K is a special line, Q is a $Q^-(5, 2)$ -quad of both \mathbb{G}_n and $DH(2n - 1, 4)$ (recall Lemma 4.6(3)). Let y be one of the three points of $R \setminus G_R$ such that $\Gamma_1(y) \cap G_R = \{x_1, x_2, x_3\}$. Then $d(y, x) = d(y, x_1) + d(x_1, x) = 1 + d(x_1, x)$ for every point x of \overline{M}_1 . It follows that u_1 is the unique point of K nearest to y . In a similar way, one proves that u_2 is the unique point of $\pi_{M_2}(K)$ nearest to y . Since u_1 and u_2 are not collinear, Lemma 4.11 tells us that K and L are parallel lines and that Q and R are parallel quads.

We claim that $u_i, i \in \{1, 2, 3\}$, is the unique point of Q nearest to x_i . Suppose that this would not be the case. Then $\pi_Q(x_i) \notin M_i$. But then the unique point of $Q \cap M_i$ collinear with $\pi_Q(x_i)$ would lie closer to x_i than $\pi_Q(x_i)$ itself, clearly a contradiction.

Since $u_1 \neq u_2$, f can take two distinct values on the (3×3) -subgrid $G_Q := Q \cap (M_1 \cup M_2 \cup M_3)$ of Q . The points of G_Q with smallest f -value form the ovoid $\{u_1, u_2, u_3\}$ of G_Q . So, the unique point u^* of Q with smallest f -value (recall Lemma 4.10) is collinear with u_1, u_2 and u_3 . Now, let x^* denote the unique point of R nearest to u^* . Since $x_i, i \in \{1, 2, 3\}$, is the unique point of R nearest to u_i , the point x^* is one of the three points of $R \setminus G_R$ collinear with x_1, x_2 and x_3 . Now, let f^* denote the valuation of \mathbb{G}_n induced by the classical valuation of $DH(2n - 1, 4)$ with center x^* and let $\epsilon \in \mathbb{Z}$ be such that $f^*(u^*) + \epsilon = f(u^*)$. We prove that $f^*(x) + \epsilon = f(x)$ for every point x of $M_1 \cup M_2 \cup Q$.

Since u^* is the unique point of Q nearest to x^* , we have $f^*(x) + \epsilon = f^*(u^*) + \epsilon + d(u^*, x) = f(u^*) + d(u^*, x) = f(x)$ for every point x of Q .

Let $i \in \{1, 2\}$. Since $d(x^*, y) = d(x^*, x_i) + d(x_i, y)$ for every $y \in \overline{M}_i$, the valuation of \widetilde{M}_i induced by f^* coincides with the valuation of \widetilde{M}_i induced by the classical valuation of \overline{M}_i with center x_i , i.e. with the valuation f_i of \widetilde{M}_i induced by f . It follows that $f(x) - f^*(x)$ is independent from the point $x \in M_i$. By the previous paragraph, $f(x) - f^*(x) = f(u_i) - f^*(u_i) = \epsilon$.

By the two previous paragraphs, $f^*(x) + \epsilon = f(x)$ for every point x of $M_1 \cup M_2 \cup Q$. Lemma 4.13 then implies that $f^*(x) + \epsilon = f(x)$ for every point x of \mathbb{G}_n . Since the minimal

values attained by f and f^* are equal to 0, we have $\epsilon = 0$ and $f = f^*$. So, f is induced by the classical valuation of $DH(2n - 1, 4)$ with center x^* . By Corollary 3.5, f is induced by a unique classical valuation of $DH(2n - 1, 4)$.

6 Proof of Theorem 1.2

We devote this short section to the proof of Theorem 1.2.

We regard \mathbb{G}_n , $n \geq 2$, as a subgeometry of $DH(2n - 1, 4)$ which is isometrically embedded into $DH(2n - 1, 4)$. Let f_1 and f_2 be two distinct valuations of \mathbb{G}_n . Then by Theorem 1.1 there exists a unique point x_i , $i \in \{1, 2\}$, of $DH(2n - 1, 4)$ such that the valuation f_i of \mathbb{G}_n is induced by the classical valuation f'_i of $DH(2n - 1, 4)$ with center x_i .

Suppose x_1 and x_2 are collinear. Then f'_1 and f'_2 are two neighboring valuations of $DH(2n - 1, 4)$ by Corollary 2.12(2). Proposition 2.8 then implies that f_1 and f_2 are neighboring valuations of \mathbb{G}_n .

Conversely, if f_1 and f_2 are neighboring valuations of \mathbb{G}_n , then by Lemma 3.6, x_1 and x_2 are collinear.

References

- [1] R. J. Blok and A. E. Brouwer. The geometry far from a residue. *Groups and geometries* (Siena, 1996), 29–38, Trends Math., Birkhäuser, Basel, 1998.
- [2] A. E. Brouwer. The uniqueness of the near hexagon on 729 points. *Combinatorica* 2 (1982), 333–340.
- [3] A. E. Brouwer. The uniqueness of the near hexagon on 759 points. *Finite geometries*, pp. 47–60, Lecture Notes in Pure and Appl. Math. 82, Dekker, New York, 1983.
- [4] A. E. Brouwer, A. M. Cohen, J. I. Hall and H. A. Wilbrink. Near polygons and Fischer spaces. *Geom. Dedicata* 49 (1994), 349–368.
- [5] A. E. Brouwer and E. W. Lambeck. An inequality on the parameters of distance regular graphs and the uniqueness of a graph related to M_{23} . *Ann. Discrete Math.* 34 (1987), 101–106.
- [6] A. E. Brouwer and H. A. Wilbrink. The structure of near polygons with quads. *Geom. Dedicata* 14 (1983), 145–176.
- [7] P. J. Cameron. Dual polar spaces. *Geom. Dedicata* 12 (1982), 75–85.
- [8] B. N. Cooperstein. On the generation of dual polar spaces of unitary type over finite fields. *European J. Combin.* 18 (1997), 849–856.
- [9] B. De Bruyn. On near-polygons and the Coxeter cap in $PG(5, 3)$. *J. Geom.* 68 (2000), 23–33.
- [10] B. De Bruyn. New near polygons from Hermitian varieties. *Bull. Belg. Math. Soc. Simon Stevin* 10 (2003), 561–577.

- [11] B. De Bruyn. *Near polygons*. Frontiers in Mathematics, Birkhäuser, Basel, 2006.
- [12] B. De Bruyn. The universal embedding of the near polygon \mathbb{G}_n . *Electron. J. Combin.* 14 (2007), Research paper 39, 12pp.
- [13] B. De Bruyn. On the Grassmann-embeddings of the hermitian dual polar spaces. *Linear Multilinear Algebra* 56 (2008), 665–677.
- [14] B. De Bruyn. An alternative definition of the notion valuation in the theory of near polygons. *Electron. J. Combin.* 16 (2009), Research paper 16, 14pp.
- [15] B. De Bruyn. On the Grassmann modules for the unitary groups. *Linear and Multilinear Algebra*, to appear.
- [16] B. De Bruyn. Isometric embeddings between dense near polygons. Preprint.
- [17] B. De Bruyn and A. Pasini. Generating symplectic and Hermitian dual polar spaces over arbitrary fields nonisomorphic to \mathbb{F}_2 . *Electron. J. Combin.* 14 (2007), Research Paper 54, 17 pp.
- [18] B. De Bruyn and P. Vandecasteele. Valuations of near polygons. *Glasg. Math. J.* 47 (2005), 347–361.
- [19] B. De Bruyn and P. Vandecasteele. The distance-2-sets of the slim dense near hexagons. *Ann. Comb.* 10 (2006), 193–210.
- [20] B. De Bruyn and P. Vandecasteele. The valuations of the near hexagons related to the Witt designs $S(5, 6, 12)$ and $S(5, 8, 24)$. *J. Combin. Des.* 14 (2006), 214–228.
- [21] B. De Bruyn and P. Vandecasteele. The valuations of the near octagon \mathbb{G}_4 . *Discrete Math.*, to appear.
- [22] A. Kasikova and E. E. Shult. Absolute embeddings of point-line geometries. *J. Algebra* 238 (2001), 265–291.
- [23] P. Li. On the universal embedding of the $U_{2n}(2)$ dual polar space. *J. Combin. Theory Ser. A* 98 (2002), 235–252.
- [24] S. E. Payne and J. A. Thas. *Finite generalized quadrangles*. Research Notes in Mathematics 110. Pitman, Boston, 1984.
- [25] M. A. Ronan. Embeddings and hyperplanes of discrete geometries. *European J. Combin.* 8 (1987), 179–185.
- [26] E. E. Shult. On Veldkamp lines. *Bull. Belg. Math. Soc. Simon Stevin* 4 (1997), 299–316.
- [27] E. E. Shult and A. Yanushka. Near n -gons and line systems. *Geom. Dedicata* 9 (1980), 1–72.
- [28] J. Tits. *Buildings of spherical type and finite BN-pairs*. Lecture Notes in Mathematics 386. Springer-Verlag, Berlin-New York, 1974.