

# Efficient covering designs of the complete graph

Yair Caro \*

and

Raphael Yuster †

Department of Mathematics

University of Haifa-ORANIM, Tivon 36006, Israel.

**AMS Subject Classification:** 05B05,05B40 (primary),  
05B30,51E05,94C30,62K05,62K10 (secondary).

Submitted: November 1, 1996; Accepted: February 3, 1997

## Abstract

Let  $H$  be a graph. We show that there exists  $n_0 = n_0(H)$  such that for every  $n \geq n_0$ , there is a covering of the edges of  $K_n$  with copies of  $H$  where every edge is covered at most twice and any two copies intersect in at most one edge. Furthermore, the covering we obtain is asymptotically optimal.

## 1 Introduction

All graphs considered here are finite, undirected and simple, unless otherwise noted. For the standard graph-theoretic notations the reader is referred to [5]. Let  $H = (V_H, E_H)$  be a graph. An  $H$ -covering design of a graph  $G = (V_G, E_G)$  is a set  $L = \{G_1, \dots, G_s\}$  of subgraphs of  $G$  such that each  $G_i$  is isomorphic to  $H$  and every edge  $e \in E_G$  appears in at least one member of  $L$ . The  $H$ -covering number of  $G$ , denoted by  $cov(G, H)$ , is the minimum number of members in an  $H$ -covering design of  $G$ . (If there is an edge of  $G$  which cannot be covered by a copy of  $H$ , we put  $cov(G, H) = \infty$ ). Clearly,  $cov(G, H) \geq |E_G|/|E_H|$ . In case equality holds, the  $H$ -covering design is called an  $H$ -decomposition (or  $H$ -design) of  $G$ . Two trivial necessary conditions for a decomposition are that  $|E_H|$  divides  $|E_G|$  and that  $gcd(H)$  divides  $gcd(G)$  where the  $gcd$  of a graph

---

\*e-mail: zeac603@uvm.haifa.ac.il

†e-mail: raphy@math.tau.ac.il

is the greatest common divisor of the degrees of all the vertices. In case  $G = K_n$ , it was shown by Wilson in [17] that the two necessary conditions are also sufficient, provided  $n \geq n_0(H)$ , where  $n_0(H)$  is a sufficiently large constant. If, however, the necessary conditions do not hold, the best one could hope for is an  $H$ -covering design of  $K_n$  where the following three properties hold:

1. **2-overlap:** Every edge is covered at most twice.
2. **1-intersection:** Any two copies of  $H$  intersect in at most one edge.
3. **Efficiency:**  $s|E_H| < \binom{n}{2} + c(H) \cdot n$ , where  $s$  is the number of members in the covering, and  $c(H)$  is some constant depending only on  $H$ .

The papers of Mills and Mullin [12] and of Brouwer [4], provide an excellent survey of covering designs. Covering designs with the 2-overlap property were first introduced in statistical designs by [10] and are also mentioned in [2], [6] and [11]. Covering designs with the 1-intersection property (also called super-simple designs) are mentioned by Adams et. al. in [1], Teirlinck [15, 16], Fort and Hedlund [8], Brouwer [3] and Schreiber [14]. The existence of efficient Covering designs of *complete hypergraphs* was first proved by Rödl in [13].

Our main result is that  $H$ -covering designs of  $K_n$ , having these three properties, exist for every fixed graph  $H$ , and for all  $n \geq n_0(H)$ :

**Theorem 1.1** *Let  $H$  be a fixed graph. There exists  $n_0 = n_0(H)$  such that if  $n \geq n_0$ ,  $K_n$  has an  $H$ -covering design with the 2-overlap, 1-intersection, and efficiency properties.*

## 2 Proof of the main result

We shall prove Theorem 1.1 whenever  $H = K_h$  is a complete graph. This suffices, since if  $H$  is not a complete graph, it is known by Wilson's theorem [17] that there exists an  $h_0 = h_0(H)$  such that  $K_{h_0}$  has an  $H$ -decomposition. By applying Theorem 1.1 to  $K_{h_0}$ , we shall obtain an  $n_0 = n_0(h_0) = n_0(H)$ , such that if  $n \geq n_0$ ,  $K_n$  has a  $K_{h_0}$ -covering design with the 2-overlap and 1-intersection properties and such that  $\binom{h_0}{2}s < \binom{n}{2} + h_0^3 \cdot n$ , where  $s$  is the number of members in the covering. Thus, there is an  $H$ -covering design of  $K_n$  with the 2-overlap and 1-intersection properties, and with  $s \frac{\binom{h_0}{2}}{|E_H|}$  elements, such that  $s \frac{\binom{h_0}{2}}{|E_H|} |E_H| < \binom{n}{2} + h_0^3 \cdot n = \binom{n}{2} + c(H) \cdot n$ .

Fix  $K_h$ , where  $h \geq 3$  (for  $h = 2$  the result is trivial), and let  $h_1$  be the minimum positive integer such that whenever  $n \geq h_1$  and  $\binom{h}{2}$  divides  $\binom{n}{2}$ , and  $h - 1$  divides  $n - 1$ ,  $K_n$  has a  $K_h$ -decomposition. As mentioned before, the existence of  $h_1$  is guaranteed by Wilson's Theorem [17]. Now let  $n \geq \max\{h^8, h_1 + h(h - 1)\}$ . We will show that  $K_n$  has a  $K_h$ -covering design, as required in

Theorem 1.1. Let  $k$  be the minimum positive integer such that  $\binom{h}{2}$  divides  $\binom{n-k}{2}$  and  $h-1$  divides  $n-k-1$ . It is easy to see that  $0 \leq k < h(h-1)$ . If  $k=0$  we are done, since in this case  $n$  satisfies the conditions in Wilson's Theorem, and there is a  $K_h$ -decomposition of  $K_n$ . Assume, therefore, that  $1 \leq k < h(h-1)$ , and put  $r = n - k$ . Note that  $r > h_1$ . Partition the vertices of  $K_n$  into two subsets. The *big* subset has  $r$  vertices, namely  $B = \{a_1, \dots, a_r\}$ . The *small* subset has  $k$  vertices, namely  $S = \{b_1, \dots, b_k\}$ . We create the members of our efficient covering design in three stages.

**Stage 1:** Let  $B_0$  be the subgraph induced by the vertices  $\{a_1, \dots, a_{r-1}\}$ . Note that  $B_0$  is a complete graph on  $r-1$  vertices, and since  $h-1$  divides  $r-1$ , there exists a  $K_{h-1}$ -factor in  $B_0$ . (Recall that an  $X$ -factor of a graph is a set of vertex-disjoint copies of  $X$  which cover all the vertices of the graph). Let  $F_1$  be such a factor. We repeat the following process for  $i = 2, \dots, k$ . Let  $B_{i-1}$  be the graph obtained from  $B_{i-2}$  after the edges of the members of  $F_{i-1}$  have been removed. Let  $F_i$  be a  $K_{h-1}$ -factor in  $B_{i-1}$ . In order to show that our process works, we need to show that a  $K_{h-1}$ -factor exists in  $B_{i-1}$ . We prove this by induction on  $i$ . For  $i=1$ , this is simply the factor  $F_1$  defined above. Assume the claim holds for all  $j < i$ . This implies that  $B_{i-1}$  is regular of degree  $(r-2) - (i-1)(h-2)$ . According to the theorem of Hajnal and Szemerédi [9] if  $(r-2) - (i-1)(h-2) \geq \frac{h-2}{h-1}(r-1)$  then  $B_{i-1}$  has a  $K_{h-1}$ -factor. Indeed,

$$(r-2) - (i-1)(h-2) \geq (r-2) - (k-1)(h-2) > (r-2) - h(h-1)(h-2) > r - h^3.$$

Since  $r - \frac{r-1}{h-1} > \frac{h-2}{h-1}(r-1)$  it suffices to show that  $r - h^3 \geq r - \frac{r-1}{h-1}$  and this holds since  $r = n - k > h^4$ . Having defined the  $K_{h-1}$ -factors  $F_1, \dots, F_k$ , we now define a set  $L_1$  of edge-disjoint copies of  $K_h$  in our  $K_n$ , which cover all the edges between  $S$  and  $\{a_1, \dots, a_{r-1}\}$ . This is done by joining the vertex  $b_i$  to every member of  $F_i$ , for  $i = 1, \dots, k$ . Note that whenever we join  $b_i$  to a member of  $F_i$  we obtain a copy of  $K_h$ . Note also that  $L_1$  has exactly  $k(r-1)/(h-1)$  members.

**Stage 2:** Since  $r \geq h_1$ , and since  $h-1$  divides  $r-1$  and  $\binom{h}{2}$  divides  $\binom{r}{2}$ , we have by Wilson's Theorem that the subgraph induced by  $B$  (which is a  $K_r$ ), has a  $K_h$ -decomposition. Fix a labeled  $K_h$ -decomposition  $D$  of this  $K_r$ . That is,  $D$  is a set of  $\binom{r}{2}/\binom{h}{2}$   $h$ -subsets of  $\{a_1, \dots, a_r\}$ , where for each  $1 \leq i < j \leq r$ , the pair  $(a_i, a_j)$  appears in exactly one member of  $D$ . If  $\pi$  is any permutation of  $\{1, \dots, r\}$  then let  $D_\pi$  be the labeled  $K_h$ -decomposition obtained from  $D$  by replacing each appearance of  $a_i$  in any member of  $D$  with  $\pi(a_i)$ , for  $i = 1, \dots, r$ . Our aim is to show that there exists a permutation  $\pi$ , and a set  $L^*$  of less than  $h^5$  members of  $L_1$  (recall that  $L_1$  is constructed in stage 1), such that every member of  $D_\pi$  intersects every member of  $L_1 \setminus L^*$  in at most one edge. In order to achieve this goal, we pick  $\pi$  randomly, where each of the  $r!$  permutations is equally likely. Consider two distinct edges  $(a_i, a_j)$  and  $(a_k, a_l)$  which both appear in the same member of  $L_1$  (note that when  $h=3$ , there is no such pair, since every member of  $L_1$  contains only two vertices of  $B$ ). We call such a pair of edges  $D_\pi$ -bad if they both appear in the same member of  $D_\pi$ . We shall

compute the probability that two *fixed* edges  $(a_i, a_j)$  and  $(a_k, a_l)$  are  $D_\pi$ -bad. Consider first the case where  $(a_i, a_j)$  and  $(a_k, a_l)$  share an endpoint, say  $a_k = a_i$ . Since  $\pi$  is random, the probability that  $(a_i, a_j)$  and  $(a_i, a_l)$  appear in the same member of  $D_\pi$  is *exactly*  $\frac{h-2}{r-2}$ . To see this, fix  $\pi(a_i)$  and  $\pi(a_j)$ , and let  $Q$  denote the unique member of  $D$  which contains both  $\pi(a_i)$  and  $\pi(a_j)$ . There are  $r - 2$  possible choices for  $\pi(a_l)$ , where  $h - 2$  of them result in a member of  $Q$ . Thus,  $D_\pi$  is *bad* with probability  $\frac{h-2}{r-2}$ , given that  $\pi(a_i)$  and  $\pi(a_j)$  are known. Note, however, that the expression  $\frac{h-2}{r-2}$  does not depend on the specific choices for  $\pi(a_i)$  and  $\pi(a_j)$ . Now consider the case where  $(a_i, a_j)$  and  $(a_k, a_l)$  are two independent edges (this is possible only if  $h - 1 \geq 4$ , since every member of  $L_1$  contains only  $h - 1$  vertices from  $B$ ). By a similar reasoning to the above, the probability that both these edges appear in the same member of  $D_\pi$  is exactly  $\frac{h-2}{r-2} \frac{h-3}{r-3}$ . There are  $(h - 1)(h - 2)(h - 3)/2$  pairs of adjacent edges of the form  $(a_i, a_j), (a_i, a_l)$  in every member of  $L_1$ . Thus, there are  $k \frac{r-1}{h-1} (h - 1)(h - 2)(h - 3)/2$  such pairs in all the members of  $L_1$ . There are  $3 \binom{h-1}{4}$  pairs of two independent edges of the form  $(a_i, a_j), (a_k, a_l)$  in every member of  $L_1$ . Thus there are  $3k \frac{r-1}{h-1} \binom{h-1}{4}$  such pairs in all the members of  $L_1$ . Therefore, if  $\mu$  is the expected number of  $D_\pi$ -bad pairs, then

$$\begin{aligned} \mu &= k \frac{r-1}{h-1} \frac{(h-1)(h-2)(h-3)}{2} \frac{h-2}{r-2} + k \frac{r-1}{h-1} 3 \binom{h-1}{4} \frac{h-2}{r-2} \frac{h-3}{r-3} < \\ & \frac{h^5}{2} + \frac{3}{24} h^7 \frac{r-1}{(r-2)(r-3)} < h^5. \end{aligned}$$

Thus, there exists a permutation  $\pi$  such that the number of  $D_\pi$ -bad pairs is less than  $h^5$ . Fix such a permutation, and let  $L_2 = D_\pi$ . Let  $L^*$  be the set of all members of  $L_1$  which contain a  $D_\pi$ -bad pair. Clearly,  $|L^*| < h^5$ . Thus, every member of  $L_2$  intersects every member of  $L_1 \setminus L^*$  in at most one edge. Put  $L_3 = L_2 \cup (L_1 \setminus L^*)$ .

**Stage 3:** Every edge of  $K_n$  appears in at most two members of  $L_3$  and any two members of  $L_3$  intersect in at most one edge. However, there may still be uncovered edges. In fact, all the  $\binom{k}{2}$  edges connecting two members of  $S$  are not covered, and all the  $k$  edges of the form  $(b_i, a_r)$ , for  $i = 1, \dots, k$ , are not covered. Furthermore, each member of  $L^*$  covers  $h - 1$  edges connecting some  $b_i \in S$  to a subset of  $h - 1$  vertices of  $\{a_1, \dots, a_{r-1}\}$ , and these edges are uncovered in  $L_3$ . Thus there are  $|L^*|(h - 1)$  uncovered edges of this form. Hence, if  $M$  denotes the set of uncovered edges, we have that

$$|M| = \binom{k}{2} + k + |L^*|(h - 1) < h^6.$$

The crucial point is that the number of uncovered edges is bounded by a constant depending only on  $h$ . We shall show how to sequentially create a set  $L_4$  of copies of  $K_h$ , beginning with  $L_4 = \emptyset$ ,

where at each stage, a new copy of  $K_h$  containing at least one non-covered edge by members of  $L_3 \cup L_4$ , is added to  $L_4$  (thus  $|L_4| < h^6$ ) and such that the following three invariants are maintained:

1. Every edge is covered at most twice by members of  $L_3 \cup L_4$ .
2. Any two members of  $L_3 \cup L_4$  intersect in at most one edge.
3. If  $L_4$  already contains  $j$  members, then any vertex of  $B \cup S$  is adjacent to at most  $jh + h^3$  edges which are covered twice by members of  $L_3 \cup L_4$ .

Note that at the beginning of the process, when  $L_4 = \emptyset$ , the first two invariants hold, since they hold for  $L_3$ . We must show that the third invariant holds initially, when  $j = 0$ . Indeed, in  $L_3$ , all the edges adjacent to a vertex of  $S$  are either non-covered, or covered once in  $L_1$ . Now consider a vertex  $a_i \in B$ . If  $i < r$ ,  $a_i$  is adjacent to exactly  $(h-2)k$  edges which are covered twice by members of  $L_1 \cup L_2$  (recall that  $a_r$  is not adjacent to any edge which is covered in  $L_1$ ). Since  $L_3 \subset L_1 \cup L_2$ , we have that any vertex in  $B \cup S$  is adjacent to at most  $(h-2)k < h^3$  edges which are covered twice by members of  $L_3$ .

Suppose  $L_4$  already contains  $j$  members, and there still exists an uncovered edge  $e = (q_1, q_2)$  in  $M$ . We shall find a set  $Q = \{q_3, \dots, q_h\}$  of  $h-2$  vertices in  $B \cup S$ , and add the complete graph  $K_h$  induced by  $\{q_1, q_2, \dots, q_h\}$  to  $L_4$ , while maintaining our three invariants. We select the elements of  $Q$  sequentially. The first element,  $q_3$ , needs to have the property that  $(q_1, q_3)$  is not covered twice, and  $(q_2, q_3)$  is not covered twice. Indeed there are at most  $2(jh + h^3)$  vertices of  $(B \cup S) \setminus \{q_1, q_2\}$  which are ruled out as candidates for  $q_3$ . Since

$$2(jh + h^3) < 2(h^7 + h^3) \leq h^8 - 2 \leq n - 2$$

we can find the desired  $q_3$ . It is important to note that there does not exist any member of  $L_3 \cup L_4$  which contains *both*  $(q_1, q_3)$  and  $(q_2, q_3)$ , since this would require it to contain  $(q_1, q_2)$  which we assume to be uncovered. Therefore, invariants 1 and 2 still hold. Suppose we have already found appropriate vertices  $q_3, \dots, q_i$ , where  $i < h$ , and we wish to find  $q_{i+1}$ . Our requirements of  $q_{i+1}$  are as follows: All the edges  $(q_t, q_{i+1})$  for  $t = 1, \dots, i$  should each be covered at most once, and for each once-covered edge  $(q_t, q_p)$  where  $1 \leq t < p \leq i$ ,  $q_{i+1}$  does not appear in the unique copy of  $L_3 \cup L_4$  containing  $(q_t, q_p)$ . These requirements rule out at most

$$i \cdot (jh + h^3) + \binom{i}{2}(h-2)$$

possible candidates for  $q_{i+1}$  from  $(B \cup S) \setminus \{q_1, \dots, q_i\}$ . In order to show that  $q_{i+1}$  can be selected we need to show that

$$n - i > i(jh + h^3) + \binom{i}{2}(h-2).$$

Indeed,

$$i(jh + h^3) + \binom{i}{2}(h-2) \leq (h-1)(h^7 + h^3) + \binom{h-1}{2}(h-2) < h^8 - (h-1) \leq n - i.$$

Our construction of  $Q$  shows that after adding the  $K_h$  subgraph induced by  $\{q_1, \dots, q_h\}$  as the  $j+1$ 'th element to  $L_4$ , invariants 1 and 2 still hold. Note also that invariant 3 holds as any vertex may only have at most  $h-1$  edges which are now covered twice, and which were not covered twice prior to this stage. (The only vertices for which this may happen are  $q_1, \dots, q_h$ ).

In order to complete our proof we only need to show that if  $L = L_3 \cup L_4$  contains  $s$  elements then  $s \binom{h}{2} < \binom{n}{2} + h^3 n$ . Clearly, it suffices to show that

$$sh(h-1) < n(n-1) + h^3(n-1). \quad (1)$$

$L_4$  contains less than  $h^6$  members.  $L_1$  contains exactly  $k(r-1)/(h-1)$  members, and  $L_2$  contains exactly  $\binom{r}{2}/\binom{h}{2}$  members. Thus,

$$s < h^6 + k \frac{r-1}{h-1} + \frac{\binom{r}{2}}{\binom{h}{2}}. \quad (2)$$

We shall prove (1) using (2) and using the facts that  $k < h(h-1)$ ,  $r = n - k$  and  $n \geq h^8$ . Indeed

$$sh(h-1) < h^7(h-1) + hk(r-1) + r(r-1) = h^8 - h^7 + hkn - hk^2 - hk + n^2 - 2kn + k^2 - n + k <$$

$$h^8 - h^3 + hkn + n^2 - 2kn - n < n(n-1) + h^3(n-1).$$

### 3- Concluding remarks and an open problem

When  $H = K_h$ , the constant  $n_0(H)$  in Theorem 1.1 is shown in the proof to be no larger than  $\max\{h^8, h_1 + h(h-1)\}$ , where  $h_1 = h_1(h)$  is the corresponding constant in Wilson's Theorem. However, the best known bound for  $h_1$  (and, consequently, for  $n_0(H)$ ), is rather large, and highly exponential in  $h$  [7]. It is plausible, however, that the statement of Theorem 1.1 is still valid for  $n_0(H)$  which is much smaller. In fact, we conjecture the following:

**Conjecture 3.1** *There exists a positive constant  $C$  such that for all  $h \geq 2$ , if  $n \geq Ch^2$  then  $K_n$  has a  $K_h$  covering design where each edge is covered at most twice and any two copies intersect in at most one edge.*

Note that a positive answer to Conjecture 3.1 requires a proof which does not use Wilson's Theorem, as improving Wilson's constant to  $O(h^2)$  is unlikely. The  $h^2$  factor in Conjecture 3.1 cannot be reduced since we have the following simple  $0.25h^2$  lower bound: Assume that  $h \geq 10$ . If  $n = \lfloor 0.25h^2 \rfloor$  then any  $K_h$ -covering of  $K_n$  contains  $\binom{n}{2} / \binom{h}{2} > h/2$  members. However, the union of  $t$   $K_h$ -subgraphs with the 1-intersection property contains at least  $h + (h-2) + \dots + (h-2t+2)$  vertices. For  $t = \lceil h/2 \rceil$  this sum is greater than  $0.25h^2 \geq n$ . Thus, any  $K_h$ -covering of  $K_n$  does not have the 1-intersection property.

## 4 Acknowledgment

The authors wish to thank T. Etzion and A. Rosa for useful discussions.

## References

- [1] P. Adams, E. Bryant and A. Khodkar, *On the existence of super-simple designs with block-size 4*, Aequationes Mathematicae 51 (1996), 230-246.
- [2] N. Alon, Y. Caro and R. Yuster, *Covering the edges of a graph by a prescribed tree with minimum overlap*, submitted.
- [3] A. E. Brouwer, *On the packing of quadruples without common triples*, Ars Combinatoria 5 (1978), 3-6.
- [4] A. E. Brouwer, *Block Designs*, in: Chapter 14 in "Handbook of Combinatorics", R. Graham, M. Grötschel and L. Lovász Eds. Elsevier, 1995.
- [5] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*, Macmillan Press, London, 1976.
- [6] Y. Caro, Y. Roditty and Y. Schonheim, *Covering the edges of the complete graph with minimum overlap*, Manuscript.
- [7] Y. Chang, *A bound for Wilson's Theorem - III*, J. Combinatorial Designs 4 (1996), 83-93.
- [8] M. K. Fort and G. A. Hedlund, *Minimal covering of pairs by triples*, Pacific J. Math. 8 (1958), 709-719.

- [9] A. Hajnal and E. Szemerédi, *Proof of a conjecture of Erdős*, in: *Combinatorial Theory and its Applications*, Vol. II (P. Erdős, A. Renyi and V. T. Sós eds.), Colloq. Math. Soc. J. Bolyai 4, North Holland, Amsterdam 1970, 601-623.
- [10] J. A. John and T. J. Mitchell, *Optimal incomplete block designs* J. Roy. Statist. Soc. 39B (1977) 39-43.
- [11] D. L. Kreher, G. F. Royle and W. D. Wallis, *A family of resolvable regular graph designs*, Discrete Math. 156 (1996), 269-275.
- [12] W. H. Mills and R. C. Mullin, *Coverings and packings*, in: *Contemporary Design Theory: A collection of Surveys*, 371-399, edited by J. H. Dinitz and D. R. Stinson. Wiley, 1992.
- [13] V. Rödl, *On a packing and covering problem*, Europ. J. of Combin. 6 (1985), 69-78.
- [14] S. Schreiber, *Covering all triples on  $n$  marks by disjoint Steiner systems*, J. Combin. Theory, Ser. A 15 (1973), 347-350.
- [15] L. Teirlinck, *On making two Steiner triple systems disjoint*, J. Combin. Theory, Ser. A 23 (1977), 349-350.
- [16] L. Teirlinck, *On large sets of disjoint quadruple systems*, Ars Combinatoria 17 (1984), 173-176.
- [17] R. M. Wilson, *Decomposition of complete graphs into subgraphs isomorphic to a given graph*, Congressus Numerantium XV (1975), 647-659.