

A central limit theorem for random walk in a random environment on marked Galton-Watson trees

Gabriel Faure*

Abstract

Models of random walks in a random environment were introduced at first by Chernoff in 1967 in order to study biological mechanisms. The original model has been intensively studied since then and is now well understood. In parallel, similar models of random processes in a random environment have been studied. In this article we focus on a model of random walk on random marked trees, following a model introduced by R. Lyons and R. Pemantle (1992). Our point of view is a bit different yet, as we consider a very general way of constructing random trees with random transition probabilities on them. We prove an analogue of R. Lyons and R. Pemantle's recurrence criterion in this setting, and we study precisely the asymptotic behavior, under restrictive assumptions. Our last result is a generalization of a result of Y. Peres and O. Zeitouni (2006) concerning biased random walks on Galton-Watson trees.

Key words: Random Walk, random environment, tree, branching random walk, central limit theorem.

AMS 2000 Subject Classification: Primary 60K37; 60F05; 60J80.

Submitted to EJP on July 13, 2010, final version accepted December 22, 2010.

*Laboratoire Analyse, Géométrie et Applications, UMR 7539, Institut Galilée, Université Paris 13, 99 avenue J.B. Clément 93430 Villetaneuse FRANCE. Research partially supported by the ANR project MEMEMO. faure@math.univ-paris13.fr

1 Introduction and statement of results.

Models of random walks in a random environment were introduced at first by Chernov in 1967 ([6]) in order to study biological mechanisms. The original model has been intensively studied since then and is now well understood. On the other hand, more recently, several attempts have been made to study extensions of this original model, for example in higher dimensions, continuous time, or different space.

It is remarkable that the random walk in \mathbb{Z}^d , $d > 1$, is still quite mysterious, in particular no precise criterion for recurrence/transience has ever been found.

In the case of trees, however, a recurrence criterion exists, and even estimates for the asymptotic behavior have been proven. To present our model and the existing results, we begin with some notations concerning trees. Let T be a tree rooted at some vertex e . For each vertex x of T we call $N(x)$ the number of his children $\{x_1, x_2, \dots, x_{N(x)}\}$, and \overleftarrow{x} his father. For two vertices $x, y \in T$, we call $d(x, y)$ the distance between x and y , that is the number of edges on the shortest path from x to y , and $|x| := d(e, x)$. Let T_n be the set of vertices such that $|x| = n$, and $T^* = T \setminus \{e\}$. We also note $x < y$ when x is an ancestor of y .

We call a marked tree a couple (T, A) , where A is a random application from the vertices of T to \mathbb{R}_+^* . Let \mathbb{T} be the set of marked trees. We introduce the filtration \mathcal{G}_n on \mathbb{T} defined as

$$\mathcal{G}_n = \sigma\{N(x), A(x_i), 1 \leq i \leq n, |x| < n, x \in T\}.$$

Following [20], given a probability measure q on $\mathbb{N} \otimes \mathbb{R}_+^{*\mathbb{N}^*}$, there exists a probability measure \mathbb{M} on \mathbb{T} such that

- the distribution of the random variable $(N(e), A(e_1), A(e_2), \dots)$ is q ,
- given \mathcal{G}_n , the random variables $(N(x), A(x_1), A(x_2), \dots)$, for $x \in T_n$, are independent and their conditional distribution is q .

We will always assume $m := E[N(e)] > 1$, ensuring that the tree is infinite with a positive probability.

We now introduce the model of random walk in a random environment. Given a marked tree T , we set for $x \in T^*$, x_i a child of x ,

$$\omega(x, x_i) = \frac{A(x_i)}{1 + \sum_{j=1}^{N(x)} A(x_j)}$$

and

$$\omega(x \overleftarrow{x}) = \frac{1}{1 + \sum_{j=1}^{N(x)} A(x_j)}.$$

Moreover we set $\omega(x, y) = 0$ whenever $d(x, y) \neq 1$,

It is easy to check that $(\omega(x, y))_{x, y \in T}$ is a family of non-negative random variables such that,

$$\forall x \in T, \sum_{y \in T} \omega(x, y) = 1,$$

and

$$\forall x \in T^*, A(x) = \frac{\omega(\overleftarrow{x}, x)}{\omega(\overleftarrow{x}, \overleftarrow{x})}, \tag{1}$$

where $\omega(e, \overleftarrow{e})$ is artificially defined as

$$\frac{1}{\omega(e, \overleftarrow{e})} = \sum_{|x|=1} A(x).$$

Further, $\omega(x, y) \neq 0$ whenever x and y are neighbors.

T will be called “the environment”, and we call “random walk on T ” the Markov chain (X_n, \mathbb{P}_T) defined by $X_0 = e$ and

$$\forall x, y \in T, \mathbb{P}_T(X_{n+1} = y | X_n = x) = \omega(x, y).$$

We call “annealed probability” the probability $\mathbb{P}_{\text{MT}} = \text{MT} \otimes \mathbb{P}_T$ taking into account the total alea.

We set, for $x \in T$, $C_x = \prod_{e < z \leq x} A(z)$. We can associate to the random walk X_n an electrical network with conductance C_x along $[\overleftarrow{x}, x]$, and a capacited network with capacity C_x along $[\overleftarrow{x}, x]$. We recall the definition of an electrical current on an electrical network. Let $G = (V, E)$ be a graph, C be a symmetric function on E , and A, Z be two disjoint subsets of V . We define the electrical current between A and Z as a function i that is antisymmetric on E such that, for any $x \in V \setminus (A \cup Z)$, the sum on the edges e starting from x of $i(e)$ equals zero (this is call Kirchhoff’s node Law), and, moreover, i satisfies the Kirchhoff’s cycle Law, that is, for any cycle $x_1, x_2, \dots, x_n = x_1$,

$$\sum_{i=1}^n \frac{i(x_i, x_{i+1})}{C(x_i, x_{i+1})} = 0.$$

A flow on a capacited network is an antisymmetric function θ that satisfies the Kirchhoff’s node Law, and such that, for all edges e , $\theta(e) < C(e)$, (for more precisions on this correspondence we refer to the chapters 2 and 3 of [17]).

We shall also frequently use the convex function ρ defined for $\alpha \geq 0$ as

$$\rho(\alpha) = E_{\text{MT}} \left[\sum_0^{N(e)} A(e_i)^\alpha \right] = E_q \left[\sum_0^N A(i)^\alpha \right].$$

Remark : This model is in fact inspired by a model introduced in [16]. In this case the tree T and the $A(x)$ were introduced separately, and the $A(x)$ were supposed to be independent. Here we can include models in which the structure of the tree and the transition probabilities are dependent. A simple example that is covered in our model is the following : Let T be a Galton-Watson tree. We chose an i.i.d. family $(B(x))_{x \in T}$ and set, for every $x \in T$, $1 \leq i \leq N(x)$, $A(x_i) = B(x)$. This way the transition probabilities to the children of any vertex are all equal, but randomly chosen. In R. Lyons and R. Pemantle’s article, a recurrence criterion was shown, our first result is a version of this criterion in our setting.

Theorem 1.1. *We suppose that there exists $0 \leq \alpha \leq 1$ such that ρ is finite in a small neighborhood of α , $\rho(\alpha) = \inf_{0 \leq t \leq 1} \rho(t) := p$ and $\rho'(\alpha) = E_q \left[\sum_{i=1}^{N(e)} A(e_i)^\alpha \log(A(e_i)) \right]$ is finite. We assume that $\sum_{i=1}^{N(e)} A(e_i)$ is not identically equal to 1.*

Then,

1. if $p < 1$ then the RWRE is a.s. positive recurrent, the electrical network has zero conductance a.s., and the capacited network admits no flow a.s..

2. if $p \leq 1$ then the RWRE is a.s. recurrent, the electrical network has zero conductance a.s. and the capacited network admits no flow a.s..
3. if $p > 1$, then, given non-extinction, the RWRE is a.s. transient, the electrical network has positive conductance a.s. and the capacited network admits flow a.s..

(By “almost surely” we mean “for MT almost every T ”).

Remark: In the case where $\sum_{i=1}^{N(e)} A(e_i)$ is identically equal to 1, which belongs to the second case, $|X_n|$ is a standard unbiased random walk, therefore X_n is null recurrent. However, there exists a flow, given by $\theta(\overleftarrow{x}, x) = C_x$.

The proof of this result is quite similar to the proof of R. Lyons and R. Pemantle, but there are some differences, coming from the fact that in their setting *i.i.d.* random variables appear along any ray of the tree, whereas it is not the case here. Results on branching processes will help us address this problem.

Theorem 1.1 does not give a full answer in the case $p = 1$, but this result can be improved, provided some technical assumptions are fulfilled. We introduce the condition

$$(H1) : \forall \alpha \in [0, 1], E_q \left[\left(\sum_0^{N(e)} A(e_i)^\alpha \right) \log^+ \left(\sum_0^{N(e)} A(e_i)^\alpha \right) \right] < \infty,$$

In the critical case, we have the following

Proposition 1.1. *We suppose $p = 1$, $m > 1$ and (H1). We also suppose that $\rho'(1) = E_q \left[\sum_{i=1}^{N(e)} A(e_i) \log(A(e_i)) \right]$ is defined and that ρ is finite in a small neighborhood of 1. Then,*

- if $\rho'(1) < 0$, then the walk is a.s. null recurrent, conditionally on the system’s survival,
- if $\rho'(1) = 0$ and for some $\delta > 0$,

$$E_{\text{MT}}[N(e)^{1+\delta}] < \infty,$$
 then the walk is a.s. null recurrent, conditionally on the system’s survival,
- if $\rho'(1) > 0$, and if for some $\eta > 0$, $\omega(x, \overleftarrow{x}) > \eta$ almost surely, then the walk is almost surely positive recurrent.

Remark: The distinction between the case $\rho'(1) = 0$ and $\rho'(1) > 0$ is quite unexpected.

The study of the critical case turns out to be quite interesting, indeed several different behaviors appear in this case. The quantity $\kappa = \inf\{t > 1, \rho(t) > 1\}$, associated to q is of particular interest. When $\rho'(1) \geq 0$, for regular trees and identically distributed $A(x)$, Y. Hu and Z. Shi showed ([9]) that there exist constants $0 < c_1 \leq c_2 < \infty$ such that

$$c_1 \leq \liminf_{n \rightarrow \infty} \frac{\max_{0 < s < n} |X_s|}{(\log n)^3} \leq \limsup_{n \rightarrow \infty} \frac{\max_{0 < s < n} |X_s|}{(\log n)^3} \leq c_2, \mathbb{P} - a.s..$$

It was recently proven by G. Faraud, Y. Hu and Z. Shi that $\frac{\max_{0 < s < n} |X_s|}{(\log n)^3}$ actually converges to an explicit constant (see [7]). Interestingly, this constant has a different form when $\rho'(1) = 0$ and when $\rho'(1) > 0$.




Figure 1: Possible shapes for ρ in the critical case

In the case $\rho'(1) < 0$, Y. Hu and Z. Shi showed ([8]) that

$$\lim_{n \rightarrow \infty} \frac{\log \max_{0 < s < n} |X_s|}{\log n} = 1 - \frac{1}{\min\{\kappa, 2\}}, \mathbb{P} - a.s..$$

Results in the case $p < 1$ have also been obtained by Y. Hu and Z. Shi ([8]), and the case $p > 1$ has been studied by E. Aidekon ([1]).

Let us go back to the critical case. Our aim is to study what happens when κ is large. When $\kappa \geq 2$, the walk behaves asymptotically like $n^{\frac{1}{2}}$. Our aim is to get a more precise estimate in this case. However we are not able to cover the whole regime $\kappa \in [2, \infty]$.

We first introduce the ellipticity assumptions

$$\exists 0 < \varepsilon_0 < \infty; \forall i, \varepsilon_0 \leq A(e_i) \leq \frac{1}{\varepsilon_0}, q - a.s. \quad (2)$$

and we assume that $(A(e_i))_{1 \leq i \leq N(e)}$ is of the form $(A'(i)\mathbb{1}_{(i \leq N(e))})_{i \geq 1}$, where $(A'(i))_{i \geq 1}$ is a i.i.d. family independent of $N(e)$ and that $E_q[N(e)^{\kappa+1}] < \infty$. (H2)

Remark : We actually only need this assumption to show Lemma 4.3, we can, for example, alternatively suppose that

$$\exists N_0; N(e) \leq N_0, q - p.s. \text{ et } P_q[N \geq 2|A(e_1)] \geq \frac{1}{N_0} \quad (3)$$

Note furthermore that those conditions imply (H1).

Theorem 1.2. *Suppose $N(e) \geq 1$, $q - a.s.$, (2), (3).*

If $p = 1$, $\rho'(1) < 0$ and $\kappa \in (8, \infty]$, then there is a deterministic constant $\sigma > 0$ such that, for MT almost every tree T , the process $\{|X_{[nt]}|/\sqrt{\sigma^2 n}\}$ converges in law to the absolute value of a standard Brownian motion, as n goes to infinity.

Remark : This result is a generalization of a central limit theorem proved by Y. Peres and O. Zeitouni [21] in the case of a biased standard random walk on a Galton-Watson tree. In this case, $A(x)$ is a constant equal to $\frac{1}{m}$, therefore $\kappa = \infty$. Our proof follows the same lines as theirs.

In the annealed setting, things happen to be easier, and we can weaken the assumption on κ .

Theorem 1.3. *Suppose $N(e) \geq 1$, $q - a.s.$, (2), (3). If $p = 1$, $\rho'(1) < 0$ and $\kappa \in (5, \infty]$, then there is a deterministic constant $\sigma > 0$ such that, under \mathbb{P}_{MT} , the process $\{|X_{[nt]}|/\sqrt{\sigma^2 n}\}$ converges in law to the absolute value of a standard Brownian motion, as n goes to infinity.*

Remark : As we will see, the annealed CLT will even be true for $\kappa \in (2, \infty)$, on a different kind of tree, following a distribution that can be described as “the invariant distribution” for the Markov chain of the “environment seen from the particle”.

We thank P. Mathieu for indicating to us the technique of C. Kipnis and S.R.S. Varadhan ([12]), that was quite an inspiration for us.

Our article will be organized as follows

- In section 2 we show Theorem 1.1.
- In section 3 we introduce a new law on trees, with particular properties.
- In section 4 we show a Central Limit Theorem for random walks on trees following the “new law”.
- In section 5 we expose a coupling between the original law and the new one.
- In section 6 we show some lemmas.
- In section 7 we show Theorem 1.3

2 Proof of Theorem 1.1.

Let us first introduce an associated martingale, which will be of frequent use in the sequence.

Let $\alpha \in \mathbb{R}^+$ and

$$Y_n^{(\alpha)} = \sum_{x \in T_n} \prod_{e < z \leq x} A(z)^\alpha = \sum_{x \in T_n} C_x^\alpha.$$

$Y_n^{(\alpha)}$ is known as Mandelbrot’s Cascade.

It is easy to see that if $\rho(\alpha) < \infty$ then $\frac{Y_n^{(\alpha)}}{\rho(\alpha)^n}$ is a non-negative martingale, with a.s. limit $Y^{(\alpha)}$.

We have the following theorem, due to J.D. Biggins (1977) (see [3, 4]) that allows us to know when $Y^{(\alpha)}$ is non trivial.

Statement 2.1 (Biggins). *Let $\alpha \in \mathbb{R}^+$. Suppose ρ is finite in a small neighborhood of α , and $\rho'(\alpha)$ exists and is finite, then the following are equivalent*

- given non-extinction, $Y^{(\alpha)} > 0$ a.s.,
- $P_{\text{MT}}[Y^{(\alpha)} = 0] < 1$,
- $E_{\text{MT}}[Y^{(\alpha)}] = 1$,
- $E_q \left[\left(\sum_0^{N(e)} A(e_i)^\alpha \right) \log^+ \left(\sum_0^{N(e)} A(e_i)^\alpha \right) \right] < \infty$, and
(H2): $:= \alpha \rho'(\alpha) / \rho(\alpha) < \log \rho(\alpha)$,
- $\frac{Y^{(\alpha)}}{\rho(\alpha)^n}$ converges in L^1 .

This martingale is related to some branching random walk, and has been intensively studied ([18, 3, 4, 13, 14, 19]). We will see that it is closely related to our problem.

Let us now prove Theorem 1.1. We shall use the following lemma, whose proof is similar to the proof presented in page 129 of [16] and omitted.

Lemma 2.1.

$$\min_{0 \leq t \leq 1} E \left[\sum_{x \in T_1} A(x)^t \right] = \max_{0 < y \leq 1} \inf_{t > 0} y^{1-t} E \left[\sum_{x \in T_1} A(x)^t \right].$$

(1) Let us begin with the subcritical case, We suppose there exists some $0 < \alpha < 1$ such that $\rho(\alpha) = \inf_{0 \leq t < 1} \rho(t) < 1$. Then, following [11] (Prop 9-131), and standard electrical/capacited network theory, if the conductances have finite sum, then the random walk is positive recurrent, the electrical network has zero conductance a.s., and the capacited network admits no flow a.s.. We have

$$\sum_{x \in T^*} C_x^\alpha = \sum_{n=0}^{\infty} \sum_{x \in T_n} C_x^\alpha = \sum_n \rho(\alpha)^n Y_n^{(\alpha)}.$$

Since $Y_n^{(\alpha)}$ is bounded (actually it converges to 0), we have

$$\sum_{x \in T^*} C_x^\alpha < \infty, \text{ MT} - a.s..$$

This implies that *a.s.*, for all but finitely many x , $C_x < 1$, and then $C_x \leq C_x^\alpha$, which gives the result.

(2) As before, we have α such that $\rho(\alpha) = \inf_{0 \leq t \leq 1} \rho(t) \leq 1$. We have to distinguish two cases. Either $\rho'(1) \geq 0$, therefore it is easy to see that, for α , (H2) is not verified, so

$$\sum_{x \in T_n} C_x^\alpha = Y_n^{(\alpha)} \rightarrow 0,$$

when n goes to ∞ . Then for n large enough, $C_x < 1$ for every $x \in T_n$, whence

$$\sum_{x \in T_n} C_x \rightarrow 0,$$

then by the *max-flow min-cut* theorem, the associated capacited network admits no flow a.s., this implies that no electrical current flows, and that the random walk is recurrent MT-a.s..

We now deal with the case where $\rho'(1) < 0$, then $\alpha = 1$. The proof is similar to [16], but, as it is quite short, we give it for the sake of clarity. We use the fact that, if the capacited network admits no flow from e , then the walk is recurrent.

We call F the maximum flows from e in T , and for $x \in T$, $|x| = 1$, we call F_x the maximum flow in the subtree $T_x = \{y \in T, x \leq y\}$, with capacity $\frac{C_y}{A(x)}$ along the edge (\overleftarrow{x}, x) . It is easy to see that F and F_x have the same distribution, and that

$$F = \sum_{|x|=1} A(x)(F_x \wedge 1). \quad (4)$$

Taking the expectation yields

$$E[F] = E[F_x \wedge 1] = E[F \wedge 1],$$

therefore $\text{ess sup } F \leq 1$. By independence, we obtain from (4) that

$$\text{ess sup } F = (\text{ess sup } \sum_{|x|=1} A(x))(\text{ess sup } F).$$

This implies that $F = 0$ almost surely, as $(\text{ess sup } \sum_{|x|=1} A(x)) > 1$, when $\sum_{|x|=1} A(x)$ is not identically equal to 1.

(3) We shall use the fact that, if the water flows when C_x is reduced exponentially in $|x|$, then the electrical current flows, and the random walk is transient a.s. (see [15]).

We have

$$\inf_{\alpha \in [0,1]} E \left[\sum_0^{N(e)} A(e_i)^\alpha \right] = p > 1$$

(p can be infinite, in which case the proof still applies).

We introduce the measure μ_n defined as

$$\mu_n(A) = E[\#\{A \cap \{\log C_x\}_{x \in T_n}\}],$$

where $\#$ denotes the cardinality.

One can easily check that

$$\phi_n(\lambda) := \int_{-\infty}^{+\infty} e^{\lambda t} d\mu_n(t) = E \left[\sum_{x \in T_n} C_x^\lambda \right] = \rho(\lambda)^n.$$

Let $y \in (0, 1]$ be such that $p = \inf_{t>0} y^{1-t} E[\sum_{x \in T_1} A(x)^t]$. Then, using Cramer-Chernov theorem (and the fact that the probability measure μ_n/m^n has the same Laplace transform as the sum of n independent random variables with law μ_1/m), we have

$$\frac{1}{n} \log \mu_n([n(-\log y), \infty)) \rightarrow \log(p/y).$$

Now, if we set $1/y < q < p/y$, there exists k such that

$$E[\#\{x \in T_k | C_x > y^k\}] > q^k.$$

Then the end of the proof is similar to the proof in [16]. We chose a small $\epsilon > 0$ such that,

$$E[\#\{x \in T_k | C_x > y^k, \text{ and } \forall e < z \leq x, A(z) > \epsilon\}] > q^k.$$

Let T^k be the tree whose vertices are $\{x \in T_{kn}, n \in \mathbb{N}\}$ such that $x = \overleftarrow{y}$ in T^k iff $x \leq y$ in T and $|y| = x + k$. We form a random subgraph $T^k(\omega)$ by deleting the edges (x, y) where

$$\prod_{x < z \leq y} A(z) < q^k \text{ or } \exists x < z \leq y, A(z) < \epsilon.$$

Let Γ_0 be the connected component of the root. The tree Γ_0 is a Galton-Watson tree, such that the expected number of children of a vertex is $q^k > 1$, hence with a positive probability Γ_0 is infinite and has branching number over q^k .

Using Kolmogoroff's 0-1 Law, conditionally to the survival there is almost surely a infinite connected component, not necessarily containing the root. This connected component has branching number at least q^k . Then we can construct almost surely a subtree T' of T , with branching number over q , such that $\forall x \in T', A(x) > \epsilon$ and if $|x| = nk, |y| = (n+1)k$ and $x < y$ then $\prod_{x < z \leq y} A(z) > q^k$. This implies the result.

We now turn to the proof of Proposition 1.1. Let π be an invariant measure for the Markov chain (X_n, P_T) (that is a measure on T such that, $\forall x \in T, \pi(x) = \sum_{y \in T} \pi(y) \omega(y, x)$), then one can easily check that

$$\pi(x) = \frac{\pi(e) \omega(e, \overleftarrow{e})}{\omega(x, \overleftarrow{x})} \prod_{0 < z \leq x} A(z),$$

with the convention that a product over an empty set is equal to 1.

Then almost surely there exists a constant $c > 0$ (dependant of the tree) such that

$$\pi(x) > c C_x.$$

Thus

$$\sum_{x \in T} \pi(x) > c \sum_n Y_n^{(1)}.$$

-If $\rho'(1) < 0$, then (H2) is verified and $Y > 0$ a.s. conditionally to the survival of the system, thus the invariant measure is infinite and the walk is null recurrent.

-If $\rho'(1) = 0$, we use a recent result from Y. Hu and Z. Shi. In [10] it was shown that, under the assumptions of Theorem 1.1, there exists a sequence λ_n such that

$$0 < \liminf_{n \rightarrow \infty} \frac{\lambda_n}{n^{1/2}} \leq \limsup_{n \rightarrow \infty} \frac{\lambda_n}{n^{1/2}} < \infty$$

and $\lambda_n Y_n^{(1)} \rightarrow_{n \rightarrow \infty} Y$, with $Y > 0$ conditionally on the system's survival. The result follows easily.

-If $\rho'(1) > 0$, there exists $0 < \alpha < 1$ such that $\rho(\alpha) = 1, \rho'(\alpha) = 0$. We set, for every $x \in T, \tilde{A}(x) := A(x)^\alpha$. We set accordingly $\tilde{C}(x) = \prod_{0 < z \leq x} \tilde{A}(z)$, and

$$\tilde{\rho}(t) := E_q \left[\sum_{i=1}^{N(e)} \tilde{A}(e_i)^t \right] = \rho(\alpha t).$$

Note that $\tilde{\rho}(1) = 1 = \inf_{0 < t \leq 1} \rho(t)$ and $\tilde{\rho}'(1) = 0$. Note that under the ellipticity condition $\omega(x, \overleftarrow{x}) > \eta$, for some constant $c > 0$

$$\sum_{x \in T} \pi(x) < c \sum_{x \in T} C_x = \sum_{x \in T} \tilde{C}_x^{1/\alpha}.$$

Using Theorem 1.6 of [10] with $\beta = 1/\alpha$ and $\tilde{C}_x = e^{-V(x)}$, we get that for any $\frac{2}{3}\alpha < r < \alpha$,

$$E_{\text{MT}} \left[\left(\sum_{x \in T_n} C_x \right)^r \right] = n^{-\frac{3r}{2\alpha} + o(1)}.$$

Note that as $r < 1$,

$$\left(\sum_n Y_n^{(1)} \right)^r \leq \sum_n (Y_n^{(1)})^r,$$

whence, using Fatou's Lemma,

$$E_{\text{MT}} \left[\left(\sum_{x \in T} C_x \right)^r \right] < \infty.$$

This finishes the proof.

3 The IMT law.

We consider trees with a marked ray, which are composed of a semi infinite ray, called $Ray = \{v_0 = e, v_1 = \overleftarrow{v_0}, v_2 = \overleftarrow{v_1} \dots\}$ such that to each v_i is attached a tree. That way v_i has several children, one of which being v_{i-1} .

As we did for usual trees, we can “mark” these trees with $\{A(x)\}_{x \in T}$. Let $\tilde{\mathbb{T}}$ be the set of such trees.

Let \mathcal{F}_n be the sigma algebra $\sigma(N_x, A_{x_i}, v_n \leq x)$ and $\mathcal{F}_\infty = \sigma(\mathcal{F}_n, n \geq 0)$. While unspecified, “measurable” will mean “ \mathcal{F}_∞ - measurable”.

Let \hat{q} be the law on $\mathbb{N} \times \mathbb{R}_+^{*\mathbb{N}^*}$ defined by

$$\frac{d\hat{q}}{dq} = \sum_1^{N(e)} A(e_i).$$

Remark : For this definition to have any sense, it is fundamental that $E_q[\sum_1^{N(e)} A_i] = 1$, which is provided by the assumptions $\rho'(1) < 0$ and $p = 1$.

Following [21], let us introduce some laws on marked trees with a marked ray. Fix a vertex v_0 (the root) and a semi infinite ray, called Ray emanating from it. To each vertex $v \in Ray$ we attach independently a set of marked vertices with law \hat{q} , except to the root e to which we attach a set of children with law $(q + \hat{q})/2$. We chose one of these vertices, with probability $\frac{A(v_i)}{\sum A(v_i)}$, and identify it with the child of v on Ray . Then we attach a tree with law MT to the vertices not on Ray . We call IMT the law obtained.

We call $\theta^v T$ be the tree T “shifted” to v , that is, $\theta^v T$ has the same structure and labels as T , but its root is moved to vertex v .

Note that as before, given a tree T in $\tilde{\mathbb{T}}$, we can define in a unique way a family $\omega(x, y)$ such that $\omega(x, y) = 0$ unless $d(x, y) = 1$,

$$\forall x \in T, \sum_{y \in T} \omega(x, y) = 1,$$

and

$$\forall x \in T, A(x) = \frac{\omega(\overleftarrow{x}, x)}{\omega(\overleftarrow{x}, \overleftarrow{x})}. \quad (5)$$

We call random walk on T the Markov chain (X_t, \mathbb{P}_T) on T , starting from v_0 and with transition probabilities $(\omega(x, y))_{x, y \in T}$.




Figure 2: The IMT law.

Let $T_t = \theta^{X_t} T$ denote the walk seen from the particle. T_t is clearly a Markov chain on $\tilde{\mathbb{T}}$. We set, for any probability measure μ on $\tilde{\mathbb{T}}$, $\mathbb{P}_\mu = \mu \otimes \mathbb{P}_T$ the annealed law of the random walk in a random environment on trees following the law μ . We have the following

Lemma 3.1. *IMT is a stationary and reversible measure for the Markov process T_t , in the sense that, for every $F : \tilde{\mathbb{T}}^2 \rightarrow \mathbb{R}$ measurable,*

$$\mathbb{E}_{\text{IMT}}[F(T_0, T_1)] = \mathbb{E}_{\text{IMT}}[F(T_1, T_0)].$$

Proof : Suppose G is a \mathcal{F}_n -measurable function, that is, G only depends on the (classical) marked tree of the descendants of v_n , to which we will refer as T^{-n} and on the position of v_0 in the $n - th$ level of T^{-n} . We shall write accordingly $G(T) = G(T^{-n}, v_0)$

We first show the following

Lemma 3.2. *If G is \mathcal{F}_n measurable, then*

$$E_{\text{IMT}}[G(T)] = E_{\text{MT}} \left[\sum_{x \in T_n} C_x G(T, x) \left(\frac{1 + \sum A(x_i)}{2} \right) \right]. \quad (6)$$

Remark : These formulae seem to create a dependency on n , which is actually irrelevant, since $E_q[\sum_{i=1}^{N(e)} A(e_i)] = 1$.

Proof : This can be seen by an induction over n , using the fact that

$$E_{\text{IMT}}[G(T^{-n}, v_0)] = E_q \left[\sum_{i=1}^N A(e_i) E \left[G(T'(i, N, A(e_j)), v_0) | i, N, A(e_j) \right] \right],$$

where $T'(x, N, A(e_i))$ is a tree composed of a vertex v_n with N children marked with the $A(e_i)$, and on each of this children is attached a tree with law MT, except on the i -th, where we attach a tree whose law is the same as $T^{-(n-1)}$.

Iterating this argument we have

$$E_{\text{IMT}}[G(T^{-n}, v_0)] = E_{\text{MT}} \left[\sum_{x \in T_n} C_x E [G(T''(x, T), x) | x, T] \right],$$

where the n first levels of $T''(x, T)$ are similar to those of T , to each $y \in T_n''$, $x \neq y$ is attached a tree with law MT, and to x is attached a set of children with law $(\hat{q} + q)/2$, upon which we attach MT trees. The result follows.

Let us go back to the proof of Lemma 3.1. Using the definition of the random walk, we get

$$\mathbb{E}_{\text{IMT}}[F(T_0, T_1)] = E_{\text{IMT}} \left[\sum_{x \in T} \omega(v_0, x) F(T, \theta^x T) \right].$$

Suppose F is $\mathcal{F}_{(n-2)} \times \mathcal{F}_{(n-2)}$ measurable; then $T \rightarrow F(T, \theta^x T)$ is at least $\mathcal{F}_{(n-1)}$ measurable. Then we can use (6) to get

$$\mathbb{E}_{\text{IMT}}[F(T_0, T_1)] = E_{\text{MT}} \left[\sum_{x \in T_n} C_x \left(\frac{1 + \sum A(x_i)}{2} \right) \sum_{y \in T} \omega(x, y) F(T, \theta^y T) \right].$$

It is easily verified that

$$\forall x, y \in T, \omega(x, y) \frac{1 + \sum A(x_i)}{2} C_x = \omega(y, x) \frac{1 + \sum A(y_i)}{2} C_y.$$

Using this equality, we get

$$\begin{aligned} \mathbb{E}_{\text{IMT}}[F(T_0, T_1)] &= E_{\text{MT}} \left[\sum_{x \in T_n} \sum_{y \in T} \omega(y, x) C_y \left(\frac{1 + \sum A(y_i)}{2} \right) F((T, x), (T, y)) \right] \\ &= E_{\text{MT}} \left[\sum_{y \in T_{n+1}} \omega(y, \overleftarrow{y}) C_y \left(\frac{1 + \sum A(y_i)}{2} \right) F((T, \overleftarrow{y}), (T, y)) \right] \\ &+ E_{\text{MT}} \left[\sum_{y \in T_{n-1}} \sum_i \omega(y, y_i) C_y \left(\frac{1 + \sum A(y_i)}{2} \right) F((T, y_i), (T, y)) \right]. \end{aligned}$$

Using (6) and the fact that F is $\mathcal{F}_{(n-2)} \times \mathcal{F}_{(n-2)}$ -measurable, we get

$$\begin{aligned} \mathbb{E}_{\text{IMT}}[F(T_0, T_1)] &= E_{\text{IMT}} \left[\omega(e, \overleftarrow{e}) F(\theta^{\overleftarrow{e}} T, T) \right] + E_{\text{IMT}} \left[\sum_i \omega(e, e_i) F(\theta^{e_i} T, T) \right] \\ &= \mathbb{E}_{\text{IMT}} [F(T_1, T_0)]. \end{aligned}$$

This finishes the proof of (3.1).

4 The Central Limit Theorem for the RWRE on IMT Trees.

In this section we introduce and show a central limit theorem for random walk on a tree following the law IMT. For $T \in \tilde{\mathbb{T}}$, let h be the horocycle distance on T (see Figure 2). h can be defined recursively by

$$\begin{cases} h(v_0) = 0 \\ h(\overleftarrow{x}) = h(x) - 1, \forall x \in T \end{cases}.$$

We have the following

Theorem 4.1. *Suppose $p = 1$, $\rho'(1) < 0$ and $\kappa \in [5, \infty]$, as well as assumptions (2) and (H2) or (3). There exists a deterministic constant σ such that, for IMT – a.e. T , the process $\{h(X_{\lfloor nt \rfloor})/\sqrt{\sigma^2 n}\}$ converges in distribution to a standard Brownian motion, as n goes to infinity.*

The proof of this result consists in the computation of a harmonic function S_x on T . We will show that the martingale S_{X_t} follows an invariance principle, and then that S_x stays very close to $h(x)$.

Let, for $v \in T$,

$$W_v = \lim_n \sum_{x \in T, v < x, d(v, x) = n} \prod_{v < z \leq x} A(z).$$

Statement 2.1 implies that $W_v > 0$ a.s. and $E[W_v | \sigma(A(x_i), N(x), x < v)] = 1$. Now, let $M_0 = 0$ and if $X_t = v$,

$$M_{t+1} - M_t = \begin{cases} -W_v & \text{if } X_{t+1} = \overleftarrow{v} \\ W_{v_i} & \text{if } X_{t+1} = v_i \end{cases}.$$

Given T , this is clearly a martingale with respect to the filtration associated to the walk. We introduce the function S_x defined as $S_e = 0$ and for all $x \in T$,

$$S_{x_i} = S_x + W_{x_i}, \quad (7)$$

in such a way that $M_t = S_{X_t}$.

Let

$$\eta = E_{\text{GW}}[W_0^2], \quad (8)$$

which is finite due to Theorem 2.1 of [13] (the assumption needed for this to be true is $\kappa > 2$). We call

$$V_t := \frac{1}{t} \sum_{i=1}^t \mathbb{E}_T[(M_{i+1} - M_i)^2 | \mathcal{F}_t]$$

the normalized quadratic variation process associated to M_t . We get

$$\mathbb{E}_T[(M_{i+1} - M_i)^2 | \mathcal{F}_t] = \omega(X_i, \overleftarrow{X_i}) W_{X_i}^2 + \sum_{j=1}^{N(X_i)} \omega(X_i, X_{ij}) W_{X_{ij}}^2 = G(T_i),$$

where X_{ij} are the children of X_i and G is a $L^1(\text{IMT})$ function on $\tilde{\mathbb{T}}$ (again due to $\kappa > 2$).

Let us define σ such that $E_{\text{IMT}}[G(T)] := \sigma^2 \eta^2$. We have the following

Proposition 4.1. *The process $\{M_{\lfloor nt \rfloor}/\sqrt{\sigma^2 \eta^2 n}\}$ converges, for IMT almost every T , to a standard Brownian motion, as n goes to infinity.*

Proof : We need the fact that when t goes to infinity,

$$V_t \rightarrow \sigma^2 \eta^2.$$

This comes from Birkhof's Theorem, using the transformation θ on $\tilde{\mathbb{T}}$, which conserves the measure IMT. The only point is to show that this transformation is ergodic, which follows from the fact that any invariant set must be independent of $\mathcal{F}_n^p = \sigma(N(x), A(x_i), v_n \leq x, h(x) < p)$, for all n, p , hence is independent of F_∞ .

The result follows then from the Central Limit Theorem for martingales. Our aim is now to show that $h(X_t)$ and M_t/η stay close in some sense, then the central limit theorem for $h(X_t)$ will follow easily.

Let

$$\epsilon_0 < 1/100, \delta \in (1/2 + 1/3 + 4\epsilon_0, 1 - 4\epsilon_0)$$

and for every t , let ρ_t be an integer valued random variable uniformly chosen in $[t, t + \lfloor t^\delta \rfloor]$.

It is important to note that, by choosing ϵ_0 small enough, we can get δ as close to 1 as we need.

We are going to show the following

Proposition 4.2. *For any $0 < \epsilon < \epsilon_0$,*

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(|M_{\rho_t}/\eta - h(X_{\rho_t})| \geq \epsilon \sqrt{t}) = 0, \text{ IMT - a.s.,}$$

further,

$$\lim_{t \rightarrow \infty} \mathbb{P}_T \left(\sup_{r,s < t, |r-s| < t^\delta} |h(X_r) - h(X_s)| > t^{1/2-\epsilon} \right) = 0, \text{ IMT - a.s..}$$

Before proving this result, we need some notations. For any vertex v of T , let

$$S_v^{\text{Ray}} = \sum_{y \text{ on the geodesic connecting } v \text{ and Ray}, y \notin \text{Ray}} W_y.$$

We need a fundamental result on marked Galton-Watson trees. For a (classical) tree T , and x in T , set

$$S_x = \sum_{e < y \leq x} W_x,$$

with W_x as before, and

$$A_n^\epsilon = \left\{ v \in T, d(v, e) = n, \left| \frac{S_v}{n} - \eta \right| > \epsilon \right\}.$$

We have the following

Lemma 4.2. *Let $2 < \lambda < \kappa - 1$, then for some constant C_1 depending on ϵ ,*

$$E_{\text{MT}} \left[\sum_{x \in A_n^\epsilon} C_x \right] < C_1 n^{1-\lambda/2}. \quad (9)$$

Proof : We consider the set \mathbb{T}^* of trees with a marked path from the root, that is, an element of \mathbb{T}^* is of the form (T, v_0, v_1, \dots) , where T is in \mathbb{T} , $v_0 = e$ and $v_i = \overleftarrow{v_{i+1}}$.

We consider the filtration $F_k = \sigma(T, v_1, \dots, v_k)$. Given an integer n , we introduce the law $\widehat{\text{MT}}_n^*$ on \mathbb{T}^* defined as follows : we consider a vertex e (the root), to this vertex we attach a set of marked children with law \hat{q} , and we chose one of those children as v_1 , with probability $P(x = v_1) = A(x) / \sum A(e_i)$. To each child of e different from v_1 we attach independently a tree with law MT, and on v_1 we iterate the process : we attach a set of children with law \hat{q} , we choose one of these children to be v_2 , and so on, until getting to the level n . Then we attach a tree with law MT to v_n .




Figure 3: the law $\widehat{\text{MT}}_n^*$.

The same calculations as in the proof of Lemma 3.2 allow us to see the following fact : for f F_n -measurable,

$$E_{\widehat{\text{MT}}_n^*} [f(T, v_0, \dots, v_n)] = E_{\text{MT}} \left[\sum_{x \in T_n} C_x f(T, p(x)) \right], \quad (10)$$

where $p(x)$ is the path from e to x . Note that, by construction, under $\widehat{\text{MT}}_n^*$ conditionally to $\tilde{F}_n^* := (C_{v_i}, 0 \leq i \leq n)$, the trees $T^{(v_i)}, 0 \leq i \leq n$ of the descendants of v_i who are not descendants of v_{i+1} are independent trees, and the law of $T^{(v_i)}$ is the law of a MT tree, except for the first level, whose law is \hat{q} conditioned on $v_{i+1}, A(v_{i+1})$.

For a tree T in \mathbb{T}^* we have

$$W_{v_k} = \sum_{v_k = \overleftarrow{x}, x \neq v_{k+1}} A(x)W_x + A(v_{k+1})W_{v_{k+1}} := W_k^* + A(v_{k+1})W_{v_{k+1}},$$

where

$$W_j^* = \lim_{n \rightarrow \infty} \sum_{x \in T, v_j < x, v_{j+1} \not\leq x, d(v_j, x) = n} \prod_{v \leq z \leq x} A(z).$$

Iterating this, we obtain

$$W_{v_k} = \sum_{j=k}^{n-1} W_j^* \prod_{i=k+1}^j A(v_i) + W_{v_n} \prod_{i=k+1}^n A(v_i),$$

with the convention that the product over an empty space is equal to one. We shall use the notation $A_i := A(v_i)$ for a tree with a marked ray.

Finally, summing over k , we obtain

$$S_{v_n} = \sum_{j=0}^{n-1} W_j^* \sum_{k=0}^j \prod_{i=k+1}^j A_i + W_{v_n} \sum_{k=0}^n \prod_{i=k+1}^n A_i. \quad (11)$$

Let $B_j = \sum_{k=0}^j \prod_{i=k+1}^j A_i$. We note for simplicity $W_{v_n} := W_n^*$. Note that

$$E_{\widehat{\text{MT}}_n^*}[W_0] = E_{\text{MT}} \left[\left(\sum_{x \in T_n} C_x \right)^2 \right] := E_{\text{MT}}[M_n^2]$$

converges to $\eta = E_{\text{MT}}[W_0^2]$ as n goes to infinity. Indeed, recalling that $E_{\text{MT}}[M_n] = 1$, we have

$$\begin{aligned} E_{\text{MT}}[(M_{n+1} - 1)^2] &= E_q \left[\left(\sum_{i=1}^{N(e)} A(e_i) U_i - 1 \right)^2 \right] \\ &= E_q \left[\left(\sum_{i=1}^{N(e)} A(e_i)(U_i - 1) + \sum_{i=1}^{N(e)} A(e_i) - 1 \right)^2 \right], \end{aligned}$$

where, conditionally to the A_i , U_i are i.i.d. random variables, with the same law as M_n . We get

$$E_{\text{MT}}[(M_{n+1} - 1)^2] = \rho(2)E_{\text{MT}}[(M_n - 1)^2] + C_2,$$

where C_2 is a finite number. It is easy to see then that $E[M_n^2]$ is bounded, and martingale theory implies that M_n converges in L^2 . Using the fact that $E_{\widehat{\text{MT}}_n^*}[W_{v_k}] = E_{\widehat{\text{MT}}_{n-k}^*}[W_0]$, a ‘‘Cesaro’’ argument implies that $E_{\widehat{\text{MT}}_n^*}[S_{v_n}]/n$ converges to η as n goes to infinity. In view of that and (10) it is clear that, for n large enough

$$\begin{aligned} E_{\text{MT}} \left[\sum_{x \in A_n^c} C_x \right] &\leq E_{\text{MT}} \left[\sum_{x \in T_n} C_x \mathbb{1}_{S_x - E_{\widehat{\text{MT}}_n^*}[S_x] > n\epsilon/2} \right] \\ &\leq P_{\widehat{\text{MT}}_n^*} \left[\left| S_{v_n} - E_{\widehat{\text{MT}}_n^*}[S_{v_n} | \tilde{F}_n^*] \right| > \frac{n\epsilon}{4} \right] \\ &\quad + P_{\widehat{\text{MT}}_n^*} \left[\left| E_{\widehat{\text{MT}}_n^*}[S_{v_n} | \tilde{F}_n^*] - E_{\widehat{\text{MT}}_n^*}[S_{v_n}] \right| > \frac{n\epsilon}{4} \right] := P_1 + P_2. \end{aligned}$$

Let us first bound P_1 . Let $\tilde{W}_j^* := W_j^* - E_{\widehat{\text{MT}}_n^*}[W_j^* | \tilde{F}_n^*]$ and $\lambda \in (2, \kappa - 1)$. We have

$$\begin{aligned} E_n^{(1)} &:= E_{\widehat{\text{MT}}_n^*} \left[\left| S_{v_n} - E_{\widehat{\text{MT}}_n^*}[S_{v_n} | \tilde{F}_n^*] \right|^\lambda \right] = E_{\widehat{\text{MT}}_n^*} \left[\left| \sum_{i=0}^n \tilde{W}_i^* B_i \right|^\lambda \right] \\ &= E_{\widehat{\text{MT}}_n^*} \left[E_{\widehat{\text{MT}}_n^*} \left[\left| \sum_{i=0}^n \tilde{W}_i^* B_i \right|^\lambda \middle| \tilde{F}_n^* \right] \right]. \end{aligned}$$

Inequality from page 82 of [22] implies

$$E_n^{(1)} \leq C(\lambda)n^{\lambda/2-1}E_{\widehat{MT}_n^*} \left[\sum_{i=0}^n E_{\widehat{MT}_n^*} \left[(\tilde{W}_i^* B_i)^\lambda | \tilde{F}_n^* \right] \right] \leq C_3 n^{\lambda/2-1} E_{\widehat{MT}_n^*} \left[\sum_{i=0}^n B_i^\lambda \right],$$

where we have admitted the following lemma

Lemma 4.3. $\forall \mu < \kappa$, there exist some constant C such that

$$E_{\widehat{MT}_n^*} [(W_i^*)^\mu | \tilde{F}_n^*] < C. \quad (12)$$

moreover, there exists some $\varepsilon_1 > 0$ such that

$$E_{\widehat{MT}_n^*} [W_i^* | \tilde{F}_n^*] > \varepsilon_1.$$

We postpone the proof of this lemma and finish the proof of Lemma 4.2. In order to bound $E_{\widehat{MT}_n^*} [B_i^\lambda]$ we need to introduce a result from [4] (lemma 4.1).

Statement 4.1 (Biggins and Kyprianou). *For any $n \geq 1$ and any measurable function G ,*

$$E_{\text{MT}} \left[\sum_{x \in T_n} C_x G(C_y, e < y \leq x) \right] = E[G(e^{S_i}; 1 \leq i \leq n)],$$

where S_n is the sum of n i.i.d variables whose common distribution is determined by

$$E[g(S_1)] = E_q \left[\sum_{i=1}^{N(e)} A(e_i) g(\log A(e_i)) \right]$$

for any positive measurable function g .

In particular, $E[e^{\lambda S_1}] = E_q[\sum_{i=1}^{N(e)} A(e_i)^{\lambda+1}] = \rho(\lambda + 1) < 1$. We are now able to compute

$$E_{\widehat{MT}_n^*} [B_n^\lambda] = E_{\text{MT}} \left[\sum_{x \in T_n} C_x \left(\sum_{e \leq y \leq x} \prod_{y < z \leq x} A(z) \right)^\lambda \right] = E \left[\left(\sum_{k=0}^n e^{S_n - S_k} \right)^\lambda \right].$$

Using Minkowski's Inequality, we get

$$E_{\widehat{MT}_n^*} [B_n^\lambda] \leq \left(\sum_{k=0}^n E \left[e^{\lambda(S_k - S_n)} \right]^{\frac{1}{\lambda}} \right)^\lambda \leq \left(\sum_{k=0}^n \rho(\lambda + 1)^{\frac{n-k}{\lambda}} \right)^\lambda \leq C_4. \quad (13)$$

We can now conclude,

$$E_n^{(1)} \leq C_5 n^{\lambda/2},$$

and by Markov's Inequality,

$$P_1 < C_6 / (\varepsilon^\lambda n^{\lambda/2}). \quad (14)$$

Now we are going to deal with

$$P_2 = P_{\widehat{MT}_n^*} \left[\left| E_{\widehat{MT}_n^*} [S_{v_n} | \tilde{F}_n^*] - E_{\widehat{MT}_n^*} [S_{v_n}] \right| > n\varepsilon/2 \right].$$

Lemma 4.3 implies that $E_{\widehat{\text{MT}}_n^*}[W_j^*|\tilde{F}_n^*]$ is bounded above and away from zero, and a deterministic function of A_{j+1} . We shall note accordingly

$$E_{\widehat{\text{MT}}_n^*}[W_j^*|\tilde{F}_n^*] := g(A_{j+1}). \quad (15)$$

Recalling (11), we have

$$E_{\widehat{\text{MT}}_n^*}[S_{v_n}|\tilde{F}_n^*] = \sum_{j=0}^n E_{\widehat{\text{MT}}_n^*}[W_j^*|\tilde{F}_n^*]B_j = \sum_{0 \leq j \leq k \leq n} \prod_{i=j}^k A_i g(A_{k+1}).$$

with the convention $g(A_{n+1}) = 1$ and $A_0 = 1$. We set accordingly

$$E_{\widehat{\text{MT}}_n^*}[S_{v_n}|\tilde{F}_n^*] := F(A_1, \dots, A_n).$$

Recalling that, due to Statement 4.1, under the law $\widehat{\text{MT}}_n^*$, the A_i are i.i.d random variables we get

$$E_{\widehat{\text{MT}}_n^*}[F(A_1, \dots, A_n)] = \sum_{0 \leq j \leq k \leq n} \prod_{i=j}^k E_{\widehat{\text{MT}}_n^*}[A_i] E_{\widehat{\text{MT}}_n^*}[g(A_{k+1})].$$

For $m \geq 0$ we call

$$\begin{aligned} F^m[A_{m+1}, \dots, A_n] \\ &:= \sum_{\substack{0 \leq j \leq k \leq n \\ k \leq m-1}} \prod_{i=j}^k E_{\widehat{\text{MT}}_n^*}[A_i] E_{\widehat{\text{MT}}_n^*}[g(A_{k+1})] + \sum_{\substack{0 \leq j \leq k \leq n \\ k \geq m}} \prod_{i=j}^m E_{\widehat{\text{MT}}_n^*}[A_i] \prod_{i'=m+1}^k A_{i'} g(A_{k+1}). \end{aligned}$$

Note that $F^0 = F$ and $F^n = E_{\widehat{\text{MT}}_n^*}[S_{v_n}]$, thus we can write

$$\begin{aligned} E_{\widehat{\text{MT}}_n^*}[S_{v_n}|\tilde{F}_n^*] - E_{\widehat{\text{MT}}_n^*}[S_{v_n}] &= F^0(A_1, \dots, A_n) - F^n \\ &= F^0(A_1, \dots, A_n) - F^1(A_2, \dots, A_n) \\ &+ F^1(A_2, \dots, A_n) - F^2(A_3, \dots, A_n) \dots \\ &+ F^{n-1}(A_n) - F^n. \end{aligned}$$

We introduce the notations $\rho := E_{\widehat{\text{MT}}_n^*}[A_1] = \rho(2) < 1$, and for a random variable X , $\tilde{X} := X - E_{\widehat{\text{MT}}_n^*}[X]$.

The last expression gives us

$$\begin{aligned}
& E_{\widehat{\text{MT}}_n^*}[S_{v_n}|\tilde{F}_n^*] - E_{\widehat{\text{MT}}_n^*}[S_{v_n}] \\
&= \tilde{g}(A_1) + \tilde{A}_1(g(A_2) + A_2g(A_3) + \dots + \prod_{i=2}^n A_i g(A_{n+1})) \\
&\quad + \rho \tilde{g}(A_2) + \tilde{A}_2(1 + \rho) \left[\sum_{j=3}^n \prod_{i=3}^j A_i g(A_{j+1}) \right] \\
&\quad + \rho^2 \tilde{g}(A_3) + \tilde{A}_3(1 + \rho + \rho^2) \left[\sum_{j=4}^n \prod_{i=4}^j A_i g(A_{j+1}) \right] + \dots \\
&\quad + \rho^{n-1} \tilde{g}(A_n) + \tilde{A}_n(1 + \rho + \rho^2 + \dots + \rho^{n-1}).
\end{aligned}$$

We deduce easily that

$$\left| E_{\widehat{\text{MT}}_n^*}[S_{v_n}|\tilde{F}_n^*] - E_{\widehat{\text{MT}}_n^*}[S_{v_n}] \right| < C_7 + C_8 \left| \sum_{k=1}^n \tilde{A}_k D_k (1 + \rho + \rho^2 + \dots + \rho^{k-1}) \right|, \quad (16)$$

where C_7, C_8 are finite constants and

$$D_k = \sum_{j=k+1}^n \prod_{i=k+1}^j A_i g(A_{j+1}).$$

To finish the proof of Lemma 4.2, we need to show that for every $\epsilon > 0$, $P_{\widehat{\text{MT}}_n^*}[\sum_{k=1}^n \tilde{A}_k D_k (1 + \rho + \rho^2 + \dots + \rho^{k-1}) > n\epsilon] < \frac{C(\epsilon)}{n^{\lambda/2-1}}$.

Recalling that $\lambda < \kappa - 1$, we can find a small $\nu > 0$ such that $\lambda(1 + \nu) < \kappa - 1$. Then we have, by Minkowski's Inequality

$$\begin{aligned}
E_{\widehat{\text{MT}}_n^*} [D_k^{\lambda(1+\nu)}] &\leq \left(\sum_{j=k+1}^n \left(E_{\widehat{\text{MT}}_n^*} \left[C_8 \prod_{i=k+1}^j A_i^{\lambda(1+\mu)} \right] \right)^{1/\lambda(1+\nu)} \right)^{\lambda(1+\nu)} \\
&\leq \left(\sum_{j=k}^n (C_9 \rho (1 + \lambda(1 + \mu))^{n-k+1})^{1/\lambda(1+\nu)} \right)^{\lambda(1+\nu)} < C_{10}.
\end{aligned} \quad (17)$$

Markov's Inequality then implies

$$P_{\widehat{\text{MT}}_n^*} \left[\max_{k \leq n} D_n > (\epsilon^2 n)^{\frac{1}{2(1+\nu)}} \right] \leq C_{11} \frac{n}{n^{\lambda/2} \epsilon^{\lambda}}. \quad (18)$$

On the other hand, we call for $0 \leq k \leq n$,

$$N_k := \sum_{j=n-k}^n D_j g(A_{n+1})(1 + \rho + \rho^2 + \dots \rho^{j-1}).$$

It is easy to check that N_k is a martingale with respect to the filtration $\mathcal{H}_k = \sigma(A_j, n-k \leq j \leq n)$. We can compute the quadratic variation of this martingale

$$\langle N_k \rangle := \sum_{j=1}^k E_{\widehat{\text{MT}}_n^*} [(N_k - N_{k-1})^2 | \mathcal{H}_{k-1}] = \rho(3) \sum_{j=1}^k (D_{n-j})^2.$$

On the other hand, the total quadratic variation of N_k is equal to

$$[N_k] := \sum_{j=1}^k (N_k - N_{k-1})^2 = \sum_{j=1}^k (\tilde{A}_{n-j} D_{n-j})^2.$$

It is easy to check that if the event in (18) is fulfilled, then there exists some constant C_{12} such that $\langle N_k \rangle < C_{12} n^{1+\frac{1}{2(1+\nu)}}$ and $[N_k] < C_{12} n^{1+\frac{1}{2(1+\nu)}}$. Therefore, using (18) and Theorem 2.1 of [2],

$$P_{\widehat{\text{MT}}_n^*} \left[\left| \sum_{k=1}^n \tilde{A}_k D_k \right| > n\epsilon \right] \leq C_{11} \frac{n}{n^{\lambda/2} \epsilon^\lambda} + 2 \exp - \frac{(\epsilon n)^2}{2C_{12} n^{1+\frac{1}{2(1+\nu)}}}. \quad (19)$$

Putting together (14) and (19), we obtain (9). This finishes the proof of Lemma 4.2. In particular, if $\kappa > 5$, we can choose $\lambda > 4$, so that

$$E_{\text{MT}} \left[\sum_{x \in \mathcal{A}_n^\epsilon} C_x \right] < n^{-\mu},$$

with $\mu > 1$. The following corollary is a direct consequence of the proof.

Corollary 4.4. *For every $a > 0$ and $2 < \lambda < \kappa - 1$,*

$$P_{\widehat{\text{MT}}_n^*} [|S_{v_k} - k\eta| > a] \leq C_1 \frac{k^{1-\lambda/2}}{a^\lambda}.$$

We now give the proof of Lemma 4.3. As we said in the introduction, for this lemma we need either the assumption (H2) or the assumption (3). We give the proof in both cases. Note that, by construction of MT^* , as, using Theorem 2.1 of [13], for every x a child of v_i , different from v_{i+1} , $W(x)$ has finite moments of order μ ,

$$E_{\widehat{\text{MT}}_n^*} [(W_i^*)^\mu | \tilde{F}_n^*] = C_0 E_{\widehat{\text{MT}}_n^*} \left[\left(\sum_{\bar{x}=v_i, x \neq v_{i+1}} A(x) \right)^\mu \middle| \tilde{F}_n^* \right] \quad (20)$$

$$= C_0 E_{\tilde{q}} \left[\left(\sum_{|x|=1, x \neq v_1} A(x) \right)^\mu \middle| A(v_1) \right] \quad (21)$$

Note that the upper bound is trivial under assumption (3). We suppose (H2), Let f be a measurable test function, we have by construction

$$\begin{aligned}
& E_{\bar{q}} \left[\left(\sum_{|x|=1, x \neq v_1} A(x) \right)^\mu f(A(v_1)) \right] \\
&= E_q \left[\sum_{i=1}^{N(e)} A(e_i) \left(\sum_{i \neq j} A(e_j) \right)^\mu f(A(e_i)) \right] \\
&\leq \sum_{n=1}^{\infty} P_q(N(e) = n) E_q \left[\sum_{i=1}^n A'(i) \left(\sum_{i \neq j} A'(1) \right)^\mu f(A'(1)) \right]
\end{aligned}$$

By standard convexity property, we get that the last term is lesser or equal to

$$\begin{aligned}
& \sum_{n=1}^{\infty} P_q(N(e) = n) E_q \left[\sum_{i=1}^n A'(i) n^{\mu-1} \sum_{i \neq j} A'(j)^\mu f(A'(i)) \right] \\
&\leq E_q[A'(1)^\mu] \sum_{n=1}^{\infty} P_q(N(e) = n) n^{\mu+1} E_q[A'(i) f(A'(i))] \\
&= E_q[A'(1)^\mu] E_q[A'(i) f(A'(i))] E_q[N(e)^{\mu+1}],
\end{aligned}$$

while, still by construction

$$\begin{aligned}
E_{\bar{q}} [f(A(v_1))] &= E_q \left[\sum_{i=1}^{N(e)} A(e_i) f(A(e_i)) \right] \\
&= \sum_{n=1}^{\infty} P_q(N(e) = n) E_q \left[\sum_{i=1}^n A'(i) f(A'(i)) \right] \\
&= E_q[N(e)] E_q[A'(1) f(A'(1))].
\end{aligned}$$

Therefore the result is direct. To prove the lower bound we begin with assumption (3). Actually we will only use the second part of this assumption, which is trivially implied by (H2), so the proof will also work for this case.

We have

$$\begin{aligned}
& E_{\hat{q}} \left[\sum_{|x|=1, x \neq v_1} A(x) f(A(v_1)) \right] = E_q \left[\sum_{i=1}^{N(e)} A(e_i) \left(\sum_{i \neq j} A(e_j) \right) f(A(e_i)) \right] \\
& \geq \epsilon_0 \sum_{i=1}^{\infty} E_q [A(e_i) f(A(e_i)) \mathbb{1}_{\{i \leq N(e)\}} (N(e) - 1)] \\
& \geq \epsilon_0 \sum_{i=2}^{\infty} E_q [A(e_i) f(A(e_i)) \mathbb{1}_{\{i \leq N(e)\}} (N(e) - 1)] + \epsilon_0 E_q [A(e_1) f(A(e_1)) (N(e) - 1)] \\
& \geq \epsilon_0 \sum_{i=2}^{\infty} E_q [A(e_i) f(A(e_i)) \mathbb{1}_{\{i \leq N(e)\}}] + \epsilon_0 E_q [A(e_1) f(A(e_1)) P(N(e) > 2|A(e_1))] \\
& \geq \frac{\epsilon_0}{N_0} E_q \left[\sum_{i=1}^{N(e)} A(e_i) f(A(e_i)) \right] = E_{\hat{q}} [f(A(v_1))],
\end{aligned}$$

indeed for $i \geq 2$, the event $\{i < N(e)\}$ implies $N(e) - 1 > 1$. This finishes the proof of Lemma 4.3.

Let us go back to IMT trees. We consider the following sets

$$\mathbf{B}_n^\epsilon = \left\{ v \in T, d(v, \text{Ray}) = n, \left| \frac{S_v^{\text{Ray}}}{n} - \eta \right| > \epsilon \right\}. \quad (22)$$

We can now prove the following

Lemma 4.5.

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(X_{\rho_t} \in \cup_{n=1}^{\infty} \mathbf{B}_n^\epsilon) = 0, \text{ IMT} - a.s..$$

Proof : we recall that a IMT tree is composed of a semi-infinite path from the root : $\text{Ray} = \{v_0 = e, v_1 = \overleftarrow{v_0} \dots\}$, and that

$$W_j^* = \lim_n \sum_{x \in T, v_j < x, v_{j-1} \not\leq x, d(v_j, x) = n} \prod_{v \leq z \leq x} A(z).$$

Recalling Lemma 4.3, under IMT, conditionally to $\{\text{Ray}, A(v_i)\}$, W_j^* are independent random variables and $E[W_j^*] > \epsilon_0$.

Let $1/2 < \gamma < \delta$. For a given tree T , we consider the event

$$\Gamma_t = \{\exists u \leq 2t | X_u = v_{\lfloor t^\gamma \rfloor}\}.$$

We have

$$\Gamma_t \subset \left\{ \inf_{u \leq 2t} M_u \leq S_{v_{\lfloor t^\gamma \rfloor}} \right\},$$

and IMT almost surely, for some ϵ ,

$$S_{v_{\lfloor t^\gamma \rfloor}} \leq - \sum_0^{\lfloor t^\gamma \rfloor} W_j^* < -\epsilon t^\gamma, \text{ for } t \text{ large enough.}$$

Since M_t is a martingale with bounded normalized quadratic variation V_t , we get that, for IMT almost every tree T ,

$$\mathbb{P}_T(\Gamma_t) \rightarrow 0.$$

Going back to our initial problem, we have

$$\mathbb{P}_T(X_{\rho_t} \in \cup_{m=1}^{\infty} \mathbf{B}_m^{\epsilon}) \leq \mathbb{P}_T(X_{\rho_t} \in \cup_{n=1}^{\infty} \mathbf{B}_n^{\epsilon}; \Gamma_t^c) + \mathbb{P}_T(\Gamma_t) \quad (23)$$

$$\leq \frac{1}{\lfloor t^{\delta} \rfloor} \mathbb{E}_T \left[\sum_{s=0}^{\lfloor t^{\gamma} \rfloor} \mathbb{1}_{X_s \in \cup_{m=1}^{\infty} \mathbf{B}_m^{\epsilon}} \right] + \mathbb{P}_T(\Gamma_t), \quad (24)$$

where $H_{v_{\lfloor t^{\gamma} \rfloor}}$ is the first time the walk hits $v_{\lfloor t^{\gamma} \rfloor}$.

As before we call $T^{(v_i)}$ the subtree constituted of the vertices $x \in T$ such that $v_i \leq x \not\leq x$. The first part of the right hand term of (23) is equal to

$$\frac{1}{\lfloor t^{\delta} \rfloor} \mathbb{E}_T \left[\sum_{i=0}^{\lfloor t^{\gamma} \rfloor} \sum_{s=0}^{H_{v_{\lfloor t^{\gamma} \rfloor}}} \mathbb{1}_{X_s \in \cup_{m=1}^{\infty} \mathbf{B}_m^{\epsilon} \cap T^{(v_i)}} \right] \leq \frac{1}{\lfloor t^{\delta} \rfloor} \sum_{i=0}^{\lfloor t^{\gamma} \rfloor} \mathbb{E}_T \left[\sum_{s=0}^{H_{v_{\lfloor t^{\gamma} \rfloor}}} \mathbb{1}_{X_s = v_i} \right] N_i,$$

where N_i is the P_T -expectation of the number of visits to $\overline{\cup_{n=1}^{\infty} \mathbf{B}_n^{\epsilon} \cap T^{(v_i)}}$ during one excursion in $T^{(v_i)}$. Lemma 4.2 implies that, under IMT conditioned on $\{Ray, A(v_i)\}$, N_i are independent and identically distributed variables, with finite expectation, up to a bounded constant due to the first level of those subtrees. We are now going to compute $\mathbb{E}_T \left[\sum_{s=0}^{H_{v_{\lfloor t^{\gamma} \rfloor}}} \mathbb{1}_{X_s = v_i} \right]$. Given T , we have

$$\sum_{s=0}^{H_{v_{\lfloor t^{\gamma} \rfloor}}} \mathbb{1}_{X_s = v_i} \leq 1 + M_i,$$

where M_i is the number of times the walk, leaving from v_i , gets back to v_i before hitting $v_{\lfloor t^{\gamma} \rfloor}$. M_i follows a geometric law, with parameter $p_i = \mathbb{P}_T^{v_i} [H_{v_{\lfloor t^{\gamma} \rfloor}} < H_{v_i}]$.

Standard computations for random walks on \mathbb{Z} , (see, for example, Theorem 2.1.12 of [24]) imply that

$$p_i = \frac{\omega(v_i, v_{i+1})}{1 + \sum_{j=i}^{\lfloor t^{\gamma} \rfloor - 1} \prod_{k=j-1}^{\lfloor t^{\gamma} \rfloor} A(v_k)},$$

and, going back to our initial problem,

$$\begin{aligned} \mathbb{P}_T(X_{\rho_t} \in \cup_{m=1}^{\infty} \mathbf{B}_m^{\epsilon}) &\leq \mathbb{P}_T(\Gamma_t) + \frac{C_{14}}{\lfloor t^{\delta} \rfloor} \sum_{i=0}^{\lfloor t^{\gamma} \rfloor} \left(1 + \sum_{j=i}^{\lfloor t^{\gamma} \rfloor - 1} \prod_{k=j-1}^{\lfloor t^{\gamma} \rfloor} A(v_k) \right) N_i \\ &\leq \mathbb{P}_T(\Gamma_t) + V_t \frac{C_{14}}{\lfloor t^{\delta} \rfloor} \sum_{i=0}^{\lfloor t^{\gamma} \rfloor} N_i, \end{aligned}$$

with $V_t = 1 + \sum_{j=0}^{\lfloor t^{\gamma} \rfloor - 1} \prod_{k=j-1}^{\lfloor t^{\gamma} \rfloor} A(v_k)$.

As in the proof of Lemma 4.2, statement 4.1 implies that $E_{\text{IMT}}[V_t^{\alpha}] < C_{15}$ for some $\alpha > 2$. Now we can choose δ close to one and γ close to $1/2$, and μ such that $1/\alpha < \mu < \delta - \gamma$

Markov's Inequality and the Borel Cantelli Lemma imply that, IMT-almost surely, there exists t_0 such that $\forall t > t_0, V_t \leq t^\mu$, and then,

$$\mathbb{P}_T(X_{\rho_t} \in \cup_{n=1}^{\infty} \mathbf{B}_n^\epsilon) \leq \mathbb{P}_T(\Gamma_t) + \frac{C_{16}}{t^{\delta-\mu}} \sum_{i=0}^{\lfloor t^\gamma \rfloor} N_i.$$

Since $\delta - \mu < \gamma$, an application of the law of large numbers finishes the proof of Lemma 4.5.

We are now able to prove the first part of Proposition 4.2. Note that under IMT, S_{v_n} follows the same law as S_{v_n} in a \mathbb{T}^* tree under $\widehat{\text{MT}}_n^*$, whence

$$S_{v_n}/n \xrightarrow[n \rightarrow \infty]{} -\eta$$

in probability. Let Q_t be the first ancestor of X_{ρ_t} on Ray . Statement 4.1 and standard RWRE theory imply that Q_t is transient, therefore

$$S_{Q_t}/h(Q_t) \xrightarrow[t \rightarrow \infty]{} \eta,$$

so that, for any positive ϵ_1 , for large t ,

$$|S_{Q_t}/\eta - h(Q_t)| \leq \epsilon_1 \sup_{s \leq 2t} |M_s|. \quad (25)$$

We can now compute

$$|M_{\rho_t}/\eta - h(X_{\rho_t})| = |S_{X_{\rho_t}}^{\text{Ray}}/\eta - d(X_{\rho_t}, \text{Ray}) + S_{Q_t}/\eta - h(Q_t)|.$$

In view of (25) on the event $\{X_{\rho_t} \notin \cup_{n=1}^{\infty} \mathbf{B}_n^\epsilon\}$, we have

$$|M_{\rho_t}/\eta - h(X_{\rho_t})| \leq 2\epsilon_1 \sup_{s \leq 2t} |M_s|.$$

The process V_t being bounded IMT *a.s.*, a standard martingale inequality implies

$$\lim_{\epsilon_1 \rightarrow 0} \limsup_{t \rightarrow \infty} \mathbb{P}_T^0(\sup_{s \leq t} |M_s| > \epsilon \sqrt{t}/(2\epsilon_1)) = 0.$$

It follows that

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(|M_{\rho_t}/\eta - h(X_{\rho_t})| \geq \epsilon \sqrt{t}) = 0, \text{ IMT} - a.s.$$

We are now going to prove the second part of Proposition 4.2. The course of the proof is similar to [21]. We have the following lemma

Lemma 4.6. *for any $u, t \geq 1$,*

$$\mathbb{P}_{\text{MT}}(|X_i| \geq u \text{ for some } i \leq t) \leq 2te^{-u^2/2t}.$$

Proof : We consider the graph T^* obtained by truncating the tree T after the level $u - 1$, and adding an extra vertex e^* , connected to all vertices in T_{u-1} . We construct a random walk X_s^* on T^* as following

$$\mathbb{P}_T^0(X_{i+1}^* = y | X_i^* = x) = \begin{cases} \omega(x, y) & \text{if } |x| < u - 1 \text{ or } |x| = u - 1, |y| = u - 2 \\ 1 - \omega(x, \overleftarrow{x}) & \text{if } |x| = u - 1, y = e^* \\ \tilde{\omega}(e^*, y) & \text{if } x = e^*, |y| = u - 1 \end{cases}.$$

We can choose $\tilde{\omega}(e^*, y)$ arbitrarily, provided $\sum_{y \in T_{u-1}} \tilde{\omega}(e^*, y) = 1$, so we will use this choice to ensure the existence of an invariant measure : indeed, if π is an invariant measure for the walk, one can easily check that, for any x such that $|x| \leq u-1$, calling $x^{(1)}$ the first vertex on the path from e to x ,

$$\pi(x) = \frac{\pi(e)\omega(e, x^{(1)})}{\omega(x, \overleftarrow{x})} \prod_{x^{(1)} < z \leq x} A(z).$$

Further, we need that, for every $x \in T_{u-1}$,

$$\pi(x)(1 - \omega(x, \overleftarrow{x})) = \pi(e^*)\tilde{\omega}(e^*, x).$$

Summing over x , and using $\sum_{y \in T_u} \tilde{\omega}(e^*, y) = 1$, we get

$$\begin{aligned} \pi(e^*) &= \pi(e) \sum_{x \in T_{u-1}} \omega(e, x^{(1)}) \prod_{x^{(1)} < z \leq x} A(z) \frac{\sum \omega(x, x_i)}{\omega(x, \overleftarrow{x})} \\ &\leq \pi(e) \sum_{x \in T_u} \prod_{x^{(1)} < z \leq x} A(z) \leq \pi(e) Y_u. \end{aligned}$$

Then,

$$\mathbb{P}_{\text{MT}}(\exists i \leq t, X_i \geq u) \leq \mathbb{P}_{\text{MT}}(\exists i \leq t, X_i^* = e^*) \leq \sum_{i=1}^t \mathbb{P}_{\text{MT}}(X_i^* = e^*).$$

By the Carne-Varnopoulos Bound (see [17], Theorem 12.1),

$$\mathbb{P}_T(X_i^* = e^*) \leq 2\sqrt{Y_u} e^{-u^2/2i}.$$

Since, by Jensen's Inequality, $E_{\text{MT}}(\sqrt{Y_n}) \leq 1$,

$$\mathbb{P}_{\text{MT}}(X_i \geq u \text{ for some } i \leq t) \leq 2te^{-u^2/2t}.$$

We have the following corollary, whose proof is omitted

Corollary 4.7.

$$\mathbb{P}_{\text{IMT}}(|h(X_i)| \geq u \text{ for some } i \leq t) \leq 4t^3 e^{-(u-1)^2/2t}.$$

Proof : see [21], Corollary 2.

We can now finish the proof of the second part of Proposition 4.2. Under \mathbb{P}_{IMT} , the increments $h(X_{i+1}) - h(X_i)$ are stationary, therefore, for any ϵ and $r, s \leq t$ with $|s - r| \leq t^\delta$,

$$\mathbb{P}_{\text{IMT}}(|h(X_r) - h(X_s)| \geq t^{1/2-\epsilon}) \leq \mathbb{P}_{\text{IMT}}(|h(X_{r-s})| \geq t^{1/2-\epsilon}) \leq 4t^3 e^{-t^{1-\delta-2\epsilon}}.$$

Whence, by Markov's Inequality, for all t large,

$$P_{\text{IMT}} \left(\mathbb{P}_T^0 (|h(X_{r-s})| \geq t^{1/2-\epsilon}) \geq e^{-t^{1-\delta-\epsilon}} \right) \leq e^{-t^{1-\delta-\epsilon}}.$$

Consequently,

$$P_{\text{IMT}} \left(\mathbb{P}_T^0 \left(\sup_{r,s \leq t, |r-s| \leq t^\delta} |h(X_r) - h(X_s)| \geq t^{1/2-\epsilon} \right) \geq e^{-t^{1-\delta-\epsilon}} \right) \leq e^{-t^{1-\delta-\epsilon}}.$$

The Borel-Cantelli Lemma completes the proof.

We are now able to finish the proof of Theorem 4.1. Due to Proposition 4.1, the process $\{M\lfloor nt\rfloor/\sqrt{\sigma^2\eta^2n}\}$ converges, for IMT almost every \mathbb{T} , to a standard Brownian motion, as n goes to infinity. Further, by Theorem 14.4 of [5], $\{M\rho_{nt}/\sqrt{\sigma^2\eta^2n}\}$ converges, for IMT almost every \mathbb{T} , to a standard Brownian motion, as n goes to infinity. Proposition 4.2 implies that the sequence of processes $\{Y_t^n\} = \{h(X_{\rho_{nt}})/\sqrt{\sigma^2n}\}$ is tight and its finite dimensional distributions converge to those of a standard Brownian motion, therefore it converges in distribution to a standard Brownian motion, and, applying again Theorem 14.4 of [5], so does $\{h(X_{\lfloor nt\rfloor})/\sqrt{\sigma^2n}\}$.

5 Proof of Theorem 1.2.

In this section we finish the proof of Theorem 1.2. Our argument relies on a coupling between random walks on MT and on IMT trees, quite similar to the coupling exposed in [21]. Let us introduce some notations : for T, S two trees, finite or infinite, we set LT the leaves of T , that is the vertices of T that have no offspring, $T^o = T/LT$ and for $v \in T$ we denote by $T \circ^v S$ the tree obtained by gluing the root of S to the vertex v of T , with vertices marked as in their original tree (the vertex coming from both v and the root of S is marked as v). Given a tree $T \in \mathbb{T}$ and a path $\{X_t\}$ on T we construct a family of finite trees T_i, U_i as follows : let $\tau_0 = \eta_0 = 0$, and U_0 the finite tree consisting of the root e of T and its offspring, marked as in T . For $i \geq 1$, let

$$\tau_i = \min\{t \geq \eta_{i-1} : X_t \in LU_{i-1}\} \quad (26)$$

$$\eta_i = \min\{t > \tau_i : X_t \in U_{i-1}^o\}. \quad (27)$$

Let T_i be the tree ‘‘explored’’ by the walk during the excursion $[\tau_i, \eta_i)$, that is to say T_i is composed of the vertices of T visited by $\{X_t, t \in [\tau_i, \eta_i)\}$, together with their offspring, marked as in T , and the root of T_i is X_{τ_i} . Let $U_i = U_{i-1} \circ^{X_{\tau_i}} T_i$ be the tree explored by the walk from the beginning. We call $\{u_t^i\}_{t=0}^{\eta_i - \tau_i - 1}$ the path in T_i defined by $u_t^i = X_{\tau_i + t}$. If T is distributed according to MT, and X_t is the path of the random walk on T , then, the walk being recurrent, \mathbb{P}_{MT} -almost surely $T = \lim U_i$.

We are now going to construct $\tilde{T} \in \tilde{\mathbb{T}}$, a tree with a semi-infinite ray emanating from the root, coupled with T , and a path $\{\tilde{X}_t\}$ on $\tilde{\mathbb{T}}$, in such a way that, if T is distributed according to MT, and X_t is the path of the random walk on T , then \tilde{T} will be distributed according to IMT and $\{\tilde{X}_t\}$ will follow the law of a random walk on \tilde{T} .

Let \tilde{U}_0 be the tree defined as follows : we choose a vertex denoted by e , as the root of \tilde{U}_0 , and a semi-infinite ray $\{e = v_0, v_1, \dots\}$. To each vertex $v_i \in \text{Ray}$ different from e we attach independently a set of marked vertices with law \hat{q} . To e we attach a set of children with distribution $(q + \hat{q})/2$ if $i \geq 1$ we chose one of those vertices, with probability $\frac{A(x)}{\sum_y A(y)}$, and identify it with v_{i-1} . We obtain a tree with a semi-infinite ray and a set of children for each vertex v_i on Ray , one of them being v_{i-1} .

We set $\tilde{\tau}_0 = \tilde{\eta}_0 = 0$. Recalling the relation (5) between the A_x and the $\omega(x, y)$, one can easily check that for any vertex x , knowing the $\{w(x, y)\}_{y \in T}$ is equivalent to knowing $\{A(x_i)\}_{x_i \text{ children of } x}$. Thus, knowing \tilde{U}_0 one can compute the $\{\omega(x, y)\}_{x \in \text{Ray}, y \in \tilde{U}_0}$ and define a random walk \tilde{X}_t on \tilde{U}_0 , stopped when it gets off Ray . We set accordingly $\tilde{\tau}_1 = \min\{t > 0 : \tilde{X}_t \in L\tilde{U}_0\}$.




Figure 4: the coupling

We are now going to “glue” the first excursion of $\{X_t\}$. Let

$$\begin{aligned}\tilde{U}_1 &= \tilde{U}_0 \circ^{\tilde{X}_{\tilde{\tau}_1}} T_1, \\ \tilde{\eta}_1 &= \tilde{\tau}_1 + \eta_1 - \tau_1, \\ \{\tilde{X}_t\}_{t=\tilde{\tau}_1}^{\tilde{\eta}_1-1} &= u_{t-\tilde{\tau}_1}^1, \\ \tilde{X}^{\tilde{\eta}_1} &= \overleftarrow{\tilde{X}^{\tilde{\eta}_1-1}}.\end{aligned}$$

One can easily check that $\{\tilde{X}_t\}_{t \leq \tilde{\eta}_1}$ follows the law of a random walk on \tilde{U}_1 .

We iterate the process, in the following way : for $i > 1$, start a random walk $\{\tilde{X}_t\}_{t \geq \tilde{\eta}_{i-1}}$ on \tilde{U}_{i-1} , and define

$$\begin{aligned}\tilde{\tau}_i &= \min\{t > 0 : \tilde{X}_t \in L\tilde{U}_{i-1}\}, \\ \tilde{U}_i &= \tilde{U}_{i-1} \circ^{\tilde{X}_{\tilde{\tau}_i}} T_i, \\ \tilde{\eta}_i &= \tilde{\tau}_i + \eta_i - \tau_i, \\ \{\tilde{X}_t\}_{t=\tilde{\tau}_i}^{\tilde{\eta}_i-1} &= u_{t-\tilde{\tau}_i}^i, \\ \tilde{X}^{\tilde{\eta}_i} &= \overleftarrow{\tilde{X}^{\tilde{\eta}_i-1}}.\end{aligned}$$

Finally, set $\tilde{U} = \bigcup_0^\infty \tilde{U}_i$ and \tilde{T} the tree obtained by attaching independents MT trees to each leaves of \tilde{U} . It is a direct consequence of the construction that

Proposition 5.1. *If T is distributed according to MT and X_t follows \mathbb{P}_T , then \tilde{T} is distributed according to IMT, and \tilde{X}_t follows $\mathbb{P}_{\tilde{T}}$.*

As a consequence, under proper assumptions on q , application of Proposition 4.1 implies that for MT almost every T the process $\{h(\tilde{X}_{\lfloor nt \rfloor})/\sqrt{\sigma^2 n}\}$ converges to a standard Brownian motion, as n goes to infinity.

We introduce $R_t = h(\tilde{X}_t) - \min_{i=1}^t h(\tilde{X}_i)$. We get immediately that

$\{R_{\lfloor nt \rfloor}/\sqrt{\sigma^2 n}\}$ converges to a Brownian motion reflected to its minimum, which has the same law as the absolute value of a Brownian motion.

In order to prove Theorem 1.2, we need to control the distance between R_t and $|X_t|$.

Let $I_t = \max\{i : \tau_i \leq t\}$ and $\tilde{I}_t = \max\{i : \tilde{\tau}_i \leq t\}$ the number of excursions started by $\{X_t\}$ and $\{\tilde{X}_t\}$ before time t . Let $\Delta_t = \sum_{i=1}^{I_t} (\tau_i - \eta_{i-1})$ and $\tilde{\Delta}_t = \sum_{i=1}^{\tilde{I}_t} (\tilde{\tau}_i - \tilde{\eta}_{i-1})$, which measure the time spent by $\{X_t\}$ and $\{\tilde{X}_t\}$ outside the coupled excursions before time t . By construction, the distance between R_t and $|X_t|$ comes only from the parts of the walks outside those excursion. In order to control these parts, we set for $0 \leq \alpha < 1/2$

$$\Delta_t^\alpha = \sum_{i=1}^{I_t} \sum_{s=\eta_{i-1}}^{\tau_i-1} \mathbb{1}_{|X_s| \leq t^\alpha};$$

similarly,

$$\tilde{\Delta}_t^\alpha = \sum_{i=1}^{\tilde{I}_t} \sum_{s=\tilde{\eta}_{i-1}}^{\tilde{\tau}_i-1} \mathbb{1}_{d(\tilde{X}_s, Ray) \leq t^\alpha}.$$

Finally, let

$$\mathbf{B}_t = \max_{0 \leq r < s \leq t, \tilde{X}_r \in \text{Ray}, \tilde{X}_s \in \text{Ray}} (h(\tilde{X}_s) - h(\tilde{X}_r)),$$

be the maximum amount the walk $\{\tilde{X}_t\}$ moves against the drift on Ray . We have the following

Proposition 5.2. *Under the assumptions of Theorem 1.2, for some $\alpha < 1/2$*

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(\Delta_t \neq \Delta_t^\alpha) = 0, \text{ MT} - a.s., \quad (28)$$

and

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(\tilde{\Delta}_t \neq \tilde{\Delta}_t^\alpha) = 0, \text{ IMT} - a.s.. \quad (29)$$

Further,

$$\limsup \frac{\Delta_t}{t} = 0, \text{ MT} - a.s., \quad (30)$$

and

$$\limsup \frac{\tilde{\Delta}_t}{t} = 0, \text{ IMT} - a.s.. \quad (31)$$

Finally,

$$\limsup \frac{\mathbf{B}_t}{\sqrt{t}} = 0, \text{ IMT} - a.s.. \quad (32)$$

Before proving this proposition, note that on the event $\{\Delta_t = \Delta_t^\alpha\} \cap \{\tilde{\Delta}_t = \tilde{\Delta}_t^\alpha\}$, we have

$$\min_{s: |s-t| \leq \Delta_t + \tilde{\Delta}_t} ||X_t| - R_s| \leq 2t^\alpha + \mathbf{B}_t.$$

Therefore we obtain that almost surely, there exists a time change θ_t such that, for t large enough,

$$\frac{|X_t - R_{\theta_t}|}{\sqrt{t}} \rightarrow_{t \rightarrow \infty} 0$$

and

$$\frac{|\theta_t - t|}{t} \rightarrow_{t \rightarrow \infty} 0.$$

As we said earlier, Proposition 4.1 implies that $\{R_{\lfloor nt \rfloor} / \sqrt{\sigma^2 n}\}$ converges, as n goes to infinity, to the law of the absolute value of a Brownian motion. Therefore so does $\{R_{\lfloor n\theta_t \rfloor} / \sqrt{\sigma^2 n}\}$. We deduce the result for $|X_t|$.

We now turn to the proof of Lemma 5.2. We introduce some notations: for $k \geq 1$, let $a_k = \sum_{j=1}^k \tau_j$, $b_k = \sum_{j=0}^{k-1} \eta_j$ and $J_k = [a_k - b_k + k, a_{k+1} - b_{k+1} + k]$. Note that $\{J_k\}_{k \geq 1}$ is a partition of \mathbb{N} , such that the length of J_k is equal to the time spent by the walk between the $k - th$ and the $k + 1 - th$ excursion. For $s \in J_k$, let $\mathbf{t}(s) = \eta_k + s - (a_k - b_k + k)$ and $Y_0 = 0, Y_1 = X_{\tau_1}$, and $Y_s = X_{\mathbf{t}(s)}$. $\{Y_s\}_{s \geq 0}$ is the walk X_n restricted off the excursions, it is clearly not Markovian, nevertheless, it is adapted to the filtration $G_s = \sigma(X_k, k \leq \mathbf{t}(s))$. For a fixed t , we set the sequence Θ_i of stopping times with respect to G_s defined by $\Theta_0 = 0$ and

$$\Theta_i = \min\{s > \Theta_{i-1} : |Y_s| - |Y_{\Theta_{i-1}}| = \lfloor (\log t)^{3/2} \rfloor\}.$$

Similarly, we set, for $k \geq 1$, $\tilde{a}_k = \sum_{j=1}^k \tilde{\tau}_j$, $\tilde{b}_k = \sum_{j=0}^{k-1} \tilde{\eta}_j$ and $\tilde{J}_k = [\tilde{a}_k - \tilde{b}_k + k, \tilde{a}_{k+1} - \tilde{b}_{k+1} + k]$, and for $s \in \tilde{J}_k$, we call $\tilde{t}(s) = \tilde{\eta}_k + s - (\tilde{a}_k - \tilde{b}_k + k)$ and $\tilde{Y}_0 = 0$, $\tilde{Y}_1 = \tilde{X}_{\tilde{\tau}_1}$, and $\tilde{Y}_s = \tilde{X}_{\tilde{t}(s)}$ the walk \tilde{X}_n restricted off the excursions. We set $\tilde{G}_s = \sigma(\tilde{X}_k, k \leq \tilde{t}(s))$. For a fixed t , we set the sequence of stopping times $\tilde{\Theta}_i$ with respect to \tilde{G}_s defined by $\tilde{\Theta}_0 = 0$ and

$$\tilde{\Theta}_i = \min \left\{ s > \tilde{\Theta}_{i-1} : \left| d(\tilde{Y}_s, \text{Ray}) - d(\tilde{Y}_{\tilde{\Theta}_{i-1}}, \text{Ray}) \right| = \lfloor (\log t)^{3/2} \rfloor \right\}.$$

We need the following lemma, whose demonstration will be postponed.

Lemma 5.1. *For all $\epsilon > 0$*

$$\lim_{t \rightarrow \infty} \mathbb{P}_T \left(\sum_{i=1}^{\lfloor t^{1/2+\epsilon} \rfloor} (\eta_i - \tau_i) < t \right) = 0, \text{ MT - a.s.}, \quad (33)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_T \left(\sum_{i=1}^{\lfloor t^{1/2+\epsilon} \rfloor} (\tilde{\eta}_i - \tilde{\tau}_i) < t \right) = 0, \text{ IMT - a.s.}, \quad (34)$$

$$\exists \epsilon' > 0 : \lim_{t \rightarrow \infty} \mathbb{P}_T \left(\exists s \leq t, W_{X_s} > t^{1/4-\epsilon'} \right) = 0, \text{ MT - a.s.}, \quad (35)$$

$$\text{and } \lim_{t \rightarrow \infty} \mathbb{P}_T \left(\exists s \leq t, W_{X_s} > t^{1/4-\epsilon'} \right) = 0, \text{ IMT - a.s.}, \quad (36)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_T (\exists k \leq I_t, \Theta_{i-1}, \Theta_i \in J_k, |Y_{\Theta_i}| > |Y_{\Theta_{i-1}}|) = 0, \text{ MT - a.s.}, \quad (37)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_T (\exists k \leq I_t, \tilde{\Theta}_{i-1}, \tilde{\Theta}_i \in \tilde{J}_k, d(\tilde{Y}_{\tilde{\Theta}_i}, \text{Ray}) > d(\tilde{Y}_{\tilde{\Theta}_{i-1}}, \text{Ray})) = 0, \text{ IMT - a.s.}, \quad (38)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_T (X_s \in \cup_{k=t^\alpha - (\log t)^2}^t \mathbf{A}_k^\epsilon \text{ for some } s \leq t) = 0, \text{ MT - a.s.}, \quad (39)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_T (X_s \in \cup_{k=t^\alpha - (\log t)^2}^t \mathbf{B}_k^\epsilon \text{ for some } s \leq t) = 0, \text{ IMT - a.s.} \quad (40)$$

Using this lemma, we can finish the proof of Proposition 5.2. We shall prove the following statement, which implies (28) : for some $\alpha \leq 1/2$,

$$\lim_{t \rightarrow \infty} \mathbb{P}_T \left(\max_{s \in \cup_{k=1}^t J_k} |Y_s| \geq t^\alpha \right) = 0, \text{ MT - a.s.} \quad (41)$$

It is a direct consequence of (33) and (37) that, MT almost surely, with \mathbb{P}_T probability approaching 1 as t goes to infinity,

$$t(\Theta_{2t^{1/2+\epsilon}}) > t,$$

whence, using lemma 5.1,

$$\begin{aligned} & \lim_{t \rightarrow \infty} \mathbb{P}_T \left(\max_{s \in \cup_{k=1}^t J_k} |Y_s| \geq t^\alpha \right) \\ & \leq \limsup_{t \rightarrow \infty} \sum_{i=0}^{2t^{1/2+\epsilon}} \mathbb{P}_T \left(\exists j > i : |Y_{\Theta_j}| \geq t^\alpha - (\log t)^2, Y_{\Theta_i} = e, \right. \\ & \quad \left. S_{Y_{\Theta_j}} \geq (\eta - \epsilon_1)t^\alpha/2, |Y_{\Theta_k}| > 0, \forall i < k \leq j; \right. \\ & \quad \left. |S_{X_s} - |X_s|| \leq \epsilon |X_s|, \forall s \leq t \right) := \limsup_{t \rightarrow \infty} \sum_{i=1}^{2t^{1/2+\epsilon}} P_{i,t}; \end{aligned}$$

where ϵ', ϵ_1 are positive numbers that can be chosen arbitrarily small.

For a fixed i and a fixed t , we set $\tilde{M}_s = S_{X_{\Theta_{i+s}}}$, and

$$K_t = \min \{s > 1 : X_r = 0 \text{ for some } r \in [s(\theta_{i+1}), s(\theta_{i+t})]\}.$$

The process $\{N_s\} = \{\tilde{M}_{s \wedge K_t} - \tilde{M}_1\}$ is a supermartingale with respect to the filtration $\tilde{G}'_s = \tilde{G}_{\Theta_{i+s}}$; indeed as long as the walk does not come back to the root, the conditional expectation of $S_{Y_{s+1}} - S_{Y_s}$ is lesser or equal to 0, and by construction the walk can only return to the root at a Θ_i .

Note that M_s and N_s depend on t , whereas this is omitted in the notation. Let A_s be the predictable process such that $N_s + A_s$ is a martingale.

Note that, on the event $\{W_{X_s} \leq t^{1/4-\epsilon'}, \forall s \leq t\}$ the increments of N_s are bounded by $t^{1/4-\epsilon'}(\log t)^{3/2}$. One can easily see that the increments of A_s are also bounded by $t^{1/4-\epsilon'}(\log t)^{3/2}$. Therefore Azuma's Inequality implies

$$P_{i,t} \leq \exp \left(-t^{2\alpha}/t^{1/2} + 2\epsilon + 2(1/4 - \epsilon') \right).$$

Recalling that we can choose ϵ arbitrarily small and α arbitrarily close to $1/2$, we get the result.

The proof of (29) is quite similar and omitted.

To prove (30) we introduce

$$T^\epsilon(t) = \min \{s : |X_s| \geq t^{1/2+\epsilon}\}. \quad (42)$$

By Lemma 4.6, we have

$$\mathbb{P}_{\text{MT}}(T_\epsilon(t) < t) \leq t e^{-t^{2\epsilon}}.$$

Using the Borel-Cantelli Lemma, we get that, MT almost surely

$$\mathbb{P}_T(T_\epsilon(t) < t) \leq e^{-t^\epsilon} \text{ for } t > t_0(T). \quad (43)$$

Let $C_{0,l}$ be the conductance between the root and the level l of the tree. Recalling that for w an offspring of v , the conductance associated to the edge $[v, w]$ is C_w , Thomson's principle implies that

$$C_{0,l}^{-1} = \inf_f \sum_{i=0}^l \sum_{v \in T_i} \sum_{w \text{ offspring of } v} \frac{f_{v,w}^2}{C_w}.$$

As one can easily check, $f_{v,w} = \frac{C_w W_w}{W_e}$ is a unit flow from the root to T_l , so we get

$$C_{0,l}^{-1} \leq \frac{1}{W_e} \sum_{i=1}^l \sum_{v \in T_i} C_w W_w^2.$$

As, conditionally to \mathcal{G}_i , W_w^2 are independent and identically distributed variables, with finite moment of order two (the assumption needed for that is $\kappa > 4$), we have

$$E_{\text{MT}} \left[\left(\sum_{v \in T_i} C_w W_w^2 - \sum_{v \in T_i} C_w E_{\text{MT}}[W_w^2] \right)^2 \right] \leq C_{17} \rho(2)^i,$$

for some constant C_{17} , then, using Markov's Inequality, for every $\nu > 0$ there exists a constant C_{18} such that

$$P_{\text{MT}} \left[\sum_{v \in T_i} C_w |W_w^2 - E[W_w^2]| > \nu \right] \leq C_{18} \rho(2)^i.$$

This is summable, so by the Borel-Cantelli Lemma, for some constant $C(T)$ dependant only on T , we get

$$\sum_{v \in T_i} C_w W_w^2 \leq C(T) \sum_{v \in T_i} C_w.$$

The last part being convergent, thus bounded, we get

$$C_{0,l}^{-1} \leq C(T)l. \tag{44}$$

If $L_0(t)$ denotes the number of visits to the root before time t , we get

$$\mathbb{E}_T [L_0(T_\epsilon(t))] = 1 + C_{0,t}^{-1/2+\epsilon},$$

indeed $L_0(T_\epsilon(t)) - 1$ follows a geometric law with parameter $1 - C_{0,t}^{-1/2+\epsilon}$.

Let $N_t(\alpha) = \sum_{k=0}^t \mathbb{1}_{|X_k| \leq t^\alpha}$. On the event that $T_\epsilon(t) > t$, we have, using Markov's property,

$$\mathbb{E}_T [N_t(\alpha); T_\epsilon(t) > t] \leq \mathbb{E}_T [L_0(T_\epsilon(t))] \pi \left(\bigcup_0^{t^\alpha} T_t \right) \leq C_{19}(T) t^{1/2+\epsilon+\alpha}.$$

Thus as $\mathbb{P}_T(T_\epsilon(t) \leq t) \leq C_{19}(T)e^{-n^\epsilon}$, using the monotonicity of $N_n(\alpha)$, we obtain $N_t(\alpha)/t \rightarrow 0$, from which the result follows, as $\Delta_t^\alpha \leq N_t^\alpha$ and $\mathbb{P}_T(\Delta_t \neq \Delta_t^\alpha) \rightarrow 0$.

Now we turn to the proof of (31). By the same calculations as in the proof of Lemma 4.5, for $\kappa > 5$, we get that $\mathbb{E}_{\text{IMT}}[\sum_{s \leq t} \mathbb{1}_{d(X_s, Ray) < t^\alpha}] \leq t^{1/2+\alpha+\epsilon}$ for any $\epsilon > 0$, from which the result follows by an application of Markov's Inequality and the Borel-Cantelli Lemma, using also the fact that the quantity in the expectation is non-decreasing in n .

The conductance from v_k to v_{k-u} is at most $C_{v_{k-u}}$, thus we have the bound

$$\mathbb{P}_T(\mathbf{B}_t > u) \leq t \sum_{k=u}^t \prod_{i=k}^{k-u} A(v_i).$$

By Theorem 2.1 and Lemma 3.2, the IMT-expectation of the right hand side is of order at most $t^2 \rho(2)^u$, therefore (32) follows by standard arguments.

6 Proof of Lemma 5.1.

It is clear that (33) and (34) are equivalent. We postpone the proof of these parts to the end of the section.

Proof of (35) : following [21], we call “fresh time” a time where the walk explore a new vertex, we have

$$\begin{aligned} \mathbb{P}_{\text{MT}}\left(\exists s \leq t, W_{X_s} > t^{1/4-\epsilon'}\right) &\leq \sum_0^t \mathbb{P}_{\text{MT}}[W_{X_s} > t^{1/4-\epsilon'}; s \text{ is a fresh time}] \\ &= \mathbb{P}_{\text{MT}}[W_0 > t^{1/4-\epsilon'}] < C_{20}t/t^{\mu(1/4+\epsilon')}, \end{aligned}$$

for $\mu < \kappa$. If $\kappa > 8$, for ϵ small enough, we can chose μ such that this is summable. Then the Borel-Cantelli Lemma implies the result.

Proof of (36) We are going to use the same arguments, excepted that we have to treat separately the vertices on *Ray*. More precisely

$$\begin{aligned} \mathbb{P}_{\text{MT}}\left(\exists s \leq t, W_{X_s} > t^{1/4-\epsilon'}\right) \\ \leq \sum_0^t \mathbb{P}_{\text{IMT}}[W_{X_s} > t^{1/4-\epsilon'}; s \text{ is a fresh time and } X_s \notin \text{Ray}] + \\ \mathbb{P}_{\text{MT}}\left(\exists s \leq t, W_{v_s} > t^{1/4-\epsilon'}\right). \end{aligned}$$

The second term is easily bounded, and the first one is similar to the previous case.

Proof of (37) : the event in the probability in (37) implies that, before time t the walk X_s gets to some vertex u , situated at least at a distance $\lfloor (\log t)^{3/2} \rfloor$, then back to the ancestor $a(u)$ of u situated at distance $\lfloor (\log t)^{3/2} \rfloor$ from u , then back again. Decomposing on the hittings of the root, we can majorate this probability by

$$\sum_{s \leq t} \mathbb{P}_T(X_t = e) \sum_{k=\lfloor (\log t)^{3/2} \rfloor}^t \sum_{u \in T_k} \mathbb{P}_T(H_u < H_e) \mathbb{P}_T^{a(u)}(H_u < t),$$

where H_u stands for the hitting time of u . Using the fact that the conductance from 0 to u is bounded by C_u , the probability we are considering is at most

$$n \sum_{k=\lfloor (\log t)^{3/2} \rfloor}^t \sum_{u \in T_k} C_u \mathbb{P}_T^{a(u)}(H_u < t).$$

Denoting by $C(v \rightarrow u)$ the conductance between v and u , we have easily

$$\mathbb{P}_T^v(H_u < t) < t \frac{C(v \rightarrow u)}{\pi(v)} < c_1 t \frac{C_u}{C_v}.$$

As a direct consequence of Theorem 4.1, we have

$$\begin{aligned} E_{\text{MT}} \left[\sum_{u \in T_k} C_u \mathbb{P}_T^{a(u)}(H_u < t) \right] &\leq c_1 t^2 E_{\text{MT}} \left[\sum_{u \in T_k} C_u \frac{C_u}{C_{a(u)}} \right] \\ &\leq c_1 t^2 \left(E_q \left[\sum A_i \exp(\log(A_i)) \right] \right)^{\lfloor (\log t)^{3/2} \rfloor} \leq c_1 t^2 \rho(2)^{\lfloor (\log t)^{3/2} \rfloor}. \end{aligned}$$

The result follows by an application of the Borel-Cantelli Lemma.

Proof of (38) : The proof is quite similar to the precedent argument, summing over the different $T^{(v_i)}$.

Proof of (39) : using $\kappa > 5$, by Lemma 4.2 we can find an $\varepsilon > 0$ such that IMT- almost surely the sequence $n^{3/2+\varepsilon}\pi(\mathbf{A}_n^\varepsilon)$ is summable, thus bounded, so there exists a constant $C'(T)$ such that for each n , $C_{e \rightarrow \mathbf{A}_n^\varepsilon} \leq C'(T)/n^{3/2+\varepsilon}$. Recalling from the proof of (30) the definition of $L_0(t)$, and $T_\varepsilon(t)$ we have

$$\mathbb{P}_T(X_t \in \mathbf{A}_{t^\alpha}^\varepsilon; t \leq T_\varepsilon(t)) \leq \mathbb{E}_T[L_0(T_\varepsilon(t))]C'(T)/t^{\alpha(3/2+\varepsilon)} \leq t^{1/2+\varepsilon'-\alpha(1+\varepsilon)}, \quad (45)$$

where ε' can be chosen arbitrarily close to 0. By choosing α close enough to 1/2, the result follows easily, using (43).

Proof of (40) : we recall from (22) the definition of the sets \mathbf{B}_n^ε . By the same argument as in the proof of Lemma 4.5, we get

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(X_s \in \mathbf{B}_{t^\alpha}^\varepsilon \text{ for some } s \leq t) \leq H_{\lfloor t^{1/2+\varepsilon} \rfloor} \sum_{i=0}^{\lfloor t^{1/2+\varepsilon} \rfloor} U_i^{t^\alpha},$$

with $H_t = 1 + \sum_{j=0}^{t-1} \prod_{k=j-1}^t A(v_k)$, and $U_i^{t^\alpha}$ is the probability to get to $B_{t^\alpha}^\varepsilon$ during one excursion in T^{v_i} . By the same argument as in the proof of Lemma 4.2, we get that, almost surely, there exists a constant $C''(T)$ such that

$$H_t \leq C''(T)t^{1/7},$$

whence

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(X_s \in \mathbf{B}_{t^\alpha}^\varepsilon \text{ for some } s \leq t) \leq C''(T)t^{1/7} \sum_{i=0}^{\lfloor t^{1/2+\varepsilon} \rfloor} U_i^{t^\alpha}.$$

Then, denoting $\sum_{i=0}^\infty U_i^t t^{1+\varepsilon'} := E_i$, the E_i are i.i.d. variables (under IMT) with finite expectation for ε' small enough and $U_i^t < \frac{1}{t^{3/2}}E_i$. Then the result follows, using the law of large numbers.

Proof of (33) and (34): Note that, under MT, the random variables $\eta_i - \tau_i$ are i.i.d.. On the other hand, as a consequence of (2), for some constant $\nu_0 > 0$,

$$\mathbb{P}_{\text{MT}}[\eta_i - \tau_i > x] \geq \nu_0 \mathbb{P}_{\text{MT}}[T_0 > x],$$

Where T_0 is the first return to the root. We recall from (42) that

$$T^\varepsilon(t) = \min\{t : |X_s| \geq t^{1/2+\varepsilon}\}$$

Then, following the proof of Lemma 10 of [21], we have,

$$\mathbb{P}_T[T_0 > t] \geq \mathbb{P}_T[T_0 > T^{\varepsilon/2}(t)]\mathbb{P}_T[T^{\varepsilon/2}(t) \geq t | T_0 > T^\varepsilon(t)]. \quad (46)$$

As a consequence of (44), for some constant depending on the tree $C_3(T)$,

$$\mathbb{P}_T[T_0 > T^{\varepsilon/2}(t)] > C_3(T)t^{-1/2-\varepsilon/2}.$$

On the other hand,

$$\mathbb{P}_T[T^{\epsilon/2}(t) < t | T_0 > T^{\epsilon/2}(t)] \leq \frac{\mathbb{P}_T[T^{\epsilon/2}(t) < t]}{\mathbb{P}_T[T_0 > T^{\epsilon/2}(t)]} \leq C_4(T)t^{1/2+\epsilon}e^{-t^{\epsilon/2}},$$

MT–almost surely, using (43) and the previous estimate. We get then that almost surely, for t large enough (the “enough” depending on T),

$$\mathbb{P}_T[T^{\epsilon/2}(t) > n | T_0 > T^{\epsilon/2}(t)] > 1/2.$$

Therefore for some positive constant $C_5(T)$,

$$\mathbb{P}_T[T_0 > t] \geq C_5(T)t^{-1/2-\epsilon/2}.$$

We deduce by taking the expectation that

$$\mathbb{P}_{\text{MT}}[T_0 > t] \geq C_{22}t^{-1/2-\epsilon/2},$$

for some positive and deterministic constant C_{22} . Now

$$\mathbb{P}_{\text{MT}} \left[\sum_{i=1}^{\lfloor t^{1/2+\epsilon} \rfloor} \eta_i - \tau_i < t \right] \leq \left(1 - \nu_0 C_{22} t^{-1/2-\epsilon/2} \right)^{t^{1/2+\epsilon}} \leq e^{-C_{23}t^{\epsilon/2}}.$$

An application of the Borel-Cantelli Lemma finishes the proof of (33) and (34). This finishes the proof of Lemma 5.1.

We now turn to our last part, namely the annealed central limit theorem. The proof has many parts in common with the proof in the quenched case, so we feel free to refer to the previous part.

7 Proof of Theorem 1.3.

We recall from section 3 the definition of the “environment seen from the particle $T_t = \theta^v(T)$ ”. As for the quenched case, we will first show a central limit theorem on IMT trees, then in a second part we will use the coupling to deduce the result for MT trees

7.1 The annealed CLT on IMT trees

We will first show the following proposition :

Proposition 7.1. *Suppose $N(e) \geq 1$, $q - a.s.$, (2). If $p = 1$, $\rho'(1) < 0$ and $\kappa \in (2, \infty]$, then there is a deterministic constant $\sigma > 0$ such that, under \mathbb{P}_{IMT} , the process $\{h(X_{\lfloor nt \rfloor})/\sqrt{\sigma^2 n}\}$ converges in law to a standard Brownian motion, as n goes to infinity.*

Remark : This result is of great theoretical interest, as it is the only context where we are able to cover the whole case $\kappa > 2$, we could actually make the proof of Theorem 1.3 without this proposition, but as it has an interest in itself, we give the proof in the general case.

Proof : Let, as in the quenched setting, $0 < \delta < 1$ and ρ_t be a random variable, independent of the walk, uniformly chosen in $[t, t + t^\delta]$. We recall from (7) the definition of $S_x, x \in T$ and from (8) the definition of η . We are going to show the following

Lemma 7.1. *Under the assumptions of Theorem 1.3,*

$$\frac{S_{X_{\rho_t}}}{h(X_{\rho_t})} \rightarrow \eta, \quad (47)$$

in probability.

We admit for the moment this lemma and finish the proof of Proposition 7.1. We have

$$h(X_{\rho_t}) = \frac{h(X_{\rho_t})}{S_{X_{\rho_t}}} S_{X_{\rho_t}} = \eta S_{X_{\rho_t}} + \left[\frac{h(X_{\rho_t})}{S_{X_{\rho_t}}} - \eta \right] S_{X_{\rho_t}}.$$

The first term converges to a Brownian motion with variance σ , by the same arguments as in the quenched setting, while the second one is a $o(S_{X_{\rho_t}})$. The result then follows easily, using the same arguments as in the proof of Theorem 1.2.

We now turn to the proof of Lemma 7.1. The proof is quite similar to the proof of Proposition 4.2: we take some small $\epsilon > 0$, then we estimate the number of visits to the points in B_n^ϵ during one excursion in T^{v_i} , and estimate the number of such excursion before time n . We rely on the following lemma, similar to Lemma 4.2

Lemma 7.2. *Suppose that the assumptions of Theorem 1.3 are true. Then for $1 < \lambda < \kappa - 1 \wedge 2$, and $n > 0$, there exists some constant C'_1 such that*

$$E_{\text{MT}} \left[\sum_{x \in A_n^\epsilon} C_x \right] < C'_1 n^{-(\lambda-1)}.$$

Proof : the proof relies on the same ideas as the proof of Proposition 4.2. First recall that, for n large enough,

$$\begin{aligned} E_{\text{MT}} \left[\sum_{x \in A_n^\epsilon} C_x \right] &\leq P_{\widehat{\text{MT}}_n} \left[\left| S_{v_n} - E_{\widehat{\text{MT}}_n} [S_{v_n} | \tilde{F}_n^*] \right| > \frac{n\epsilon}{4} \right] \\ &\quad + P_{\widehat{\text{MT}}_n} \left[\left| E_{\widehat{\text{MT}}_n} [S_{v_n} | \tilde{F}_n^*] - E_{\widehat{\text{MT}}_n} [S_{v_n}] \right| > \frac{n\epsilon}{4} \right] := P_1 + P_2. \end{aligned}$$

To bound P_1 , we recall that, under the law $\widehat{\text{MT}}_n^*$,

$$S_{v_n} - E_{\widehat{\text{MT}}_n} [S_{v_n} | \tilde{F}_n^*] = \sum_{i=0}^n \tilde{W}_i^* B_i,$$

where W_i are centered and independent random variables with bounded moments of order $\lambda + 1$ and

$$B_j = \sum_{k=0}^j \prod_{i=k+1}^j A_i.$$

Using Inequality 2.6.20 from page 82 of [23], we obtain that, for some constant C_2

$$E \left[\left| S_{v_n} - E_{\widehat{\text{MT}}_n^*} [S_{v_n} | \tilde{F}_n^*] \right|^\lambda \right] < C_2 \sum_{k=0}^n E[B_k^\lambda].$$

Then, using the same arguments as in the proof of Proposition 4.2, we get that $E[B_k^\lambda]$ is bounded independently of n and k , whence

$$E \left[\left| S_{v_n} - E_{\widehat{\text{MT}}_n^*} [S_{v_n} | \tilde{F}_n^*] \right|^\lambda \right] < C_3 n.$$

Using Markov's Inequality, there exists C_4 such that

$$P_1 < \frac{C_4}{2} n^{-(\lambda-1)}. \quad (48)$$

On the other hand, recalling (16),

$$\left| E_{\widehat{\text{MT}}_n^*} [S_{v_n} | \tilde{F}_n^*] - E_{\widehat{\text{MT}}_n^*} [S_{v_n}] \right| < C_5 + \left| \sum_{k=1}^n \tilde{A}_k D_k g(A_{n+1}) (1 + \rho + \rho^2 + \dots + \rho^{k-1}) \right|,$$

where C_5 is a finite constant and

$$D_k = \sum_{j=k+1}^n \prod_{i=k+1}^j A_i g(A_{j+1}),$$

where g is a bounded function. We recall that

$$N_k := \sum_{j=n-k}^n \tilde{A}_j D_j (1 + \rho + \rho^2 + \dots + \rho^{j-1})$$

is a martingale with respect to the filtration $\mathcal{A}_k = \sigma(A_j, n-k \leq j \leq n)$, whence, using Burkholder's Inequality,

$$E_{\widehat{\text{MT}}_n^*} [(N_n)^\lambda] \leq C_6 E_{\widehat{\text{MT}}_n^*} \left[\left(\sum_{i=0}^n (D_i)^2 \right)^{\lambda/2} \right].$$

We recall that $1 < \lambda < (\kappa - 1) \wedge 2$, whence, by concavity, the last expression is bounded above by

$$C_6 E_{\widehat{\text{MT}}_n^*} \left[\sum_{i=0}^n (D_i)^\lambda \right] < C_7 n.$$

Therefore, using Markov's Inequality, we get that

$$P_2 < n^{1-\lambda}.$$

This, together with (48), finishes the proof of Lemma 7.2.

We now finish the proof of Lemma 7.1. Let us go back to IMT trees. We recall the definition of the sets \mathbf{B}_n^ϵ :

$$\mathbf{B}_n^\epsilon = \left\{ v \in T, d(v, \text{Ray}) = n, \left| \frac{S_v^{\text{Ray}}}{n} - \eta \right| > \epsilon \right\}. \quad (49)$$

We are going to prove that

$$\lim_{t \rightarrow \infty} \mathbb{P}_T(X_{\rho_t} \in \cup_{n=1}^{\infty} \mathbf{B}_n^\epsilon) = 0, \text{ IMT - a.s..}$$

We introduce $\gamma > 1/2$, and recall the definition of the event

$$\Gamma_t = \{\exists u \leq 2t | X_u = v_{\lfloor t^\gamma \rfloor}\}.$$

It is easy to see, using the same arguments as in the proof of Lemma 4.5, that

$$P_{\text{IMT}}[\Gamma_t] \xrightarrow{t \rightarrow \infty} 0.$$

Furthermore, we introduce the event

$$\Gamma'_t = \{\exists 0 \leq u \leq t, d(X_u, \text{Ray}) > n^\gamma\};$$

then it is a direct consequence of Lemma 4.6 that

$$P_{\text{IMT}}[\Gamma'_t] \xrightarrow{t \rightarrow \infty} 0.$$

As for the quenched case, we have

$$\begin{aligned} \mathbb{P}_{\text{IMT}}(X_{\rho_t} \in \cup_{m=1}^{\infty} \mathbf{B}_m^\epsilon) &\leq \mathbb{P}_{\text{IMT}}(X_{\rho_t} \in \cup_{m=1}^{n^\gamma} \mathbf{B}_m^\epsilon; \Gamma_t^c \cap \Gamma'_t{}^c) + \mathbb{P}_{\text{IMT}}(\Gamma_t) + \mathbb{P}_{\text{IMT}}(\Gamma'_t) \\ &\leq \frac{1}{\lfloor t^\delta \rfloor} E_{\text{IMT}} \left[\mathbb{E}_T \left[\sum_{s=0}^{H_{v_{\lfloor t^\gamma \rfloor}}} \mathbb{1}_{X_s \in \cup_{m=1}^{n^\gamma} \mathbf{B}_m^\epsilon} \right] \right] + o(1), \end{aligned} \quad (50)$$

where $H_{v_{\lfloor t^\gamma \rfloor}}$ is the first time the walk hits $v_{\lfloor t^\gamma \rfloor}$.

We recall that $T^{(v_i)}$ the subtree constituted of the vertices $x \in T$ such that $v_i \leq x$, $v_{i-1} \not\leq x$, the same computations as in the proof of Lemma 4.5 imply

$$\mathbb{P}_{\text{IMT}}(X_{\rho_t} \in \cup_{m=1}^{\infty} \mathbf{B}_m^\epsilon) \leq \frac{1}{\lfloor t^\delta \rfloor} E_{\text{IMT}} \left[\sum_{i=0}^{\lfloor t^\gamma \rfloor} \mathbb{E}_T \left[\sum_{s=0}^{H_{v_{\lfloor t^\gamma \rfloor}}} \mathbb{1}_{X_s = v_i} \right] \tilde{N}_i \right], \quad (51)$$

Where \tilde{N}_i is the \mathbb{P}_T -expectation of the number of visits to $\cup_{m=1}^{n^\delta} \mathbf{B}_m^\epsilon \cap T^{(v_i)}$ during one excursion in $T^{(v_i)}$. Lemma 7.2, and the method of 4.3 imply that, under IMT conditioned on $\{\text{Ray}, A(v_i)\}$, \tilde{N}_i are independent and identically distributed variables, with expectation at most equal to $C'_1 \sum_{i=0}^{n^\gamma} i^{1-\lambda}$ for some $\lambda > 1$. By choosing γ close enough to 0, we get $E_{\text{IMT}}[\tilde{N}_i | \{\text{Ray}, A(v_i)\}] \leq C'_1 n^{1/2-\epsilon}$ for some $\epsilon > 0$. We recall that

$$\mathbb{E}_T \left[\sum_{s=0}^{H_{v_{\lfloor t^\gamma \rfloor}}} \mathbb{1}_{X_s = v_i} \right] \leq C'' \left(1 + \sum_{j=0}^{\lfloor t^\gamma \rfloor - 1} \prod_{k=j-1}^{\lfloor t^\gamma \rfloor} A(v_k) \right).$$

The latter expression has bounded expectation under IMT, as an easy consequence of Statement 4.1 and Lemma 3.2.

We deduce that

$$\mathbb{P}_{\text{IMT}}(X_{\rho_t} \in \cup_{m=1}^{\infty} \mathbf{B}_m^{\varepsilon}) \leq C_5 n^{\frac{1}{2} - \varepsilon + \gamma - \delta}.$$

Since γ can be chosen as close to $1/2$ as needed, the exponent can be taken lower than 0. The end of the proof is then completely similar to the quenched case.

7.2 The annealed CLT on MT trees.

We now turn to the proof of Theorem 1.3. We use the coupling and the notations presented in section 5. Our main proposition in this part will be the following:

Proposition 7.2. *Under the assumptions of Theorem 1.3, for some $\alpha < 1/2$*

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{MT}}(\Delta_t \neq \Delta_t^{\alpha}) = 0, \tag{52}$$

and

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{IMT}}(\tilde{\Delta}_t \neq \tilde{\Delta}_t^{\alpha}) = 0. \tag{53}$$

Further, under MT,

$$\limsup \frac{\Delta_t}{t} = 0, \tag{54}$$

and under IMT,

$$\limsup \frac{\tilde{\Delta}_t}{t} = 0. \tag{55}$$

Finally, under IMT,

$$\limsup \frac{\mathbf{B}_t}{\sqrt{t}} = 0. \tag{56}$$

(Here \limsup denotes the limit in law.)

Before proving the latter proposition, we introduce some technical estimates, whose proof will be postponed.

Lemma 7.3. For all $\epsilon > 0$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{MT}} \left(\sum_{i=1}^{t^{1/2+\epsilon}} (\eta_i - \tau_i) < t \right) = 0, \quad (57)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{IMT}} \left(\sum_{i=1}^{t^{1/2+\epsilon}} (\tilde{\eta}_i - \tilde{\tau}_i) < t \right) = 0, \quad (58)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{MT}}(\exists k \leq I_t, \Theta_{i-1}, \Theta_i \in J_k, |Y_{\Theta_i}| > |Y_{\Theta_{i-1}}|) = 0, \quad (59)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{IMT}}(\exists k \leq I_t, \tilde{\Theta}_{i-1}, \tilde{\Theta}_i \in \tilde{J}_k, d(\check{Y}_{\tilde{\Theta}_i}, \text{Ray}) > d(\check{Y}_{\tilde{\Theta}_{i-1}}, \text{Ray})) = 0, \quad (60)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{MT}}(X_s \in \cup_{k=t^\alpha - (\log t)^2}^{t^\alpha} \mathbf{A}_k^\epsilon \text{ for some } s \leq t) = 0, \quad (61)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{IMT}}(X_s \in \cup_{k=t^\alpha - (\log t)^2}^{t^\alpha} \mathbf{B}_k^\epsilon \text{ for some } s \leq t) = 0. \quad (62)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{MT}}(W_{X_s} > t^{1/4-\epsilon} \text{ for some } 0 \leq s \leq t) = 0 \quad (63)$$

$$\lim_{t \rightarrow \infty} \mathbb{P}_{\text{IMT}}(W_{X_s} > t^{1/4-\epsilon} \text{ for some } 0 \leq s \leq t) = 0 \quad (64)$$

We now turn to the proof of (52). As a consequence of (57) and (59) that, with \mathbb{P}_{MT} probability approaching 1 as n goes to infinity,

$$\mathbf{t}(\Theta_{2t^{1/2+\epsilon}}) > t,$$

whence, using Lemmas 7.3 and 4.6,

$$\begin{aligned} & \lim_{t \rightarrow \infty} \mathbb{P}_{\text{MT}} \left(\max_{s \in \cup_{k=1}^{I_t} J_k} |Y_s| \geq t^\alpha \right) \\ & \leq \limsup_{t \rightarrow \infty} \sum_{i=0}^{2t^{1/2+\epsilon}} \mathbb{P}_{\text{MT}}(\exists j > i : |Y_{\Theta_j}| \geq t^\alpha - (\log t)^2, Y_{\Theta_i} = e, \\ & S_{Y_{\Theta_j}} \geq (\eta - \epsilon_1)t^\alpha/2, |Y_{\Theta_k}| > 0, \forall i < k \leq j; W_{X_s} \leq t^{1/4-\epsilon} \forall 0 \leq s \leq t \\ & |S_{X_s} - |X_s|| \leq \epsilon t^{1/4-\epsilon'} |X_s|, \forall s \leq t) := \limsup_{t \rightarrow \infty} \sum_{i=1}^{2t^{1/2+\epsilon}} \mathbb{P}_{i,t}; \end{aligned}$$

where ϵ, ϵ_1 are positive numbers that can be chosen arbitrarily small. We recall that the process $\{N_s\} = \{S_{X_{\theta_i+s \wedge K_t}}\}$ is a supermartingale. and that there exists a previsible and non-decreasing process A_s such that $N_s + A_s$ is a martingale. Furthermore, on the event $\{W_{X_s} \leq t^{1/4-\epsilon} \forall 0 \leq s \leq t\}$, the increments of this martingale are bounded above by $t^{1/2-\epsilon}$. Azuma's Inequality implies the result, as in the quenched case.

The proof of (53) is similar and omitted.

We recall that in the proofs of (30),(31) and (32) we only used the assumption $\kappa > 5$, therefore the proof of (54),(55) and (56) are direct consequence, by dominated convergence.

We now turn to the proof of Lemma 7.3. The proofs of (57), (58), (59), (60) and (61) follow directly from equations (33), (34), (37), (38) and (39), whose proofs did not use any assumption other than $\kappa > 5$, by dominated convergence.

To prove (62), note that, similarly to the proof of 7.1,

$$\begin{aligned} \mathbb{P}_{\text{IMT}}(X_s \in \cup_{k=t^\alpha - (\log t)^2}^{t^\alpha} \mathbf{B}_k^\epsilon \text{ for some } s \leq t) \\ = E_{\text{IMT}} \left[\sum_{i=0}^{\lfloor t^\gamma \rfloor} \mathbb{E}_T \left[\sum_{s=0}^{H_{v_{\lfloor t^\gamma \rfloor}}} \mathbb{1}_{X_s=v_i} \right] N'_i \right], \end{aligned}$$

where N'_i is the \mathbb{P}_T -expectation of the number of visits to $\cup_{k=t^\alpha - (\log t)^2}^{t^\alpha} \mathbf{B}_k^\epsilon \cap T^{(v_i)}$ during one excursion in $T^{(v_i)}$. Lemma 4.2 and the method of Lemma 4.3 imply that, under IMT conditioned on $\{\text{Ray}, A(v_i)\}$, N'_i are independent and identically distributed variables, up to a bounded constant, with expectation at most equal to $C'(\log t)^2 t^{-\alpha(\lambda-1)}$ for some $\lambda > 2$. We also recall that

$$\mathbb{E}_T \left[\sum_{s=0}^{H_{v_{\lfloor t^\gamma \rfloor}}} \mathbb{1}_{X_s=v_i} \right] \leq C'' \left(1 + \sum_{j=0}^{\lfloor t^\gamma \rfloor - 1} \prod_{k=j-1}^{\lfloor t^\gamma \rfloor} A(v_k) \right).$$

has bounded expectation under IMT, as an easy consequence of Statement 4.1 and Lemma 3.2. By choosing γ close enough to 0 and α close to 1, we get the result.

The proofs of (63) and (64) are easily deduced from the proofs of (35) and (36), the only difference being that we do not need to apply the Borel-Cantelli Lemma.

Acknowledgement: We would like to warmly thank Y. Hu for his help and encouragement.

References

- [1] E. Aidekon. Transient random walks in random environment on a Galton–Watson tree. *Probability Theory and Related Fields*, 142(3):525–559, 2008. MR2438700
- [2] B. Bercu and A. Touati. Exponential inequalities for self-normalized martingales with applications. *Annals of Applied Probability*, 18(5):1848–1869, 2008. MR2462551
- [3] J.D. Biggins. Martingale convergence in the branching random walk. *Journal of Applied Probability*, pages 25–37, 1977. MR0433619
- [4] J.D. Biggins and A.E. Kyprianou. Seneta-Heyde norming in the branching random walk. *Annals of Probability*, 25(1):337–360, 1997. MR1428512
- [5] P. Billingsley. *Convergence of Probability measures*. Wiley, 1999. MR1700749
- [6] A.A. Chernov. Replication of a multicomponent chain, by the lightning mechanism. *Biophysics*, 12(2):336–341, 1967.
- [7] G. Faraud, Y. Hu, and Z. Shi. An almost sure convergence for stochastically biased random walk on a Galton-Watson tree . <http://arxiv.org/abs/1003.5505>, 2010.

- [8] Y. Hu and Z. Shi. A subdiffusive behaviour of recurrent random walk in random environment on a regular tree. *Probability theory and related fields*, 138(3):521–549, 2007. MR2299718
- [9] Y. Hu and Z. Shi. Slow movement of random walk in random environment on a regular tree. *Annals of Probability*, 35(5):1978–1997, 2007. MR2349581
- [10] Y. Hu and Z. Shi. Minimal position and critical martingale convergence in branching random walks, and directed polymers on disordered trees. *Annals of Probability*, 37(2):742–789, 2009. MR2510023
- [11] J.G. Kemeny, A.W. Knapp, and J.L. Snell. *Denumerable Markov Chains*. 2nd ed. Springer, 1976. MR0407981
- [12] C. Kipnis and S.R.S. Varadhan. Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions. *Communications in Mathematical Physics*, 104(1):1–19, 1986. MR0834478
- [13] Q. Liu. On generalized multiplicative cascades. *Stochastic processes and their applications*, 86(2):263–286, 2000. MR1741808
- [14] Q. Liu. Asymptotic properties and absolute continuity of laws stable by random weighted mean. *Stochastic processes and their applications*, 95(1):83–107, 2001. MR1847093
- [15] R. Lyons. The Ising model and percolation on trees and tree-like graphs. *Communications in Mathematical Physics*, 125:337–353, 1989. MR1016874
- [16] R. Lyons and R. Pemantle. Random walks in a random environment and first-passage percolation on trees. *Annals of Probability*, 20:125–136, 1992. MR1143414
- [17] R. Lyons and Y. Peres. *Probability on trees and networks*. 2005.
- [18] B. Mandelbrot. Multiplications aleatoires iterees et distributions invariantes par moyenne ponderee aleatoire: quelques extensions. *Comptes rendus de l'académie des Sciences*, 278:355–358, 1974. MR0431352
- [19] M.V. Menshikov and D. Petritis. On random walks in random environment on trees and their relationship with multiplicative chaos. *Mathematics and computer science II (Versailles, 2002)*, pages 415–422, 2002. MR1940150
- [20] J. Neveu. Arbres et processus de Galton-Watson. *Annales de l'Institut H. Poincare*, 22(2):199–207, 1986. MR0850756
- [21] Y. Peres and O. Zeitouni. A central limit theorem for biased random walks on Galton–Watson trees. *Probability Theory and Related Fields*, 140(3):595–629, 2008. MR2365486
- [22] V.V. Petrov. *Sums of Independent Random Variables*. Springer-Verlag, 1975. MR0388499
- [23] V.V. Petrov. *Limit theorems of probability theory: sequences of independent random variables*. Oxford University Press, USA, 1995. MR1353441
- [24] O. Zeitouni. Lecture notes on random walks in random environment. Ecole d'été de probabilités de Saint-Flour 2001. *Lecture Notes in Mathematics*, 1837:189–312, 2003. MR2071631