

Stochastic order methods applied to stochastic travelling waves

Roger Tribe and Nicholas Woodward*

Abstract

This paper considers some one dimensional reaction diffusion equations driven by a one dimensional multiplicative white noise. The existence of a stochastic travelling wave solution is established, as well as a sufficient condition to be in its domain of attraction. The arguments use stochastic ordering techniques.

Key words: travelling wave, stochastic order, stochastic partial differential equations.

AMS 2000 Subject Classification: Primary 60H15.

Submitted to EJP on July 8, 2010, final version accepted January 13, 2011.

*Department of Mathematics, University of Warwick, CV4 7AL, UK. Corresponding author: r.p.tribe@warwick.ac.uk.

1 Introduction

We consider the following one dimensional reaction diffusion equation, driven by a one dimensional Brownian motion:

$$du = u_{xx} dt + f(u) dt + g(u) \circ dW. \quad (1)$$

We shall assume throughout that

$$f, g \in C^3([0, 1]) \quad \text{and} \quad f(0) = f(1) = g(0) = g(1) = 0, \quad (2)$$

and consider solutions whose values $u(t, x)$ lie in $[0, 1]$ for all $x \in R$ and $t \geq 0$. The noise term $g(u) \circ dW$ models the fluctuations due to an associated quantity that affects the entire solution simultaneously (for example temperature effects). In this setting we consider modelling with a Stratonovich integration to be more natural, as we can consider it as the limit of smoother noisy drivers. The use of a non-spatial noise allows us the considerable simplification of considering solutions that are monotone functions on R .

We consider three types of reaction f . We call f : (i) of *KPP* type if $f > 0$ on $(0, 1)$ and $f'(0), f'(1) \neq 0$; (ii) of *Nagumo* type if there exists $a \in (0, 1)$ with $f < 0$ on $(0, a)$, $f > 0$ on $(a, 1)$ and $f'(0), f'(a), f'(1) \neq 0$; (iii) of *unstable* type if there exists $a \in (0, 1)$ with $f > 0$ on $(0, a)$, $f < 0$ on $(a, 1)$ and $f'(0), f'(a), f'(1) \neq 0$. The deterministic behaviour (that is when $g = 0$) is well understood (see Murray [7] chapter 13 for an overview). Briefly, for f of Nagumo type there is a unique travelling wave, for f of KPP type a family of travelling waves, and for f of unstable type one expects solutions that split into two parts, one travelling right and one left, with a large flatish region inbetween around the level a . For f of KPP or Nagumo type the solution starting at the initial condition $H(x) = I(x < 0)$ converges towards the slowest travelling wave. Various sufficient conditions (and in Bramson [1] necessary and sufficient conditions) are known on other initial conditions that guarantee the solutions converge to a travelling wave.

The aim of this paper is to start to investigate a few of these results for the stochastic equation (1). There are many tools used in the deterministic literature. In this paper, we develop only the key observation that the deterministic solutions started from the Heavyside initial condition $H(x) = I(x < 0)$ become more stretched over time. The most transparent way to view this, as explained in Fife and McLeod [2], is in phase space, where it corresponds to a comparison result. More precisely, the corresponding phase curves $p(t, u)$, defined via $u_x(t, x) = p(t, u(t, x))$, are increasing in time. This idea is exploited extensively in [2] and subsequent papers. For the stochastic equation (1), the solution paths are not almost surely increasing. However we will use similar arguments to show that the solutions are stochastically ordered, and that this is an effective substitute.

To state our results we briefly describe a state space for solutions. Our state space will be

$$\mathcal{D} = \{\text{decreasing, right continuous } \phi : R \rightarrow [0, 1] \text{ satisfying } \phi(-\infty) = 1, \phi(\infty) = 0\}.$$

We will use a wavefront marker defined, for a fixed $a \in (0, 1)$, by

$$\Gamma(\phi) = \inf\{x : \phi(x) < a\} \quad \text{for } \phi \in \mathcal{D}. \quad (3)$$

To center the wave at its wavefront we define

$$\tilde{\phi}(x) = \phi(\Gamma(\phi) + x) \quad \text{for } \phi \in \mathcal{D}. \quad (4)$$

We call $\tilde{\phi}$ the wave ϕ centered at height a . We have suppressed the dependence on a in the notation for the wavefront marker and the centered wave. We give \mathcal{D} the $L^1_{loc}(R)$ topology. We write $\mathcal{M}(\mathcal{D})$ for the space of (Borel) probability measures on \mathcal{D} with the topology of weak convergence of measures.

A stochastic travelling wave is a solution $u = (u(t) : t \geq 0)$ to (1) with values in \mathcal{D} and for which the centered process $(\tilde{u}(t) : t \geq 0)$ is a stationary process with respect to time. The law of a stochastic travelling wave is the law of $\tilde{u}(0)$ on \mathcal{D} . We will show (see section 3.3) that the centered solutions themselves form a Markov process. Then an equivalent definition is that the law of a stochastic travelling wave is an invariant measure for the centered process.

The hypotheses for our results below are stated in terms of the drift f_0 in the equivalent Ito integral formulation, namely

$$du = u_{xx} dt + f_0(u) dt + g(u) dW \quad \text{where } f_0 = f + \frac{1}{2} g g'.$$

While we suspect that the existence, uniqueness and domains of stochastic travelling waves are determined by the Stratonovich drifts f , our methods use the finiteness of certain moments and require assumptions about f_0 . It is easy to find examples where the type of f and f_0 can be different, for example f of KPP type and f_0 of Nagumo type, or f of Nagumo type and f_0 of unstable type.

We now state our main results. The framework for describing stretching on \mathcal{D} is explained in section 2.3, where we define a pre-order on \mathcal{D} that reflects when one element is more stretched than another, and where we also recall the ideas of stochastic ordering for laws on a metric space equipped with a pre-order. These ideas are exploited to deduce the convergence in law in the following theorem, which is proved in section 3.

Theorem 1. *Suppose that f_0 is of KPP, Nagumo or unstable type. In the latter two cases suppose that $f_0(a) = 0$ and $g(a) \neq 0$. Let u be the solution to (1) started from $H(x) = I(x < 0)$. Then the laws $\mathcal{L}(\tilde{u}(t))$ are stochastically increasing (for the stretching pre-order on \mathcal{D} - see 2.3), and converge to a law $\nu \in \mathcal{M}(\mathcal{D})$. Furthermore ν is the law of a stochastic travelling wave.*

Note that the unstable type reactions are therefore stabilized by the noise. This becomes intuitive when one realizes that a large flatish patch near the level a will be destroyed by the noise since $g(a) \neq 0$.

It is an immediate consequence of the stochastic ordering that for any solution whose initial condition $u(0)$ is stochastically less stretched than ν , the laws $\mathcal{L}(\tilde{u}(t))$ will also converge to ν (that is they are in the domain of attraction of ν - see Proposition 16). It is not clear how to check whether an initial condition has this property. However, our stochastic ordering techniques do yield a simple sufficient condition, albeit for not quite the result one would want and also not in the unstable case, as described in the following theorem which is proved in section 4.

Theorem 2. *Suppose that f_0 is of KPP or Nagumo type, and in the latter two case suppose that $f_0(a) = 0$ and $g(a) \neq 0$. let u be the solution to (1) with initial condition $u(0) = \phi \in \mathcal{D}$ which equals 0 for all sufficiently positive x and equals 1 for all sufficiently negative x . Then*

$$\frac{1}{t} \int_0^t \mathcal{L}(\tilde{u}(s)) ds \rightarrow \nu \quad \text{as } t \rightarrow \infty$$

where ν is the law of the stochastic travelling wave from Theorem 1.

2 Preliminaries, including stretching and stochastic ordering

2.1 Regularity and moments for solutions

We state a theorem that summarizes the properties of solutions to (1) that we require. Recall we are assuming the hypothesis (2). A mild solution is one satisfying the semigroup formulation.

Theorem 3. *Let W be an (\mathcal{F}_t) Brownian motion defined on a filtered space $(\Omega, (\mathcal{F}_t), \mathcal{F}, P)$ where \mathcal{F}_0 contains the P null sets. Given any $u_0 : R \times \Omega \rightarrow [0, 1]$ that is $\mathcal{B}(R) \times \mathcal{F}_0$ measurable, there exists a progressively measurable mild solution $(u(t, x) : t \geq 0, x \in R)$ to (1), driven by W and with initial condition u_0 . The paths of $(u(t) : t \geq 0)$ lie almost surely in $C([0, \infty), L^1_{loc}(R))$ and solutions are pathwise unique in this space. If P_ϕ is the law on $C([0, \infty), L^1_{loc}(R))$ of the solution started at ϕ , then the family $(P_\phi : \phi \in L^1_{loc}(R))$ form a strong Markov family. The associated Markov semigroup is Feller (that is it maps bounded continuous functions on L^1_{loc} into bounded continuous functions).*

There is a regular version of any solution, where the paths of $(u(t, x) : t > 0, x \in R)$ lie almost surely in $C^{0,3}((0, \infty) \times R)$. The following additional properties hold for such regular versions.

(i) *Two solutions u, v with initial conditions satisfying $u_0(x) \leq v_0(x)$ for all $x \in R$, almost surely, remain coupled, that is $u(t, x) \leq v(t, x)$ for all $t \geq 0$ and $x \in R$, almost surely.*

(ii) *If $u_0 \in \mathcal{D}$, almost surely, then $u(t) \in \mathcal{D}$ for all $t \geq 0$, almost surely. Moreover, $u(t, x) > 0$ and $u_x(t, x) < 0$ for all $t > 0$ and $x \in R$, almost surely.*

(iii) *For $0 < t_0 < T$ and $L, p > 0$*

$$E [|u_x(t, x)|^p + |u_{xx}(t, x)|^p + |u_{xxx}(t, x)|^p] \leq C(p, t_0, T) \quad \text{for } x \in R \text{ and } t \in [t_0, T]$$

and hence (as explained below)

$$E \left[\sup_{|x| \leq L} |u_x(t, x)|^p + \sup_{|x| \leq L} |u_{xx}(t, x)|^p \right] \leq C(p, t_0, T)(L + 1) \quad \text{for } t \in [t_0, T].$$

(iv) *Suppose the initial condition $u_0 = \phi$ satisfies $\phi(x) = 0$ for all large x and $\phi(x) = 1$ for all large $-x$. Then for $0 < t_0 < T$ and $p, \eta > 0$*

$$E [|u_x(t, x)|^p + |u_{xx}(t, x)|^p] \leq C(\phi, \eta, p, t_0, T)e^{-\eta|x|} \quad \text{for } x \in R \text{ and } t \in [t_0, T]$$

and hence (as explained below)

$$\sup_{t \in [t_0, T]} E \left[\sup_x |u_x(t, x)|^p \right] < \infty.$$

Moreover, $|u_x(t, x)| \rightarrow 0$ as $|x| \rightarrow \infty$, almost surely, for $t > 0$.

Remark. Henceforth, all results refer to the regular versions of solutions, that is with paths in $C([0, \infty), L^1_{loc}(R)) \cap C^{0,3}((0, \infty) \times R)$ almost surely.

The results in Theorem 3 are mostly standard, and we omit the proofs but give a few comments for some of the arguments required. The moments in parts (iii) and (iv) at fixed (t, x) can be

established via standard Green's function estimates, though a little care is needed since we allow arbitrary initial conditions. Indeed the constants for the p th moment of the k th derivative blow up like $t_0^{-pk/2}$ (as for the deterministic equation), though we shall not use this fact. One can then derive all the bounds on the supremum of derivatives by bounding them in terms of integrals of a higher derivative and using the pointwise estimates. For example, in part (iv),

$$\sup_x |u_x(t, x)|^p \leq C(p, \eta) \left(|u_x(t, 0)|^p + \int_R e^{+\eta|x|} |u_{xx}(t, y)|^p dy \right).$$

The supremum over $[-L, L]$ in part (iii) can be bounded by a sum of suprema over intervals $[k, k+1]$ of length one, and each of these bounded using higher derivatives. This leads to the dependency $L + 1$ in the estimate, which we do not believe is best possible but is sufficient for our needs.

One route to reach the strict positivity and strict negativity in part (ii) is to follow the argument in Shiga [11]. In [11] Theorem 1.3, there is a method to show that $u(t, x) > 0$ for all $t > 0, x \in R$ for an equation as in (1) but where the noise is space-time white noise. However the proof applies word for word for an equation driven by a single noise once the basic underlying deviation estimate in [11] Lemma 4.2 is established. This method applies to the equation

$$dv = v_{xx} dt + f'(u)v dt + g'(u)v \circ dW.$$

for the derivative $v = u_x$ over any time interval $[t_0, \infty)$. This yields the strict negativity $u_x(t, x) < 0$ for all $t > 0, x \in R$, almost surely, (which of course implies the strict positivity of u). The underlying large deviation estimate is for

$$P \left[|N(t, x)| \geq \epsilon e^{-(T-t)|x|} \text{ for some } t \leq T/2, x \in R \right]$$

where $N(t, x) = \int_0^t \int G_{t-s}(x - y)b(s, y) dy dW_s$ is the stochastic part of the Green's function representation for $u(t, x)$. This estimate can also be derived using the method suggested in Shiga, where he appeals to an earlier estimate in Lemma 2.1 of Mueller [6]. The method in [6], based on dyadic increments as in the Levy modulus for Brownian motion, can also be applied without significant changes to our case, since it reduces to estimates on the quadratic variation of increments of $N(t, x)$ and these are all bounded (up to a constant) for our case by the analogous expressions in the space-time white noise case.

2.2 Wavefront markers, and pinned solutions

We remark on the L_{loc}^1 topology on \mathcal{D} . First, the space \mathcal{D} is Polish. Indeed, for $\phi_n, \phi \in \mathcal{D}$, the convergence $\phi_n \rightarrow \phi$ is equivalent to the convergence of the associated measures $-d\phi_n \rightarrow -d\phi$ in the weak topology on the space of finite measures on R . Note that using the Prohorov metric for this weak convergence gives a compatible metric on \mathcal{D} that is translation invariant, in that $d(\phi, \psi) = d(\phi(\cdot - a), \psi(\cdot - a))$ for any a . Second, the convergence $\phi_n \rightarrow \phi$ is equivalent to $\phi_n(x) \rightarrow \phi(x)$ for almost all x (indeed $\phi_n(x) \rightarrow \phi(x)$ provided that x is a continuity point of ϕ).

The wave marker Γ , defined by (3), is upper semicontinuous on \mathcal{D} . The wavemarker $\Gamma(u(t))$ and the centered solution $\tilde{u}(t, x)$ are semi-martingales for $t \geq t_0 > 0$ and $x \in R$. Here is the calculus behind this.

Lemma 4. Let u be a solution to (1) with $u(0) \in \mathcal{D}$ almost surely. For $t > 0$ let $m(t, \cdot)$ denote the inverse function for the map $x \rightarrow u(t, x)$. Then the process $(m(t, x) : t > 0, x \in (0, 1))$ lies in $C^{0,3}((0, \infty) \times (0, 1))$, almost surely. For each $x \in (0, 1)$ and $t_0 > 0$, the process $(m(t, x) : t \geq t_0)$ satisfies

$$dm(t, x) = \frac{m_{xx}(t, x)}{m_x^2(t, x)} dt - f(x)m_x(t, x) dt - g(x)m_x(t, x) \circ dW_t. \quad (5)$$

Also $\Gamma(u(t)) = m(t, a)$ and the centered process \tilde{u} solves, for $t \geq t_0$,

$$d\tilde{u} = \tilde{u}_{xx} dt + f(\tilde{u}) dt + g(\tilde{u}) \circ dW + \tilde{u}_x \circ d\Gamma(u). \quad (6)$$

Proof The (almost sure) existence and regularity of m follow from Theorem 3 (noting that $x \rightarrow u(t, x)$ is strictly decreasing for $t > 0$ by Theorem 3 (ii)). The equation for $m(t, x)$ would follow by chain rule calculations if W were a smooth function. To derive it using stochastic calculus we choose $\phi : (0, 1) \rightarrow \mathbb{R}$ smooth and compactly supported and develop $\int m(t, x)\phi(x) dx$. To shorten the upcoming expressions, we use, for real functions h_1, h_2 defined on an interval with R , the notation $\langle h_1, h_2 \rangle$ for the integral $\int h_1(x)h_2(x)dx$ over this interval, whenever it is well defined. Using the substitution $x \rightarrow u(t, x)$ we have, for $t > 0$,

$$\langle m(t), \phi \rangle = \int_0^1 m(t, x)\phi(x) dx = \int_R x\phi(u(t, x))u_x(t, x) dx = \langle \phi(u(t))u_x(t), x \rangle$$

(where we write x for the identity function). Expanding $\phi(u(t, x))u_x(t, x)$ via Itô's formula we obtain

$$\begin{aligned} d\langle m, \phi \rangle &= d\langle \phi(u)u_x, x \rangle \\ &= \langle \phi(u)(u_{xxx} + f'(u)u_x), x \rangle dt + \langle \phi_x(u)(u_{xx} + f(u))u_x, x \rangle dt \\ &\quad + \langle \phi(u)g'(u)u_x + \phi_x(u)g(u)u_x, x \rangle \circ dW. \end{aligned}$$

To assist in our notation we let $\hat{u}, \hat{u}_x, \hat{u}_{xx}, \dots$ denote the composition of the maps $x \rightarrow u, u_x, u_{xx}$ with the map $x \rightarrow m(t, x)$ (e.g. $\hat{u}_x(t, x) = u_x(t, m(t, x))$). Using this notation we have, for $x \in (0, 1), t > 0$,

$$\hat{u}(t, x) = x, \quad \hat{u}_x m_x = 1, \quad \hat{u}_{xx} m_x^2 + \hat{u}_x m_{xx} = 0. \quad (7)$$

We continue by using the reverse substitution $x \rightarrow m(t, x)$ to reach

$$\begin{aligned} d\langle m, \phi \rangle &= \langle \hat{u}_{xxx} m m_x + f' m, \phi \rangle dt + \langle \hat{u}_{xx} m + f m, \phi_x \rangle dt \\ &\quad + \langle g' m, \phi \rangle \circ dW + \langle g m, \phi_x \rangle \circ dW \\ &= -\langle \hat{u}_{xx} m_x + f m_x, \phi \rangle dt - \langle g m_x, \phi \rangle \circ dW \\ &= \langle (m_{xx}/m_x^2) - f m_x, \phi \rangle dt - \langle g m_x, \phi \rangle \circ dW. \end{aligned}$$

In the second equality we have integrated by parts. In the final equality we have used the identities in (7). This yields the equation for m . The decomposition for \tilde{u} follows by applying the Itô-Ventzel formula (see Kunita [3] section 3.3) using the decompositions for $du(t, x)$ and $d\Gamma(u(t)) = dm(t, a)$. \square

2.3 Stretching and stochastic stretching

Definitions. For $\phi : R \rightarrow R$ we set


$$\theta_0(\phi) = \inf\{x : \phi(x) > 0\},$$

where we set $\inf\{\emptyset\} = \infty$. We write $\tau^a \phi$ for the translated function $\tau^a \phi(\cdot) = \phi(\cdot - a)$. For $\phi, \psi : R \rightarrow R$ we say that

ϕ crosses ψ if $\phi(x) \geq \psi(x)$ for all $x \geq \theta_0(\phi - \psi)$,

ϕ is more stretched than ψ if $\tau^a \phi$ crosses ψ for all $a \in R$.

We write $\phi \succ \psi$ to denote that ϕ is more stretched than ψ , and as usual we write $\phi \prec \psi$ when $\psi \succ \phi$. In the diagram below, we plot a wave ϕ and two of its translates, all three curves crossing another wave ψ .



Remarks

1. Any function $\phi \in \mathcal{D}$ is more stretched than the Heavyside function $H(x) = I(x < 0)$. If $\lambda > 1$ and $\phi \in \mathcal{D}$ then ϕ is more stretched than $x \rightarrow \phi(\lambda x)$.
2. The upcoming lemma shows that the relation $\phi \succ \psi$ is quite natural. For $\phi \in \mathcal{C}^1 \cap \mathcal{D}$ with $\phi_x < 0$, one can associate a phase curve $p^\phi : (0, 1) \rightarrow R$ defined by $\phi_x(x) = p^\phi(\phi(x))$. The relation of stretching between two such functions becomes simple comparison between the associated phase curves. Another way to define the relation for functions in \mathcal{D} is to define it on such nice paths via comparison of the associated phase curves, and then take the smallest closed extension.
3. It is useful for us to have a direct definition of stretching without involving the associated phase curves. For example the key Lemma 7 below uses this direct definition. Moreover, in a future work, we will treat the case of spatial noise, where solutions do not remain decreasing and working in phase space is difficult. Note that Lemma 7 applies when functions are not necessarily decreasing.
4. We will show that the stretching relation is a pre-order on \mathcal{D} , which means that it is reflexive ($\phi \succ \phi$) and transitive ($\phi \succ \psi$ and $\psi \succ \rho$ imply that $\phi \succ \rho$). We recall that a partial order would in addition satisfy the anti-symmetric property: $\phi \succ \psi$ and $\psi \succ \phi$ would imply that $\phi = \psi$.

Lemma 5. (i) The relation $\phi \succ \psi$ is a pre-order on \mathcal{D} .

(ii) If $\phi \succ \psi$ then $\tau^a \phi \succ \psi$ and $\phi \succ \tau^a \psi$ for all a . Moreover, if $\phi \succ \psi$ and $\psi \succ \phi$ then $\phi = \tau^a \psi$ for some a .

(iii) The relation $\phi \succ \psi$ is closed, that is the set $\{(\phi, \psi) \in \mathcal{D}^2 : \phi \succ \psi\}$ is closed in the product topology.

(iv) Suppose that $\phi, \psi \in \mathcal{D} \cap \mathcal{C}^1(R)$ and that $\phi_x, \psi_x < 0$. Write ϕ^{-1}, ψ^{-1} for the inverse functions on $(0, 1)$. Then $\phi \succ \psi$ if and only if $\phi_x^{-1} \leq \psi_x^{-1}$ if and only if $p^\phi \geq p^\psi$.

(v) Suppose that $\phi, \psi \in \mathcal{D} \cap \mathcal{C}^1(R)$ and that $\phi_x, \psi_x < 0$. If $\phi \succ \psi$ and $\phi(x) = \psi(x)$ then $\phi(y) \geq \psi(y)$ for all $y \geq x$ and $\phi(y) \leq \psi(y)$ for all $y \leq x$.

Proof We start with parts (iv) and (v), which use only straightforward calculus (and are exploited in [2]). Fix $\phi, \psi \in \mathcal{D} \cap \mathcal{C}^1(R)$ satisfying $\phi_x, \psi_x < 0$. Suppose $\phi \succ \psi$ and $z \in (0, 1)$. We may choose unique x_1, x_2 so that $\phi(x_1) = \psi(x_2) = z$. Let $a = x_2 - x_1$. Then if $x_2 \geq \theta_0(\tau^a \phi - \psi)$

$$\begin{aligned} \phi_x(x_1) &= \lim_{h \downarrow 0} h^{-1}(\phi(x_1 + h) - \phi(x_1)) \\ &= \lim_{h \downarrow 0} h^{-1}(\tau^a \phi(x_2 + h) - z) \\ &\geq \lim_{h \downarrow 0} h^{-1}(\psi(x_2 + h) - z) = \psi_x(x_2), \end{aligned}$$

while if $x_2 \leq \theta_0(\tau^a \phi - \psi)$

$$\begin{aligned} \phi_x(x_1) &= \lim_{h \downarrow 0} h^{-1}(\phi(x_1) - \phi(x_1 - h)) \\ &= \lim_{h \downarrow 0} h^{-1}(z - \tau^a \phi(x_2 - h)) \\ &\geq \lim_{h \downarrow 0} h^{-1}(z - \psi(x_2 - h)) = \psi_x(x_2). \end{aligned}$$

This shows that $p^\phi(z) \geq p^\psi(z)$. This is equivalent to $\phi_x^{-1}(z) \leq \psi_x^{-1}(z)$ since $p^\phi(z)\phi_x^{-1}(z) = 1$. Conversely suppose that $p^\phi \geq p^\psi$ on $(0, 1)$. Write ϕ^{-1}, ψ^{-1} for the inverse functions. Then for $0 < z_1 < z_2 < 1$

$$\phi^{-1}(z_2) - \phi^{-1}(z_1) = \int_{z_1}^{z_2} \phi_x^{-1}(x) dx \leq \int_{z_1}^{z_2} \psi_x^{-1}(x) dx = \psi^{-1}(z_2) - \psi^{-1}(z_1).$$

If $\phi(x_0) = \psi(x_0) = z_0$ then the above implies that

$$\begin{cases} \phi^{-1}(z) \leq \psi^{-1}(z) & \text{for } z \geq z_0, \\ \phi^{-1}(z) \geq \psi^{-1}(z) & \text{for } z \leq z_0, \end{cases} \quad \text{and} \quad \begin{cases} \phi(x) \geq \psi(x) & \text{for } x \geq x_0, \\ \phi(x) \leq \psi(x) & \text{for } x \leq x_0. \end{cases} \quad (8)$$

This proves part (v). We claim that ϕ crosses ψ . If $x_0 = \theta_0(\phi - \psi) \in (-\infty, \infty)$ then the assumed regularity of ϕ, ψ implies that $\phi(x_0) = \psi(x_0)$ and (8) shows that $\phi(x) \geq \psi(x)$ for $x \geq x_0$. If $\theta_0(\phi - \psi) = -\infty$ then $\phi(x) \geq \psi(x)$ for all x (since if $\phi(x_1) < \psi(x_1)$ we may choose $x_0 < x_1$ with $\phi(x_0) = \psi(x_0)$ which contradicts (8)). This shows that ϕ crosses ψ . Since $p^{\tau^a \phi} = p^\phi$ we may apply the same argument to $\tau^a \phi$ to conclude that $\phi \succ \psi$ completing the proof of part (iv).

For part (iii) suppose that $\phi_n \succ \psi_n$ for $n \geq 1$ and $\phi_n \rightarrow \phi, \psi_n \rightarrow \psi$. Let $E \subseteq R$ be the nullset off which $\phi_n(x) \rightarrow \phi(x), \psi_n(x) \rightarrow \psi(x)$. Suppose $\theta_0(\phi - \psi) < \infty$. We may choose $x \in E^c$, arbitrarily close to $\theta_0(\phi - \psi)$, satisfying $\phi(x) > \psi(x)$. Then $\phi_n(x) > \psi_n(x)$ and hence $\theta_0(\phi_n - \psi_n) \leq x$ for large n . We deduce that $\limsup \theta_0(\phi_n - \psi_n) \leq \theta_0(\phi - \psi)$. Now if $x \in E^c$ and $x > \theta_0(\phi - \psi)$ then

$x > \theta_0(\phi_n - \psi_n)$ for large n and so $\phi(x) \geq \psi(x)$. By the right continuity of ϕ, ψ we have that ϕ crosses ψ . We may repeat this argument for $\tau^a \phi, \psi$ to deduce that $\phi \succ \psi$.

For part (i) we write ϕ_ϵ for the convolution $\phi * G_\epsilon$ with the Gaussian kernel $G_\epsilon(x) = (4\pi\epsilon)^{-1/2} \exp(-x^2/4\epsilon)$. Note that $\phi_\epsilon \rightarrow \phi$ almost everywhere and hence also in \mathcal{D} . Suppose that $\phi \succ \rho \succ \psi$. It will follow from Lemma 7 below, in the special case that $f = g = 0$, that $\phi_\epsilon \succ \rho_\epsilon \succ \psi_\epsilon$. Moreover $\phi_\epsilon, \rho_\epsilon, \psi_\epsilon \in \mathcal{D} \cap \mathcal{C}^1(\mathbb{R})$ and $(\phi_\epsilon)_x, (\rho_\epsilon)_x, (\psi_\epsilon)_x < 0$. Using part (iv) we find that $\phi_\epsilon \succ \psi_\epsilon$ and by part (iii) that $\phi \succ \psi$, completing the proof of transitivity. Reflexivity follows from $\theta_0(\phi - \phi) = \infty$.

The first statement in part (ii) is immediate from the definition. Suppose $\phi \succ \psi \succ \phi$. Then, as in part (i), $\phi_\epsilon \succ \psi_\epsilon \succ \phi_\epsilon$ and so $\tilde{\phi}_\epsilon \succ \tilde{\psi}_\epsilon \succ \tilde{\phi}_\epsilon$. Since $\tilde{\phi}_\epsilon(0) = \tilde{\psi}_\epsilon(0) = a$ it follows as in (8) (with the choice $x_0 = 0, z_0 = a$) that $\tilde{\phi}_\epsilon = \tilde{\psi}_\epsilon$. This implies that $\phi_\epsilon = \tau^{a_{\epsilon}} \psi_\epsilon$ and $\phi = \tau^a \psi$ for some a_ϵ, a . \square

Notation. For two probability measures $\mu, \nu \in \mathcal{M}(\mathcal{D})$ we write $\mu \stackrel{s}{\succ} \nu$ if μ is stochastically larger than ν , where we take the stretching pre-order on \mathcal{D} .

Notation. For a measure $\mu \in \mathcal{M}(\mathcal{D})$ we define the centered measure $\tilde{\mu}$ as the image of μ under the map $\phi \rightarrow \tilde{\phi}$.

Remark 5. We recall here the definition of stochastic ordering. A function $F : \mathcal{D} \rightarrow \mathbb{R}$ is called increasing if $F(\phi) \geq F(\psi)$ whenever $\phi \succ \psi$. Then $\mu \stackrel{s}{\succ} \nu$ means that $\int_{\mathcal{D}} F(\phi) d\mu \geq \int_{\mathcal{D}} F(\phi) d\nu$ for all non-negative, measurable, increasing $F : \mathcal{D} \rightarrow \mathbb{R}$. An equivalent definition is that there exists a pair of random measures X, Y (with values in $\mathcal{M}(\mathcal{D})$), defined on a single probability space and satisfying $X \succ Y$ almost surely. The equivalence is sometimes called Strassen's theorem, and is often stated for partial orders, but holds when the relation is only a pre-order on a Polish space. Indeed, there is an extension to countably many laws: if $\mu_1 \stackrel{s}{\prec} \mu_2 \stackrel{s}{\prec} \dots$ then there exist variables $(U_n : n \geq 1)$ with $\mathcal{L}(U_n) = \mu_n$ and

$$U_1 \prec U_2 \prec \dots \quad \text{almost surely.} \quad (9)$$

See Lindvall [4] for these results (where a mathscinet review helps by clarifying one point in the proof).

Lemma 6. (i) The relation $\mu \stackrel{s}{\succ} \nu$ is a pre-order on $\mathcal{M}(\mathcal{D})$.

(ii) If $\mu \stackrel{s}{\succ} \nu$ then $\tilde{\mu} \stackrel{s}{\succ} \tilde{\nu}$ and $\mu \stackrel{s}{\succ} \tilde{\nu}$. Moreover, if $\nu \stackrel{s}{\succ} \mu$ and $\mu \stackrel{s}{\succ} \nu$ then $\tilde{\mu} = \tilde{\nu}$.

(iii) The relation $\mu \stackrel{s}{\succ} \nu$ is closed, that is the set $\{(\mu, \nu) : \mu \stackrel{s}{\succ} \nu\}$ is closed in the product topology on $(\mathcal{M}(\mathcal{D}))^2$.

Proof For part (i) note that the transitivity follows from Strassen's theorem (9) and the transitivity for \succ on \mathcal{D} . For part (ii) suppose that $\mu \stackrel{s}{\succ} \nu \stackrel{s}{\succ} \mu$. Then by (9) we may pick variables so that $\mathcal{L}(X) = \mathcal{L}(Z) = \mu, \mathcal{L}(Y) = \nu$ and $X \succ Y \succ Z$ almost surely. The smoothed and then centered variables satisfy $\tilde{X}_\epsilon \succ \tilde{Y}_\epsilon \succ \tilde{Z}_\epsilon$ almost surely. Note that $\tilde{X}_\epsilon(0) = \tilde{Y}_\epsilon(0) = \tilde{Z}_\epsilon(0) = a$ almost surely. Applying (8) we find that

$$\tilde{X}_\epsilon(x) \geq \tilde{Y}_\epsilon(x) \geq \tilde{Z}_\epsilon(x) \text{ for } x \geq 0 \quad \text{and} \quad \tilde{X}_\epsilon(x) \leq \tilde{Y}_\epsilon(x) \leq \tilde{Z}_\epsilon(x) \text{ for } x \leq 0.$$

But \tilde{X}_ϵ and \tilde{Z}_ϵ have the same law and this implies that $\tilde{X}_\epsilon = \tilde{Z}_\epsilon$ almost surely. Undoing the smoothing shows that $\tilde{X} = \tilde{Z}$ almost surely and hence $\tilde{\mu} = \tilde{\nu}$.

For part (iii) we suppose that $\mu_n \xrightarrow{s} \nu_n$ and $\mu_n \rightarrow \mu$, $\nu_n \rightarrow \nu$. Choose pairs (X_n, Y_n) with $\mathcal{L}(X_n) = \mu_n$ and $\mathcal{L}(Y_n) = \nu_n$ and $X_n \succ Y_n$ almost surely. The laws of (X_n, Y_n) are tight so that we may find a subsequence n_k and versions $(\hat{X}_{n_k}, \hat{Y}_{n_k})$ that converge almost surely to a limit (X, Y) . Now pass to the limit as $k \rightarrow \infty$ to deduce that $\mu \succ \nu$. \square

3 Existence of the stochastic travelling wave

3.1 The solution from $H(x) = I(x < 0)$ stretches stochastically

It is straightforward to extend the basic stretching lemma from McKean [5] to deterministic equations with time dependent reactions, as follows. Since it plays a key role in this paper, we present the proof with the small changes that are needed.

Lemma 7. *Consider the deterministic heat equation*

$$u_t(t, x) = u_{xx}(t, x) + H(u(t, x), t, x), \quad t > 0, x \in R,$$

where $H : [0, 1] \times [0, T] \times R \rightarrow R$ is a measurable function that is Lipschitz in the first variable, uniformly over (t, x) .

Suppose u and v are mild solutions, taking values in $[0, 1]$, and $u, v \in \mathcal{C}^{1,2}([0, T] \times R)$. Suppose that $u(0)$ crosses $v(0)$. Then $u(t)$ crosses $v(t)$ at all $t \in [0, T]$.

Proof Consider $w : [0, T] \times R \rightarrow [-1, 1]$ defined as $w = u - v$. Define

$$R(t, x) = \left(\frac{H(u(t, x), t, x) - H(v(t, x), t, x)}{u(t, x) - v(t, x)} \right) \mathcal{I}(u(t, x) \neq v(t, x)).$$

R is bounded and w is a mild solution to $w_t = w_{xx} + wR$. We now wish to exploit a Feynman-Kac representation for w . Let $(B(t) : t \geq 0)$ be a Brownian motion, time scaled so that its generator is the Laplacian, and defined on a filtered probability space $(\Omega, (\mathcal{F}_s : s \geq 0), (P_x : x \in R))$, where under P_x , B starts at x . Fix $t > 0$. Then for $s \in [0, t)$,

$$M(s) = w(t - s, B(s)) \exp \left(\int_0^s R(t - r, B(r)) dr \right)$$

is a continuous bounded (\mathcal{F}_s) martingale and hence has an almost sure limit $M(t)$ as $s \uparrow t$. As $s \uparrow t$ one has $w(t - s, B(s)) \rightarrow w(0, B(t))$ (use the fact that $w(r, x) \rightarrow w(0, x)$ as $r \downarrow 0$ for almost all x). For any (\mathcal{F}_s) stopping time τ satisfying $\tau \leq t$ we obtain from $E_x[M(0)] = E_x[M(\tau \wedge s)]$ by letting $s \uparrow t$, that

$$w(t, x) = E_x[M(\tau)] = E_x \left[w(t - \tau, B(\tau)) \exp \left(\int_0^\tau R(t - r, B(r)) dr \right) \right]. \quad (10)$$

Suppose $w(t, x_1) > 0$ for some x_1 , in particular $x_1 \geq \theta_0(w(t))$. Consider the stopping time $\tau = \inf_{0 \leq s \leq t} \{s : |M(s)| = 0\} \wedge t$. Then $E_{x_1}[M(\tau)] = E_{x_1}[M(t)I(\tau = t)] = w(t, x_1) > 0$. From this we can

construct a deterministic continuous path $(\xi(s) : s \in [0, t])$ such that $\xi(0) = x_1$ and $w(t-s, \xi(s)) > 0$ for $0 \leq s \leq t$. Now take $x_2 > x_1$. Consider another stopping time defined by

$$\tau^* = \inf_{0 \leq s \leq t} \{s : B(s) = \xi(s)\} \wedge t.$$

We claim $M(\tau^*) \geq 0$ almost surely under P_{x_2} . Indeed, on $\{\tau^* < t\}$ this is immediate from the construction of ξ . On $\{\tau^* = t\}$ we have $B(\tau^*) = B(t) \geq \xi(t)$. Since $w(0, \xi(t)) > 0$ we know that $\xi(t) \geq \theta_0(u(0) - v(0))$ and the assumption that $u(0)$ crosses $v(0)$ ensures that $w(0, B(t)) \geq 0$ and hence $M(\tau^*) \geq 0$. Applying (10), with $x = x_2$ and τ replaced by τ^* , we find that $w(t, x_2) \geq 0$ when $x_2 \geq x_1$. If $\theta_0(u(t) - v(t)) \in (-\infty, \infty)$ then we can choose x_1 arbitrarily close to $\theta_0(u(t) - v(t))$ and the proof is finished. In the cases $\theta_0(u(t) - v(t)) = -\infty$ we may pick x_1 arbitrarily negative and in the case $\theta_0(u(t) - v(t)) = +\infty$ there is nothing to prove. \square

By using a Wong-Zakai result for approximating the stochastic equation (1) by piecewise linear noises, we shall now deduce the following stretching lemma for our stochastic equations with white noise driver.

Proposition 8. *Suppose that u, v are two solutions to (1) with respect to the same Brownian motion. Then, for all $t > 0$,*

- (i) *if $u(0)$ crosses $v(0)$ almost surely then $u(t)$ crosses $v(t)$ almost surely;*
- (ii) *if $u(0), v(0) \in \mathcal{D}$ and $u(0) \succ v(0)$ almost surely then $u(t) \succ v(t)$ almost surely.*

Proof Define a piecewise linear approximation to a Brownian motion W by, for $\epsilon > 0$,

$$W_t^\epsilon = W_{k\epsilon} + \epsilon^{-1}(t - k\epsilon)(W_{(k+1)\epsilon} - W_{k\epsilon}) \quad \text{for } t \in [k\epsilon, (k+1)\epsilon] \text{ and } k = 0, 1, \dots$$

Then the equation

$$\frac{du^\epsilon}{dt} = u_{xx}^\epsilon + f(u^\epsilon) + g(u^\epsilon) \frac{dW^\epsilon}{dt}, \quad u^\epsilon(0) = u(0)$$

can be solved successively over each interval $[k\epsilon, (k+1)\epsilon]$, path by path. If u solves (1) with respect to W then we have the convergence

$$u^\epsilon(t, x) \rightarrow u(t, x) \quad \text{in } L^2 \text{ for all } x \in R, t \geq 0.$$

We were surprised not to be able to find such a result in the literature that covered our assumptions. The closest papers that we found were [8], whose assumptions did not cover Nemitski operators for the reaction and noise, and [12], which proves convergence in distribution for our model on a finite interval. Nevertheless this Wong-Zakai type result is true and can be established by closely mimicking the original Wong-Zakai proof for stochastic ordinary differential equations. The details are included in section 2.6 of the thesis [13]. (We note that the proof there, which covers exactly equation (1), would extend easily to equations with higher dimensional noises. Also it is in this proof that the hypothesis that f, g have continuous third derivatives is used.)

In a similar way we construct v^ϵ with $v^\epsilon(0) = v(0)$. For all k , all paths of u^ϵ and v^ϵ lie in $C^{1,2}([k\epsilon, (k+1)\epsilon] \times R)$. By applying Lemma 7 repeatedly over the intervals $[k\epsilon, (k+1)\epsilon]$ we see that $u^\epsilon(t)$ crosses $v^\epsilon(t)$ for all $t \geq 0$ along any path where $u(0)$ crosses $v(0)$. We must check that this is preserved in the limit. Fix $t > 0$. There exists $\epsilon_n \rightarrow 0$ so that for almost all paths

$$u^{\epsilon_n}(t, x) \rightarrow u(t, x) \quad \text{and} \quad v^{\epsilon_n}(t, x) \rightarrow v(t, x) \quad \text{for all } x \in Q.$$

Fix such a path where in addition $u(0)$ crosses $v(0)$. Suppose that $\theta_0(u(t) - v(t)) < \infty$. Arguing as in part (iii) of Lemma 5 we find that $\limsup_{n \rightarrow \infty} \theta_0(u^{\epsilon_n}(t) - v^{\epsilon_n}(t)) \leq \theta_0(u(t) - v(t))$. Now choose $y \in Q$ with $y > \theta_0(u(t) - v(t))$. Taking n large enough that $y > \theta_0(u^{\epsilon_n}(t) - v^{\epsilon_n}(t))$ we find, since $u^{\epsilon_n}(t)$ crosses $v^{\epsilon_n}(t)$, that $u^{\epsilon_n}(t, y) \geq v^{\epsilon_n}(t, y)$. Letting $n \rightarrow \infty$ we find $u(t, y) \geq v(t, y)$. Now the continuity of the paths ensures that $u(t)$ crosses $v(t)$. For part (ii) it remains to check that $\tau^a u(t)$ crosses $v(t)$. But this follows from part (i) after one remarks that if u solves (1) then so too does $(\tau^a u(t) : t \geq 0)$. \square

Corollary 9. (i) Let u, v be solutions to (1) satisfying $\mathcal{L}(u(0)) \succ^s \mathcal{L}(v(0))$. Then $\mathcal{L}(u(t)) \succ^s \mathcal{L}(v(t))$ for all $t \geq 0$.

(ii) Let u be the solution to (1) started from $H(x) = I(x < 0)$. Then $\mathcal{L}(\tilde{u}(t)) \succ^s \mathcal{L}(\tilde{u}(s))$ for all $0 \leq s \leq t$.

Proof For part (i), we may by Strassen's theorem find versions $\bar{u}(0)$ and $\bar{v}(0)$ that satisfy $\bar{u}(0) \succ \bar{v}(0)$ almost surely. The result then follows by from Lemma 8 (ii) (and uniqueness in law of solutions).

For part (ii) we shall, when $\mu \in \mathcal{M}(\mathcal{D})$, write Q_t^μ for the law of $u(t)$ for a solution u to (1) whose initial condition $u(0)$ has law μ . We write \tilde{Q}_t^μ for the centered law of $\tilde{u}(t)$. We write Q_t^H and \tilde{Q}_t^H in the special case that $\mu = \delta_H$. Since H is less stretched than any $\phi \in \mathcal{D}$ we know that $Q_s^H \succ^s Q_0^H = \delta_H$ for any $s \geq 0$. Now set $\mu = Q_s^H$ and apply part (i) to see that

$$Q_{t+s}^H = Q_t^\mu \stackrel{s}{\succ} Q_t^H$$

where the first equality is the Markov property of solutions. This shows that $t \rightarrow Q_t^H$ is stochastically increasing. By Lemma 6 (ii) the family $t \rightarrow \tilde{Q}_t^H$ is also increasing. \square

The stochastic monotonicity will imply the convergence in law of $\tilde{u}(t)$ on a larger space, as explained in the proposition below. Define

$$\mathcal{D}_c = \{\text{decreasing, right continuous } \phi : R \rightarrow [0, 1]\}. \quad (11)$$

Then D_c is a compact space under the L_{loc}^1 topology: given a sequence $\phi_n \in \mathcal{D}_c$ then along a suitable subsequence n' the limit $\lim_{n' \rightarrow \infty} \phi_{n'}(x)$ exists for all $x \in Q$; then $\phi_{n'} \rightarrow \phi$ where $\phi(x) = \lim_{y \downarrow x} \psi(y)$ is the right continuous regularization of $\psi(x) = \limsup_{n' \rightarrow \infty} \phi_{n'}(x)$.

Proposition 10. Let u be the solution to (1) started from $H(x) = I(x < 0)$. Then $\tilde{u}(t)$, considered as random variables with values in \mathcal{D}_c , converge in distribution as $t \rightarrow \infty$ to a limit law $\nu_c \in \mathcal{M}(\mathcal{D}_c)$.

Proof Choose $t_n \uparrow \infty$. Then by Strassen's Theorem (9) we can find \mathcal{D} valued random variables U_n with law $\mathcal{L}(U_n) = \mathcal{L}(\tilde{u}(t_n))$ and satisfying $U_1 \prec U_2 \prec \dots$ almost surely. Note that $U_n(0) = a$ and that U_n has continuous strictly negative derivatives (by Theorem 3 (ii)). The stretching pre-order, together with Lemma 5 (v), implies that almost surely

$$U_{n+1}(x) \geq U_n(x) \text{ for } x \geq 0 \quad \text{and} \quad U_{n+1}(x) \leq U_n(x) \text{ for } x \leq 0.$$

Thus the limit $\lim_{n \rightarrow \infty} U_n(x)$ exists, almost surely, and we set U to be the right continuous modification of $\limsup U_n$. This modification satisfies $U_n(x) \rightarrow U(x)$ for almost all x , almost surely. Hence $U_n \rightarrow U$ in \mathcal{D}_c , almost surely, and the laws $\mathcal{L}(\tilde{u}_{t_n})$ converge to $\mathcal{L}(U)$ in distribution. We set ν_c to

be the law $\mathcal{L}(U)$ on D_c . To show that $\mathcal{L}(\tilde{u}_t) \rightarrow \nu$ it suffices to show that the limit does not depend on the choice of sequence (t_n) . Suppose (s_n) is another sequence increasing to infinity. If (r_n) is a third increasing sequence containing all the elements of (s_n) and (t_n) then the above argument shows that $\mathcal{L}(\tilde{u}_{r_n})$ is convergent and hence the limits of $\mathcal{L}(\tilde{u}_{s_n})$ and $\mathcal{L}(\tilde{u}_{t_n})$ must coincide. \square

Remark We do not yet know that the limit ν_c is supported on \mathcal{D} . We must rule out the possibility that the wavefronts get wider and wider and the limit ν_c is concentrated on flat profiles. We do this by a moment estimate in the next section. Once this is known, standard Markovian arguments in section 3.3 will imply that $\nu = \nu_c|_{\mathcal{D}}$, the restriction to \mathcal{D} , is the law of a stochastic travelling wave.

3.2 A moment bound

We will require the following simple first moment bounds. Under hypothesis (2) we may choose K_1, K_2 so that

$$-K_1(1-x) \leq f_0(x) \leq K_2x \quad \text{for all } x \in [0, 1].$$

Lemma 11. *Let u be a solution to (1) with initial condition $u(0) = \phi \in \mathcal{D}$.*

(i) *For any $T > 0$ there exist $C(T) < \infty$ so that*

$$\int_{\mathbb{R}} E [|u(t, x) - \phi(x)|] dx \leq C(T) \left(1 + \int_{\mathbb{R}} \phi(1-\phi)(x) dx \right) \quad \text{for } t \leq T.$$

(ii) *When the initial condition is $H(x) = I(x < 0)$ we have for $x > 0$*

$$\begin{aligned} E[u(t, x)] &\leq \min\{1, e^{K_2 t} x^{-1} G_t(x)\}, \\ E[1 - u(t, -x)] &\leq \min\{1, e^{K_1 t} x^{-1} G_t(x)\}, \end{aligned}$$

and there exists $C(K_1, K_2, a) < \infty$ so that

$$E[|\Gamma(u(t))|] \leq C(K_1, K_2, a) + 2(K_1^{1/2} + K_2^{1/2})t \quad \text{for all } t \geq 0.$$

Proof For part (i) we may, by translating the solution if necessary, assume that ϕ crosses $1/2$ at the origin, that is $\phi(x) \leq 1/2$ for $x \geq 0$ and $\phi(x) > 1/2$ for $x < 0$. Taking expectations in (1) and applying $f_0(x) \leq K_2x$ we find that $E[u(t, x)]$ solves

$$(\partial/\partial t)E[u(t, x)] \leq \Delta E[u(t, x)] + K_2 E[u(t, x)].$$

This leads to

$$E[u(t, x)] \leq e^{K_2 t} \int_{\mathbb{R}} G_t(x-y)\phi(y) dy$$

where $G_t(x) = (4\pi t)^{-1/2} \exp(-x^2/4t)$. Hence for $t \leq T$

$$\begin{aligned} \int_0^\infty E[u(t, x)] dx &\leq e^{K_2 t} \int_0^\infty \int_R^\infty G_t(x-y) \phi(y) dy dx \\ &= e^{K_2 t} \left(\int_{-\infty}^0 \int_0^\infty + \int_0^\infty \int_0^\infty \right) G_t(x-y) \phi(y) dx dy \\ &\leq e^{K_2 t} \int_{-\infty}^0 \int_0^\infty G_t(x-y) dx dy + e^{K_2 t} \int_0^\infty \phi(y) \int_R^\infty G_t(x-y) dx dy \\ &\leq C(T) \left(1 + \int_0^\infty \phi(y) dy \right). \end{aligned}$$

The process $v(t, x) = 1 - u(t, -x)$ solves (1) with $f_0(u)$ and $g(u)$ replaced by $-f_0(1-v)$ and $-g(1-v)$. So the same argument estimates $E[1 - u(t, -x)]$ and yields the bound (with a possibly different constant depending on K_1)

$$\int_{-\infty}^0 E[(1-u)(t, x)] dx \leq C(T) \left(1 + \int_{-\infty}^0 (1-\phi)(y) dy \right).$$

Note that

$$\int_R^\infty |u(t, x) - \phi(x)| dx \leq \int_0^\infty u(t, x) + \phi(x) dx + \int_{-\infty}^0 (1-u)(t, x) + (1-\phi)(x) dx.$$

Combining this with the bounds above and also

$$\int_0^\infty \phi(x) dx + \int_{-\infty}^0 (1-\phi)(x) dx \leq 2 \int_R^\infty \phi(1-\phi)(x) dx$$

(which uses that ϕ crosses $1/2$ at the origin) completes the proof of part (i).

For part (ii) we have more explicit bounds. Use a Gaussian tail estimate to the bound for $x > 0$

$$E[u(t, x)] \leq e^{K_2 t} \int_x^\infty G_t(y) dy \leq 2tx^{-1} e^{K_2 t} G_t(x).$$

In some regions the estimate $E[u(t, x)] \leq 1$ is better. Since $\{\Gamma(u(t)) \geq x\} = \{u(t, x) \geq a\}$ almost

surely for $t > 0$, we have via Chebychev's inequality

$$\begin{aligned}
E[(\Gamma(u(t)))_+] &= \int_0^\infty P[\Gamma(u(t)) \geq x] dx \\
&= \int_0^\infty P[u(t, x) \geq a] dx \\
&\leq \int_0^{2K_2^{1/2}t} dx + \int_{2K_2^{1/2}t}^\infty a^{-1} E[u(t, x)] dx \\
&\leq 2K_2^{1/2}t + \int_{2K_2^{1/2}t}^\infty 2t(ax)^{-1} e^{K_1 t} x^{-1} G_t(x) dx \\
&\leq 2K_2^{1/2}t + a^{-1} e^{K_2 t} (4\pi t)^{-1/2} \int_{2K_2^{1/2}t}^\infty (x/2K_2 t) e^{-x^2/4t} dx \\
&= 2K_2^{1/2}t + (aK_2)^{-1} (4\pi t)^{-1/2}.
\end{aligned}$$

Similarly $\{\Gamma(u(t)) \leq -x\} = \{1 - u(t, -x) \geq 1 - a\}$ and the same argument yields the bound on $E[1 - u(t, -x)]$ and also $E[(\Gamma(u(t)))_-] \leq 2K_1^{1/2}t + (aK_1)^{-1} (4\pi t)^{-1/2}$. These estimates combine to control $E[|\Gamma(u(t))|]$ as desired for $t \geq 1$. A slight adjustment bounds the region $t \leq 1$. \square

We briefly sketch a simple idea from [5] for the deterministic equation $u_t = u_{xx} + u(1 - u)$ started at H , which we will adapt for our stochastic equation. The associated centered wave satisfies

$$\tilde{u}_t = \tilde{u}_{xx} + \tilde{u}_x \dot{\gamma} + \tilde{u}(1 - \tilde{u})$$

where γ_t is the associated wavefront marker. Integrating over $(-\infty, 0] \times [t_0, t]$, for some $0 < t_0 < t$ yields the estimate

$$\begin{aligned}
0 &\geq \int_{-\infty}^0 [\tilde{u}(t, x) - \tilde{u}(t_0, x)] dx \\
&= \int_{t_0}^t \tilde{u}_x(s, 0) ds - (1 - a)(\gamma_t - \gamma_0) + \int_{t_0}^t \int_{-\infty}^0 \tilde{u}(1 - \tilde{u})(s, x) dx ds.
\end{aligned}$$

This allows one, for example, to control the size of the back tail $\int_{-\infty}^0 (1 - \tilde{u})(t, x) dx$. Integrating over $[0, \infty)$ gives information on the front tail. The following lemma gives the analogous tricks for the stochastic equation.

Lemma 12. *Let u be the solution to (1) started from $H(x) = I(x < 0)$. Let \tilde{u} be the solution centered*

at height $a \in (0, 1)$. Then for $0 < t_0 < t$, almost surely,

$$\begin{aligned} & \int_0^\infty (\tilde{u}(t, x) - \tilde{u}(t_0, x)) dx \\ &= - \int_{t_0}^t \tilde{u}_x(s, 0) ds - a(\Gamma(u(t)) - \Gamma(u(t_0))) \\ & \quad + \int_{t_0}^t \int_0^\infty f_0(\tilde{u})(s, x) dx ds + \int_{t_0}^t \int_0^\infty g(\tilde{u})(s, x) dx dW_s, \end{aligned} \quad (12)$$

$$\begin{aligned} & \int_{-\infty}^0 (\tilde{u}(t, x) - \tilde{u}(t_0, x)) dx \\ &= \int_{t_0}^t \tilde{u}_x(s, 0) ds - (1-a)(\Gamma(u(t)) - \Gamma(u(t_0))) \\ & \quad + \int_{t_0}^t \int_{-\infty}^0 f_0(\tilde{u})(s, x) dx ds + \int_{t_0}^t \int_{-\infty}^0 g(\tilde{u})(s, x) dx dW_s. \end{aligned} \quad (13)$$

Proof Integrating (6) first over $[t_0, t]$ and then over $[0, U]$ we find

$$\begin{aligned} & \int_0^U (\tilde{u}(t, x) - \tilde{u}(t_0, x)) dx \\ &= \int_0^U \int_{t_0}^t \tilde{u}_{xx}(s, x) ds dx + \int_0^U \int_{t_0}^t \tilde{u}_x(s, x) \circ d\Gamma(u(s)) dx \\ & \quad + \int_0^U \int_{t_0}^t f_0(\tilde{u})(s, x) ds dx + \int_0^U \int_{t_0}^t g(\tilde{u})(s, x) dW_s dx \\ &= \int_{t_0}^t (\tilde{u}_x(s, U) - \tilde{u}_x(s, 0)) ds + \int_{t_0}^t (\tilde{u}(s, U) - a) \circ d\Gamma(u(s)) \\ & \quad + \int_{t_0}^t \int_0^U f_0(\tilde{u})(s, x) dx ds + \int_{t_0}^t \int_0^U g(\tilde{u})(s, x) dx dW_s. \end{aligned} \quad (14)$$

The interchange of integrals uses Fubini's theorem path by path for the first and third terms on the right hand side and a stochastic Fubini theorem for the second and fourth term (for example the result on p176 of [9] applies directly for the fourth term and also the second term after localizing at the stopping times $\sigma_n = \inf\{t \geq t_0 : \sup_{y \in [0, U]} |\tilde{u}_x(s, y)| \geq n\}$).

To prove the lemma we shall let $U \rightarrow \infty$ in each of the terms. Bound $|f(z)| \leq Cz(1-z)$ for some C . Using the first moment bounds from Lemma 11 (ii) we see that

$$\begin{aligned} & \int_0^t \int_R E[|f_0(u)(s, x)|] dx ds \\ & \leq C \int_0^t \int_R E[u(1-u)(s, x)] dx ds \\ & \leq C \int_0^t \int_0^\infty E[u(s, x)] dx ds + C \int_0^t \int_{-\infty}^0 E[(1-u(s, x))] dx ds < \infty. \end{aligned}$$

This gives the domination that justifies $\int_{t_0}^t \int_0^U f_0(\tilde{u})(s, x) dx ds \rightarrow \int_{t_0}^t \int_0^\infty f_0(\tilde{u})(s, x) dx ds$ almost surely. The first moments also imply

$$\begin{aligned} \int_0^t E \left[\left| \int_0^\infty u(s, x) dx \right|^2 \right] ds &= \int_0^t \int_0^\infty \int_0^\infty E[u(s, x)u(s, y)] dx dy ds \\ &\leq \int_0^t \int_0^\infty \int_0^\infty E[u(s, x)] \wedge E[u(s, y)] dx dy ds < \infty. \end{aligned}$$

A similar argument bounds the integral over $(-\infty, 0]$. Bounding $|g(z)| \leq Cz(1-z)$ we find that $E[\int_0^t (\int_R |g(u)(s, x)| dx)^2 ds]$ is finite. This shows that the integral $\int_0^t \int_0^\infty g(\tilde{u})(s, x) dx dW_s$ makes sense and the Ito isometry allows one to check that the L^2 convergence

$$\int_0^t \int_0^U g(\tilde{u})(s, x) dx dW_s \rightarrow \int_0^t \int_0^\infty g(\tilde{u})(s, x) dx dW_s.$$

Theorem 3 (iv) shows that $E[\int_{t_0}^t \sup_x |u_x(s, x)| ds] < \infty$ and this gives the required domination to turn $\tilde{u}_x(s, U) \rightarrow 0$ into $\int_{t_0}^t \tilde{u}_x(s, U) ds \rightarrow 0$.

This leaves the second term in (14) and the lemma will follow once we have shown that $\int_{t_0}^t \tilde{u}(s, U) \circ d\Gamma(u(s)) \rightarrow 0$ almost surely as $U \rightarrow \infty$. To see this expand the integral using the decomposition for $d\Gamma(u(t)) = dm(t, a)$ in (5) to see that

$$\begin{aligned} \int_{t_0}^t \tilde{u}(s, U) \circ d\Gamma(u(s)) &= \int_{t_0}^t \tilde{u}(s, U) \left(\frac{m_{xx}(s, a)}{m_x^2(s, a)} - f(a)m_x(s, a) \right) ds \\ &\quad - g(a) \int_{t_0}^t \tilde{u}(s, U) m_x(s, a) dW_s \\ &\quad - \frac{1}{2} g(a) \int_{t_0}^t \tilde{u}(s, U) d [\tilde{u}(\cdot, U) m_x(\cdot, a), W]_s \end{aligned} \quad (15)$$

where we have converted from a Stratonovich to an Ito integral and we are writing $[\cdot, \cdot]_t$ for the cross quadratic variation. We claim that each of these terms converge to zero almost surely. Note that the strict negativity of the derivative $u_x(t, x)$ and the relations (7) imply that the path $s \rightarrow (m_{xx}(s, a)/m_x^2(s, a)) - f(a)m_x(s, a)$ is (almost surely) continuous on $[t_0, t]$. So the first term on the right hand side of (15) converges (almost surely) to zero by dominated convergence using $\tilde{u}(s, U) \rightarrow 0$ as $U \rightarrow \infty$. The second term in (15) also converges to zero by applying the same argument to the quadratic variation $g^2(a) \int_{t_0}^t \tilde{u}^2(s, U) m_x^2(s, a) ds$. A short calculation leads to the explicit formula for the cross variation

$$\begin{aligned} d [\tilde{u}(\cdot, U) m_x(\cdot, a), W]_t &= g(\tilde{u}(t, U)) m_x(t, a) dt - g(a) m_{xx}(t, a) \tilde{u}(t, U) dt \\ &\quad - g(a) \tilde{u}(t, U) m_x^2(t, a) dt - g'(a) m_x(t, a) \tilde{u}(t, U) dt. \end{aligned}$$

Again, since also $g(\tilde{u}(t, U)) \rightarrow 0$ as $U \rightarrow \infty$, a dominated convergence argument shows that the final term in (15) converges to zero as $U \rightarrow \infty$.

This completes the proof of the first equation in the lemma. The second is similar by integrating over $[-L, 0]$ and letting $L \rightarrow \infty$. \square

Proposition 13. Suppose that f_0 is of KPP, Nagumo or unstable type. In the latter two cases suppose that $f_0(a) = 0$ and $g(a) \neq 0$. Let u be the solution to (1) started from $H(x) = I(x < 0)$ and let ν_c be the limit law constructed from u in Proposition 10. Then $\nu_c(\mathcal{D}) = 1$.

In the KPP and Nagumo cases we have the increasing limits as $t \uparrow \infty$

$$E \left[\int u(1-u)(t, x) dx \right] \uparrow \int_{\mathcal{D}} \int_R \phi(1-\phi)(x) dx \nu_c(d\phi) < \infty. \quad (16)$$

In the unstable case

$$\begin{aligned} \int_{\mathcal{D}} \int_0^\infty \phi(a-\phi)(x) dx \nu_c(d\phi) &\leq \sup_t E \left[\int_0^\infty \tilde{u}(a-\tilde{u})(t, x) dx \right] < \infty, \\ \int_{\mathcal{D}} \int_{-\infty}^0 (1-\phi)(\phi-a)(x) dx \nu_c(d\phi) &\leq \sup_t E \left[\int_{-\infty}^0 (1-\tilde{u})(\tilde{u}-a)(t, x) dx \right] < \infty. \end{aligned} \quad (17)$$

Proof We start with the case where f_0 is of KPP type. In this case there is a constant C so that $Cf_0(x) \geq x(1-x)$ on $[0, 1]$. In a similar (but easier) way to Lemma 12, one may integrate (1) over $s \in [t_0, t]$ and then $x \in R$ to find

$$\int_R (u(t, x) - u(t_0, x)) dx = \int_{t_0}^t \int_R f_0(u)(s, x) dx ds + \int_{t_0}^t \int_R g(u)(s, x) dx dW_s.$$

Taking expectations and rearranging one finds

$$\begin{aligned} t^{-1} \int_{t_0}^t \int_R E[u(1-u)(s, x)] dx ds &\leq Ct^{-1} \int_{t_0}^t \int_R E[f_0(u)(s, x)] dx ds \\ &\leq Ct^{-1} \int_R E[u(t, x) - u(t_0, x)] dx \\ &\leq Ct^{-1} \int_0^\infty E[u(t, x) + u(t_0, x)] dx \\ &\quad + Ct^{-1} \int_{-\infty}^0 E[(1-u)(t, x) + (1-u)(t_0, x)] dx. \end{aligned}$$

Using the first moments from Lemma (11) (ii) on each of the four terms of the right hand side we find that

$$\limsup t^{-1} \int_{t_0}^t \int_R E[u(1-u)(s, x)] dx ds < \infty.$$

For example

$$\begin{aligned} \int_0^\infty E[u(t, x)] dx &\leq \int_0^\infty \min\{1, e^{K_2 t} x^{-1} G_t(x)\} dx \\ &\leq \int_0^{\sqrt{2K_2 t}} 1 dx + \int_{\sqrt{2K_2 t}}^\infty e^{K_2 t} \frac{x}{2K_2 t^2} G_t(x) dx \\ &= \sqrt{2K_2 t} + \frac{1}{2K_2 t}. \end{aligned}$$

The other terms are similar.

Writing $z \rightarrow m(t, z)$ for the inverse function to $x \rightarrow u(t, x)$ we have

$$\int_{\mathcal{R}} u(1-u)(t, x) dx = - \int_0^1 z(1-z)m_z(t, z) dz.$$

The stochastic ordering of $\mathcal{L}(u(t))$ and Lemma (5) (iv) show that $t \rightarrow E[m_z(t, z)]$ is decreasing for $t > 0$. Thus $t \rightarrow \int_{\mathcal{R}} E[u(1-u)(s, x)] dx$ is increasing and we conclude that

$$\sup_{t \geq 0} \int_{\mathcal{R}} E[u(1-u)(t, x)] dx < \infty.$$

For fixed $N > 0$ the functionals $\phi \rightarrow \int_{-N}^N \phi(1-\phi)(x) dx$ are bounded and continuous on \mathcal{D}_c . So by the convergence of $\mathcal{L}(\tilde{u}(t))$ to ν_c in $\mathcal{M}(\mathcal{D}_c)$ we see that

$$\int_{-N}^N E[\tilde{u}(1-\tilde{u})(t, x)] dx \rightarrow \int_{\mathcal{D}_c} \int_{-N}^N \phi(1-\phi)(x) dx \nu_c(d\phi).$$

The last two displayed equations imply that

$$\int_{\mathcal{D}_c} \int_{\mathcal{R}} \phi(1-\phi)(x) dx \nu_c(d\phi) \leq \lim_{t \uparrow \infty} \int E[u(1-u)(t, x)] dx < \infty. \quad (18)$$

This in turn implies that ν_c only charges \mathcal{D} .

For $0 \leq N \leq M$ the function

$$I(N, t) = \int_N^M E[\tilde{u}(t, x)] dx$$

is increasing in M and also in t (since $\mathcal{L}(\tilde{u}(t))$ are stochastically increasing). We may therefore interchange the t and M limits to see that

$$\begin{aligned} \lim_{t \uparrow \infty} \int_N^\infty E[\tilde{u}(t, x)] dx &= \lim_{t \uparrow \infty} \lim_{M \uparrow \infty} \int_N^M E[\tilde{u}(t, x)] dx \\ &= \lim_{M \uparrow \infty} \lim_{t \uparrow \infty} \int_N^M E[\tilde{u}(t, x)] dx \\ &= \lim_{M \uparrow \infty} \int_{\mathcal{D}} \int_N^M \phi(x) dx \nu_c(d\phi) \\ &= \int_{\mathcal{D}} \int_N^\infty \phi(x) dx \nu_c(d\phi). \end{aligned}$$

Similarly, as $t \uparrow \infty$

$$\int_{-\infty}^{-N} E[(1-\tilde{u})(t, x)] dx \uparrow \int_{\mathcal{D}} \int_{-\infty}^{-N} (1-\phi) dx \nu_c(d\phi).$$

Since $\int_{\mathcal{D}} \int_R \phi(1 - \phi)(x) dx \nu_c(d\phi) < \infty$ we may, for any $\epsilon > 0$, choose $N_0(\epsilon)$ so that

$$\begin{aligned} \sup_t \int_{N_0(\epsilon)}^{\infty} E[\tilde{u}(t, x)] dx &\leq \int_{\mathcal{D}} \int_{N_0(\epsilon)}^{\infty} \phi dx \nu_c(d\phi) \leq \epsilon, \\ \sup_t \int_{-\infty}^{-N_0(\epsilon)} E[(1 - \tilde{u})(t, x)] dx &\leq \int_{\mathcal{D}} \int_{-\infty}^{-N_0(\epsilon)} (1 - \phi) dx \nu_c(d\phi) \leq \epsilon. \end{aligned} \quad (19)$$

This control on the tails allows us to improve on (18) to the desired result (16).

Now we consider the case where f_0 is of Nagumo type, and this is the only place we exploit the bi-stability of f_0 (that is $f'(0), f'(1) < 0$). We may fix a smooth strictly concave $h : [0, 1] \rightarrow R$ satisfying $h(0) = h(1) = 0$ and $h'(a) = 0, h'(0) > 0, h'(1) < 0$. (A running example to keep in mind is the case $f_0 = x(1 - x)(x - \frac{1}{2})$ and $h = x(1 - x)$.) Then

$$dh(u) = h'(u)u_{xx} dt + (h'f_0 + \frac{1}{2}h''g^2)(u) dt + (h'g)(u) dW_t. \quad (20)$$

The properties of h and the fact that f_0 is of Nagumo type together imply that $h'f_0 \leq 0$ on $[0, 1]$ and $h'f_0$ only vanishes at $0, a, 1$. Since $g(a) > 0$ we have $(h'f_0 + \frac{1}{2}h''g^2) < 0$ on $(0, 1)$. The derivatives at $x = 0, 1$ are non-zero and this implies that here is an $\epsilon > 0$ so that $(h'f_0 + h''g^2) \leq -\epsilon h$. The aim is to obtain a differential inequality of the form $dm \leq C - \epsilon m$ for $m(t) = \int_R E[h(u)(t, x)] dx$. Integrate (20) over $[t_0, t]$ and then $[-N, N]$ to obtain

$$\begin{aligned} &\int_{-N}^N E[h(u)(t, x)] dx - \int_{-N}^N E[h(u)(t_0, x)] dx \\ &= \int_{t_0}^t \int_{-N}^N E[h'(u)(s, x)u_{xx}(s, x)] dx ds + \int_{t_0}^t \int_{-N}^N E[(h'f_0 + \frac{1}{2}h''g^2)(u)(s, x)] dx ds \\ &= \int_{t_0}^t (E[h'(u)(s, N)u_x(s, N)] - E[h'(u)(s, -N)u_x(s, -N)]) ds \\ &\quad - \int_{t_0}^t \int_{-N}^N E[h''(u)(s, x)u_x^2(s, x)] dx ds + \int_{t_0}^t \int_{-N}^N E[(h'f_0 + \frac{1}{2}h''g^2)(u)(s, x)] dx ds \end{aligned}$$

where we have integrated by parts in the last equality. Letting $N \rightarrow \infty$ is justified (and is similar but simpler than Lemma 12) and we find

$$\begin{aligned} &\int_R E[h(u)(t, x)] dx - \int_R E[h(u)(t_0, x)] dx \\ &\leq \|h''\|_{\infty} \int_{t_0}^t \int_R E[u_x^2(s, x)] dx ds - \epsilon \int_{t_0}^t \int_R E[h(u)(s, x)] dx ds \\ &\leq \|h''\|_{\infty} \int_{t_0}^t E[\sup_z |u_x(s, z)|] ds - \epsilon \int_{t_0}^t \int_R E[h(u)(s, x)] dx ds \end{aligned}$$

where in the last step we use that

$$\int_R u_x^2(s, x) dx \leq \sup_z |u_x(s, z)| \int_R |u_x(s, x)| dx = \sup_z |u_x(s, z)|.$$

The stochastic monotonicity of $s \rightarrow \mathcal{L}(\tilde{u}(s))$ and Lemma 5 (iv) imply that the supremum $\sup_z |u_x(s, x)|$ is stochastically decreasing. Since $E[\sup_z |u_x(t_0, z)|]$ is finite by Theorem 3 (iv) we have the desired differential inequality for $m(t) = \int_R E[h(u)(t, x)] dx$. This implies that m stays bounded and since $Ch(z) \geq z(1-z)$ for some C we find

$$\sup_{t \geq t_0} \int_R E[u(1-u)(t, x)] dx \leq C \sup_{t \geq t_0} \int_R E[h(u)(s, x)] dx < \infty.$$

As in the previous KPP case this implies (16) and that v_c only charges \mathcal{D} .

Now we consider the case where f_0 is of unstable type. Rearranging the conclusion of Lemma 12 we see, after taking expectations, that

$$\begin{aligned} t^{-1} \left| \int_{t_0}^t \int_0^\infty E[f_0(\tilde{u})(s, x)] dx ds \right| &\leq t^{-1} \int_0^\infty E[\tilde{u}(t, x) + \tilde{u}(t_0, x)] dx \\ &\quad + t^{-1} \int_{t_0}^t E[|\tilde{u}_x(s, 0)|] ds \\ &\quad + at^{-1} E[|\Gamma(u(t))| + |\Gamma(u(t_0))|]. \end{aligned} \quad (21)$$

We claim that the limsup as $t \rightarrow \infty$ is finite for all three terms on the right hand side. The first term can be bounded using

$$\begin{aligned} \int_0^\infty \tilde{u}(s, x) dx &\leq (1-a)^{-1} \int_R \tilde{u}(1-\tilde{u})(s, x) dx \\ &= (1-a)^{-1} \int_R u(1-u)(s, x) dx \\ &\leq (1-a)^{-1} \int_0^\infty u(s, x) dx + (1-a)^{-1} \int_{-\infty}^0 (1-u)(s, x) dx \end{aligned}$$

and then controlled by first moments as in the KPP case. For the second term the claim follows from the fact that $s \rightarrow E[|\tilde{u}_x(s, 0)|]$ is decreasing and finite from Theorem 3 (iv). For the third term the claim follows from Lemma 11. We conclude that the limsup of the left hand side of (21) is finite. Applying a similar argument to the second equation of Lemma 12 we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} t^{-1} \left| \int_{t_0}^t \int_0^\infty E[f_0(\tilde{u})(s, x)] dx ds \right| &< \infty, \\ \limsup_{t \rightarrow \infty} t^{-1} \left| \int_{t_0}^t \int_{-\infty}^0 E[f_0(\tilde{u})(s, x)] dx ds \right| &< \infty. \end{aligned}$$

Note that f_0 is of a single sign on each of the intervals $[0, a]$ and $[a, 1]$. Indeed there exists C so that

$$C|f_0(x)| \geq x(a-x) \text{ for } x \in [0, a] \text{ and } C|f_0(x)| \geq (1-x)(x-a) \text{ for } x \in [a, 1].$$

Therefore we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} t^{-1} \int_{t_0}^t \int_0^\infty E[\tilde{u}(a - \tilde{u})(s, x)] dx ds &< \infty, \\ \limsup_{t \rightarrow \infty} t^{-1} \int_{t_0}^t \int_{-\infty}^0 E[(1 - \tilde{u})(\tilde{u} - a)(s, x)] dx ds &< \infty. \end{aligned}$$

We have

$$\begin{aligned} \int_0^\infty \tilde{u}(a - \tilde{u})(t, x) dx &= - \int_0^a z(a - z)m_z(t, z) dz \\ \int_{-\infty}^0 (1 - \tilde{u})(\tilde{u} - a)(t, x) dx &= - \int_a^1 (1 - z)(z - a)m_z(t, z) dz. \end{aligned}$$

For fixed $N > 0$ the functionals $\phi \rightarrow \int_0^N (a - \phi)\phi dx$ and $\phi \rightarrow \int_{-N}^0 (1 - \phi)(\phi - a) dx$ are bounded and continuous on \mathcal{D}_c . The same reasoning as in the previous cases yields (17). The construction of ν_c in Lemma 10 as the law of a variable U shows that

$$\phi(x) \geq a \text{ for } x < 0 \text{ and } \phi(x) \leq a \text{ for } x > 0, \text{ for } \nu_c \text{ almost all } \phi.$$

This and (17) imply that ν_c charges only \mathcal{D} or the single point $\phi \equiv a$.

The argument that there is no mass on the point $\phi \equiv a$ is a little fiddly, and we start with a brief sketch. We argue that if $\phi \equiv a$ with ν_c positive mass then there are arbitrarily wide patches in $\tilde{u}(t)$, for large t , that are flattish, that is lie close to the value a . But the height of this large flatish patch will evolve roughly like the SDE $dY = f_0(Y)dt + g(Y)dW$ with $Y_0 = a$. Since $g(a) \neq 0$ the sde will move away from the value a with non-zero probability and this would lead to an arbitrary large value of $E[\int_R |1 - \tilde{u}| |\tilde{u} - a| \tilde{u} dx]$ for all large times which contradicts (17). To implement this argument we will use the following estimate.

Lemma 14. *Let u be a solution to (1) driven by a Brownian motion W . Let Y be the solution to the SDE $dY = f_0(Y)dt + g(Y)dW$ with $Y_0 = a$. Then there exists a constant $c_0(T)$ so that for all $\eta \in (0, 1)$*

$$\int_R e^{-\eta|x|} E[|u(t, x) - Y_t|^2] dx \leq c_0(T) \int_R e^{-\eta|x|} E[|u(0, x) - a|^2] dx \quad \text{for } t \in [0, T].$$

Note that the constant c_0 does not depend on $\eta \in (0, 1)$. Considered as a constant function in x , the process Y_t is a solution to (1). This lemma therefore follows by a standard Gronwall argument in order to estimate the L^2 difference between two solutions for an equation with Lipschitz coefficients. The use of weighted norms for equations on the whole space, that is the norm $\int_R e^{-\eta|x|} E[|u(t, x) - \nu(t, x)|^2] dx$, is also standard - see, for example, the analogous estimate in the proof of Shiga [11] Theorem 2.2 for the (harder) case of an equation driven by space-time white noise.

Suppose (aiming for a contradiction) that $\nu_c(\phi \equiv a) = \delta_1 > 0$. By the convergence $\mathcal{L}(\tilde{u}(t)) \rightarrow \nu_c$ we have, for any $\eta > 0$,

$$\begin{aligned} P \left[\int_R |\tilde{u}(t, x) - a|^2 e^{-\eta|x|} dx \leq 1 \right] &\geq E \left[\left(1 - \int_R |\tilde{u}(t, x) - a|^2 e^{-\eta|x|} dx \right)_+ \right] \\ &\rightarrow \int_{\mathcal{D}_c} \left(1 - \int_R |\phi(x) - a|^2 e^{-\eta|x|} dx \right)_+ \nu_c(d\phi) \\ &\geq \delta_1. \end{aligned}$$

Suppose the solution u is defined on a filtered space $(\Omega, \mathcal{F}, (\mathcal{F}_t), P)$ and with respect to an (\mathcal{F}_t) Brownian motion W . Then for $t \geq T(\eta)$, we may choose sets $\Omega_t \in \mathcal{F}_t$ satisfying $P[\Omega_t] = \delta_1/2$ and

$$\int_{\mathbb{R}} |u(t, x) - a|^2 e^{-\eta|x|} dx \leq 1 \quad \text{on } \Omega_t.$$

Fix $t \geq T(\eta)$. Let $(Y_s : s \in [t, t+1])$ solve the SDE $dY = f_0(Y)dt + g(Y)dW$ with initial condition $Y_t = a$. Since $g(a) \neq 0$ there must exist $t^* \in (0, 1)$, $\delta_2, \delta_3 > 0$ satisfying $4\delta_2 < a \wedge (1-a)$ such that $P[|Y_{t^*} - a| \in (\delta_2, 2\delta_2)] > \delta_3$. The requirement that $4\delta_2 < a \wedge (1-a)$ ensures that when $|Y_{t^*} - a| \in (\delta_2, 2\delta_2)$ then $Y_{t^*} \geq \frac{a}{2}$ and $1 - Y_{t^*} \geq \frac{1-a}{2}$. Then

$$\begin{aligned} & \int_{\mathbb{R}} E[u|1-u||u-a|(t+t^*, x)] dx \\ & \geq \int_{\mathbb{R}} e^{-\eta|x|} E[u|1-u||u-a|(t+t^*, x)I(\Omega_t)] dx \\ & \geq \int_{\mathbb{R}} e^{-\eta|x|} E[Y_{t+t^*}|1-Y_{t+t^*}||Y_{t+t^*}-a|I(\Omega_t)] dx \\ & \quad - L_0 \int_{\mathbb{R}} e^{-\eta|x|} E[|u(t+t^*, x) - Y_{t+t^*}|I(\Omega_t)] dx \\ & =: I - II \end{aligned} \tag{22}$$

where L_0 is the Lipschitz constant of $z|1-z||a-z|$ on $[0, 1]$. We now estimate the terms I and II . Firstly,

$$\begin{aligned} I &= \frac{2}{\eta} E[Y_{t+t^*}|1-Y_{t+t^*}||Y_{t+t^*}-a|I(\Omega_t)] \\ &= \frac{2}{\eta} P[\Omega_t] E[Y_{t^*}|1-Y_{t^*}||Y_{t^*}-a| \mid Y_0 = a] \quad (\text{by the Markov property}) \\ &\geq \frac{a(1-a)\delta_1}{4\eta} E[|Y_{t^*} - a| \mid Y_0 = a] \\ &\geq \frac{a(1-a)\delta_1\delta_2\delta_3}{4\eta}. \end{aligned}$$

Secondly, using Cauchy-Schwarz,

$$\begin{aligned}
II &= L_0 E \left[I(\Omega_t) E \left[\int_R e^{-\eta|x|} |u(t+t^*, x) - Y_{t+t^*}| dx \middle| \mathcal{F}_t \right] \right] \\
&\leq L_0 \sqrt{\frac{2}{\eta}} E \left[I(\Omega_t) \left| E \left[\int_R e^{-\eta|x|} |u(t+t^*, x) - Y_{t+t^*}|^2 dx \middle| \mathcal{F}_t \right] \right|^{1/2} \right] \\
&\leq L_0 c_0(1) \sqrt{\frac{2}{\eta}} E \left[I(\Omega_t) \left| \int_R e^{-\eta|x|} |u(t, x) - a|^2 dx \right|^{1/2} \right] \quad (\text{by Lemma 14}) \\
&\leq L_0 c_0(1) \sqrt{\frac{2}{\eta}} E[I(\Omega_t)] \quad (\text{by the choice of } \Omega_t) \\
&= L_0 c_0(1) \sqrt{\frac{2}{\eta}} \frac{\delta_1}{2}.
\end{aligned}$$

Thus, substituting these estimates into (22), we find for $t \geq T(\eta)$

$$\begin{aligned}
\int_R E [\tilde{u} |1 - \tilde{u}| |\tilde{u} - a|(t+t^*, x)] dx &= \int_R E [u |1 - u| |u - a|(t+t^*, x)] dx \\
&\geq \frac{a(1-a)\delta_1\delta_2\delta_3}{4\eta} - L c_0(1) \sqrt{\frac{2}{\eta}} \frac{\delta_1}{2}.
\end{aligned}$$

By taking η small this bound can be made arbitrarily large, which contradicts (17). \square

3.3 Proof of Theorem 1

Let ν_c be the limit law constructed from u in Proposition 10. We let ν be the restriction of ν_c to \mathcal{D} . Proposition 13 shows that in all cases ν is a probability. Moreover the fact that $\mathcal{L}(\tilde{u}(t)) \rightarrow \nu_c$ in $\mathcal{M}(\mathcal{D}_c)$ implies that $\mathcal{L}(\tilde{u}(t)) \rightarrow \nu$ in $\mathcal{M}(\mathcal{D})$.

We first check that the centered solutions are still a Markov process. This can be done, as follows, by using the Dynkin criterion (see [10] Theorem 13.5) which gives a simple transition kernel condition for when a function of a Markov process remains Markov. Let $D_0 = \{\phi \in \mathcal{D} : \Gamma(\phi) = 0\}$ with the induced topology. Define the centering map $J : \mathcal{D} \rightarrow \mathcal{D}_0$ by $J(\phi) = \check{\phi}$. Let $(P_t(\phi, d\psi) : t \geq 0)$ be the Markov transition kernels for solutions to (1). Then the Dynkin criterion is that for all measurable $A \subseteq \mathcal{D}_0$ and all $\psi \in \mathcal{D}_0$ the values $P_t(\phi, J^{-1}A)$ are equal for all $\phi \in J^{-1}(\psi)$. By Lemma 5 (ii), elements of $J^{-1}(\psi)$ are translates of each other and the Dynkin criterion follows from translation invariance of solutions. As a consequence, there are transition kernels $\check{P}_t(\phi, d\psi)$ for the centered process on D_0 . We write (P_t) (respectively (\check{P}_t)) for the associated semigroups generated by these kernels and acting on measurable $F : \mathcal{D} \rightarrow R$ (respectively $F : \mathcal{D}_0 \rightarrow R$), and we write (P_t^*) and (\check{P}_t^*) for the dual semigroups acting on $\mathcal{M}(\mathcal{D})$ (respectively $\mathcal{M}(\mathcal{D}_0)$).

We aim to show that ν is the law of a stationary travelling wave by applying Markov semigroup arguments applied to the centered solutions $(\tilde{u}(t) : t \geq 0)$. Some difficulties arise since the wavefront marker Γ is only semi-continuous on \mathcal{D} , and hence \mathcal{D}_0 is a measurable but not a closed subset of \mathcal{D} .

For example, we do not yet know that $\Gamma(\phi) = 0$ for ν almost all ϕ (though we will see that this is true).

The centered law $\tilde{\nu}$ charges only D_0 and we will therefore consider it (with a slight abuse of notation) as an element of $\mathcal{M}(\mathcal{D}_0)$, where it is the image of ν under the centering map J . Take $F : \mathcal{D} \rightarrow \mathbb{R}$ that is bounded, continuous and translation invariant (that is $F(\phi) = F(\tau^a \phi)$ for all a). Then the Feller property and translation invariance of solutions imply that $P_t F$ remains bounded, continuous and translation invariant. Let F_0 be the restriction of F to D_0 . The translation invariance of F implies that $\tilde{P}_t F_0(\phi) = P_t F(\phi)$. Write Q_t^H and \tilde{Q}_t^H for the law of $u(t)$ and $\tilde{u}(t)$ on D , when u is started at H . Then

$$\begin{aligned}
\int_{\mathcal{D}_0} F_0 d(\tilde{P}_s^* \tilde{\nu}) &= \int_{\mathcal{D}_0} \tilde{P}_s F_0 d\tilde{\nu} \\
&= \int_{\mathcal{D}} P_s F d\nu \quad (\text{by translation invariance of } P_s F) \\
&= \lim_{t \rightarrow \infty} \int_{\mathcal{D}} P_s F d\tilde{Q}_t^H \quad (\text{by the convergence } \tilde{Q}_t^H \rightarrow \nu) \\
&= \lim_{t \rightarrow \infty} \int_{\mathcal{D}} P_s F dQ_t^H \quad (\text{by translation invariance of } P_s F) \\
&= \lim_{t \rightarrow \infty} \int_{\mathcal{D}} F dQ_{t+s}^H \quad (\text{by the Markov property of } u) \\
&= \lim_{t \rightarrow \infty} \int_{\mathcal{D}} F d\tilde{Q}_{t+s}^H \quad (\text{by translation invariance of } F) \\
&= \int_{\mathcal{D}} F d\nu \\
&= \int_{\mathcal{D}_0} F_0 d\tilde{\nu}.
\end{aligned}$$

This equality may now be extended, by a monotone class argument, to hold for all bounded functions that are measurable with respect to the sigma field generated by continuous translation invariant F . Lemma 15 below shows that this includes all bounded measurable translation invariant $F : \mathcal{D} \rightarrow \mathbb{R}$. Then taking $F(\phi) = I(\tilde{\phi} \in A)$ for a measurable $A \subseteq \mathcal{D}$ we find

$$\begin{aligned}
\tilde{P}_s^* \tilde{\nu}(A \cap \mathcal{D}_0) &= \tilde{P}_s^* \tilde{\nu}(\phi \in \mathcal{D}_0 : \tilde{\phi} \in A) \\
&= \int_{\mathcal{D}_0} I(\tilde{\phi} \in A) d(\tilde{P}_s^* \tilde{\nu}) \\
&= \int_{\mathcal{D}_0} I(\tilde{\phi} \in A) d\tilde{\nu} \\
&= \tilde{\nu}(A \cap D_0).
\end{aligned}$$

This yields $\tilde{P}_s^* \tilde{\nu} = \tilde{\nu}$ showing that $\tilde{\nu}$ is the law of a stationary travelling wave.

Finally we check that ν was already centered. By the regularity of solutions at any time $t > 0$ we know that $\phi \in \mathcal{C}^1$ and $\phi_x < 0$ for $\tilde{P}_t^* \tilde{\nu}$ almost all ϕ , and hence for $\tilde{\nu}$ almost all ϕ or indeed for ν

almost all ϕ . But the construction of ν showed that $\phi(x) \leq a$ for $x > 0$ and $\phi(x) \geq a$ for $x < 0$ for ν almost all ϕ . Combining these shows that $\Gamma(\phi) = 0$ for ν almost all ϕ and thus ν charges only \mathcal{D}_0 . Thus $\tilde{\nu} = \nu$ and this completes the proof.

Remark. We have fixed the centering height a throughout and suppressed its dependence in the notation. However, we wish to show that the choice of height is unimportant and, in this remark only, we shall now indicate this dependence. The construction in Proposition 10 of the stretched limit ν_c held for any centering height. We write $\nu_c^{\bar{a}}$ for this law when centered at height \bar{a} and also $\Gamma^{\bar{a}}$ for the wavefront marker at height \bar{a} and $\tilde{u}^{\bar{a}}$ for the the solution started at H centered using $\Gamma^{\bar{a}}$. The moments in Proposition 13 rely on the specific properties of f_0 and g and the distinguished point a in the definition of the three types of reaction f_0 . But these moments imply, in any of the three cases, that the law $\nu_c^{\bar{a}}$ charges only \mathcal{D} and that the restriction $\nu_c^{\bar{a}}$ to \mathcal{D} is the law of a stationary travelling wave for any centering height \bar{a} . We claim, for $\bar{a}_1, \bar{a}_2 \in (0, 1)$, that the image of $\nu^{\bar{a}_1}$ under the map $\phi \rightarrow \phi(\cdot + \Gamma^{\bar{a}_2}(\phi))$ is $\nu^{\bar{a}_2}$. Indeed $\nu^{\bar{a}_1} \stackrel{s}{\succ} \delta_H$ and so by Corollary 9 (i) and the stationarity of $\nu^{\bar{a}_1}$ we have $\nu^{\bar{a}_1} \stackrel{s}{\succ} \mathcal{L}(\tilde{u}^{\bar{a}_2}(t))$. Now letting $t \rightarrow \infty$ we have $\nu^{\bar{a}_1} \stackrel{s}{\succ} \nu^{\bar{a}_2}$. But reversing the roles of \bar{a}_1 and \bar{a}_2 we find $\nu^{\bar{a}_2} \stackrel{s}{\succ} \nu^{\bar{a}_1}$ and Lemma 6 (ii) implies that the centered copies (at any height) of $\nu^{\bar{a}_1}$ and $\nu^{\bar{a}_2}$ must coincide.

Lemma 15. *Translation invariant measurable $F : \mathcal{D} \rightarrow R$ are measurable with respect to the sigma field generated by the continuous translation invariant functions on \mathcal{D} .*

Proof. We make use of a smoother wave marker than the wave marker Γ^a for the height a . Define $\hat{\Gamma} = \int_0^1 h(a)\Gamma^a da$, where $h : (0, 1) \rightarrow R$ is continuous and compactly supported in $(0, 1)$. Then $\Gamma(\phi)$ is finite and $\Gamma(\tau^a \phi) = \Gamma(\phi) + a$ if we assume in addition that $\int h dx = 1$. Then the map $\phi \rightarrow \hat{\Gamma}(\phi)$ is continuous (since $\Gamma^a(\phi)$ is discontinuous at ϕ only for the countably many a where $\{x : \phi(x) = a\}$ has non-empty interior). We let $\hat{D}_0 = \{\phi \in \mathcal{D} : \hat{\Gamma}(\phi) = 0\}$, so that \mathcal{D}_0 is a closed subset of \mathcal{D} , and give it the induced subspace topology and Borel sigma field. For this proof only we let $\tilde{\phi}$ be the wave centered at the new wave-marker $\hat{\Gamma}$. One may now check that the map

$$\mathcal{D} \ni \phi \rightarrow J(\phi) = (\hat{\Gamma}(\phi), \tilde{\phi}) \in R \times \hat{\mathcal{D}}_0$$

is a homeomorphism. Also, every continuous (respectively measurable) translation invariant $F : \mathcal{D} \rightarrow R$ is of the form $F(\phi) = \hat{F}(\tilde{\phi})$ for some continuous (respectively measurable) $\hat{F} : \hat{\mathcal{D}}_0 \rightarrow R$. Using this one finds that

$$\begin{aligned} \tau &:= \{O \subseteq \mathcal{D} : O \text{ is open and } I(O) \text{ is translation invariant}\} \\ &= \{J^{-1}(R \times O) : O \text{ open in } \hat{D}_0\} \end{aligned}$$

and

$$\begin{aligned} &\{B \subseteq \mathcal{D} : B \text{ is measurable and } I(B) \text{ is translation invariant}\} \\ &= \{J^{-1}(R \times B) : B \text{ measurable in } \hat{D}_0\}. \end{aligned}$$

This implies that translation invariant measurable $F : \mathcal{D} \rightarrow R$ are measurable with respect to the sigma field $\sigma(\tau)$. But for an open $O \subseteq \mathcal{D}$ for which $I(O)$ is translation invariant, by using a translation invariant metric on \mathcal{D} (see the start of section 2.2), the continuous functions $F_n(\phi) = \min\{nd(\phi, O^c), 1\}$ are translation invariant and converge to $I(O)$ as $n \rightarrow \infty$, completing the proof. \square

4 The domain of attraction

Throughout this section ν is the law of the stationary travelling wave constructed in Theorem 1. The stochastic monotonicity results imply that solutions starting from a certain set of initial conditions are attracted to ν , as follows.

Proposition 16. *Suppose that f_0 is of KPP, Nagumo or unstable type. In the latter two cases suppose that $f_0(a) = 0$ and $g(a) \neq 0$. Let u be a solution to (1) whose initial condition $u(0)$ has law $\mu \in \mathcal{M}(\mathcal{D})$. If $\nu \succ^s \mu \succ^s \delta_H$ then $\mathcal{L}(\tilde{u}(t)) \rightarrow \nu$ as $t \rightarrow \infty$.*

Proof Since $\nu \succ^s \mu \succ^s \delta_H$, Corollary 9 (i) implies that

$$\nu \succ^s \mathcal{L}(\tilde{u}(t)) \succ^s \tilde{Q}_t^H \quad (23)$$

for all t . The simple idea is to let $t \rightarrow \infty$ and use $\tilde{Q}_t^H \rightarrow \nu$.

We face a familiar set of technical details to check. For any sequence $t_n \rightarrow \infty$ there is a subsequence t'_n along which $\mathcal{L}(\tilde{u}(t))$ converges in $\mathcal{M}(\mathcal{D}_c)$ to a limit which we call μ_c . For $\lambda > 0$, the map $\phi \rightarrow 1 - \exp(-\lambda \int_R \phi(1 - \phi)dx)$ is increasing on \mathcal{D} (in the stretching pre-order) and lower semi-continuous on \mathcal{D}_c . Using $\nu \succ^s \mathcal{L}(\tilde{u}(t_{n'}))$ and $\mathcal{L}(\tilde{u}(t_{n'})) \rightarrow \mu_c$ we find

$$\int_{D_c} \left(1 - \exp(-\lambda \int_R \phi(1 - \phi)dx) \right) \mu_c(d\phi) \leq \int_D \left(1 - \exp(-\lambda \int_R \phi(1 - \phi)dx) \right) \nu(d\phi)$$

Proposition 13 shows that in all cases $\int \phi(1 - \phi)dx < \infty$ for ν almost all ϕ . Therefore as $\lambda \downarrow 0$ the right hand side approaches 0. This shows that μ_c charges only D or the single points $\phi \equiv 0$ or $\phi \equiv 1$. However the construction of μ_c as the limit of centered laws ensures that $\phi(x) \leq a$ for $x > 0$ and $\phi(x) \geq a$ for $x < 0$ for μ_c almost all ϕ . We conclude that μ_c only charges \mathcal{D} .

Let μ be the restriction of μ_c to D , so that $\mathcal{L}(\tilde{u}(t_{n'})) \rightarrow \mu$ in $\mathcal{M}(\mathcal{D})$. Using Lemma 6 (iii) we may pass to the limit in (23) as $n' \rightarrow \infty$ to find $\nu \succ^s \mu \succ^s \mu$. Lemma 6 (ii) now gives that $\tilde{\mu} = \tilde{\nu}$. But we know that ν charges only ϕ for which $\phi \in C^1(\mathbb{R})$ and $\phi_x < 0$. Hence so does μ and $\tilde{\mu} = \mu = \nu$. This shows that $\mathcal{L}(\tilde{u}(t_{n'})) \rightarrow \nu$ and since (t_n) was arbitrary the proof is finished. \square

Checking the hypotheses of Proposition (16) does not look easy. Moreover one would expect the domain of attraction for ν to contain many other initial conditions. In the upcoming section we discuss a natural class of initial conditions that should be in the domain.

4.1 Solutions with finite initial width

Consider initial conditions which have a wavefront of finite width, that is satisfying

$$\text{there exist } l \leq r \text{ so that } I(x < l) \leq \phi(x) \leq I(x < r) \text{ for all } x \in \mathbb{R}. \quad (24)$$

We can exploit comparison theorems to deduce estimates for a solution started from such ϕ from the estimates for the solution started at the Heaviside function. The next lemma collects what we shall need.

Lemma 17. Let u be a solution to (1) started from an initial condition satisfying (24). Let u_l, u_r be the solutions to (1) driven by the same white noise and with initial conditions $u_l(0) = I(x < l)$ and $u_r(0) = I(x < r)$. Then, almost surely,

$$u_l(t, x) \leq u(t, x) \leq u_r(t, x) \quad \text{for all } t, x, \quad (25)$$

and

$$\Gamma(u_l(t)) \leq \Gamma(u(t)) \leq \Gamma(u_r(t)) = \Gamma(u_l(t)) + (r - l) \quad \text{for all } t. \quad (26)$$

Suppose that f_0 is of KPP or Nagumo type, with $f_0(a) = 0$ and $g(a) \neq 0$ in the latter case. Then for any $\epsilon > 0$ there exists $N(\epsilon)$ so that

$$\sup_{t \geq 0} \int_{N(\epsilon)}^{\infty} E[\tilde{u}(t, x)] dx \leq \epsilon \quad \text{and} \quad \sup_{t \geq 0} \int_{-\infty}^{-N(\epsilon)} E[1 - \tilde{u}(t, x)] dx \leq \epsilon. \quad (27)$$

In particular

$$\sup_{t \geq 0} \int_{\mathbb{R}} E[u(t, x)(1 - u(t, x))] dx < \infty. \quad (28)$$

Proof Theorem 3 (i) shows that coupled solutions u_l, u, u_r exists as desired. Note that

$$u_r(t, x) = u_l(t, x - (r - l)) \quad \text{for all } x, t, \text{ almost surely,}$$

(by uniqueness of solutions). This yields $\Gamma(u_r(t)) = \Gamma(u_l(t)) + (r - l)$. Furthermore

$$\begin{aligned} & \int_{\mathbb{R}} u(1 - u)(t, x) dx \\ & \leq \int_{-\infty}^{\Gamma(u(t))} (1 - u_l(t, x)) dx + \int_{\Gamma(u(t))}^{\infty} u_r(t, x) dx \\ & \leq \int_{-\infty}^{\Gamma(u_l(t))} (1 - u_l(t, x)) dx + \int_{\Gamma(u_r(t))}^{\infty} u_r(t, x) dx + (r - l) \\ & \leq \frac{1}{a} \int_{\mathbb{R}} u_l(1 - u_l)(t, x) dx + \frac{1}{1 - a} \int_{\mathbb{R}} u_r(1 - u_r)(t, x) dx + (r - l). \end{aligned}$$

So (28) follows from Proposition 13.

The uniform control on the tails was obtained for the solution started from H in (19). Define $N = N(\epsilon)$ by $N(\epsilon) = N_0(\epsilon) + (r - l)$ where $N_0(\epsilon)$ is chosen as in (19). Then

$$\begin{aligned} E \left[\int_{N(\epsilon)}^{\infty} \tilde{u}(t, x) dx \right] &= E \left[\int_{\Gamma(u(t)) + N(\epsilon)}^{\infty} u(t, x) dx \right] \\ &\leq E \left[\int_{\Gamma(u_l(t)) + N(\epsilon)}^{\infty} u_r(t, x) dx \right] \quad \text{using (25) and (26),} \\ &= E \left[\int_{\Gamma(u_r(t)) + N_0(\epsilon)}^{\infty} u_r(t, x) dx \right] \leq \epsilon. \end{aligned}$$

A similar estimate holds for the left hand tail which completes the proof. \square

A key step to proving Theorem 2 is an implicit formula for the expected wave-speed.

Proposition 18. *Suppose that f_0 is of KPP or Nagumo type, and that in the latter case suppose that $f_0(a) = 0$ and $g(a) \neq 0$. Suppose u is a solution to (1) with a trapped initial condition $u_0 = \phi$ as in (24). Then*

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{1}{t} E[\Gamma(u(t))] &= \int_{\mathcal{D}} \int_R f_0(\phi)(x) dx \nu(d\phi) \\ &= \frac{1}{a} \int_{\mathcal{D}} \left(\int_0^\infty f_0(\phi)(x) dx - \phi_x(0) \right) \nu(d\phi), \\ &= \frac{1}{1-a} \int_{\mathcal{D}} \left(\int_{-\infty}^0 f_0(\phi)(x) dx + \phi_x(0) \right) \nu(d\phi), \end{aligned} \quad (29)$$

where ν is the law of the stationary travelling wave constructed in Theorem 1.

Remark. The proof breaks down in the unstable case and indeed we expect these formulae to be incorrect for the unstable case (for example the last integral in (29) would always be positive in the unstable case). Indeed an examination of the proof suggests that we must have $\int \int \phi(1 - \phi) dx \nu(dx) = \infty$ in the unstable case, else we could establish (29).

Proof By Lemma 17 it is enough to establish the formulae for the solution u started at $u(0) = H$. Combining (12) and (13) we have

$$\begin{aligned} &\Gamma(u(t)) - \Gamma(u(t_0)) \\ &= \int_{t_0}^t \int_R f_0(u)(s, x) dx ds + \int_{t_0}^t \int_R g(u)(s, x) dx dW_s - \int_R (\tilde{u}(t, x) - \tilde{u}(t_0, x)) dx. \end{aligned} \quad (30)$$

The aim is to take expectations, divide by t and then take the limit $t \rightarrow \infty$. We may bound the final term by

$$\left| \int_R (\tilde{u}(t, x) - \tilde{u}(t_0, x)) dx \right| \leq C(a) \int_R u(1-u)(t, x) + u(1-u)(t_0, x) dx$$

and the moments (28) show that this term will vanish in the limit. For any N we have

$$E \left[\int_{-N}^N f_0(\tilde{u})(t, x) dx \right] \rightarrow \int_{\mathcal{D}} \int_{-N}^N f_0(\phi) dx \nu(d\phi) \quad \text{as } t \rightarrow \infty$$

and the tail estimates (27) allows one to obtain the same limit when $N = \infty$. Using this in (30) we may as planned deduce the first of the formulae in the Lemma. For the second and third formula we argue similarly with each of (12) and (13) separately. We essentially need only one new fact, that

$$t^{-1} \int_{t_0}^t E[\tilde{u}_x(s, 0)] ds \rightarrow \int_{\mathcal{D}} \phi_x(0) \nu(d\phi), \quad (31)$$

which requires us to show that $\mathcal{L}(\tilde{u}(t))$ converges in a stronger topology. Choose $t_n \uparrow \infty$. The upcoming Lemma 19 implies the tightness of $(\tilde{u}(t, x) : |x| \leq L)_{t \geq t_0}$ on $\mathcal{C}^1([-L, L], R)$. So we may find a subsequence $(t_{n'})$ along which $(\tilde{u}(t_{n'}, x) : |x| \leq L)$ converge in distribution on $\mathcal{C}^1([-L, L], R)$. The limit law must agree with that of $(\phi(x) : |x| \leq L)$ under ν . Moreover the moments from Lemma 19 show that the variables $\tilde{u}_x(t_{n'}, 0)$ are uniformly integrable. Therefore $E[\tilde{u}_x(t_{n'}, 0)] \rightarrow \int_{\mathcal{D}} \phi_x(0) \nu(d\phi)$. Since this is true for any choice of subsequence (t_n) we may deduce (31) and complete the proof. \square

Lemma 19. *Suppose that f_0 is of KPP or Nagumo type, and that in the latter case suppose that $f_0(a) = 0$ and $g(a) \neq 0$. Suppose u is a solution to (1) with a trapped initial condition $u_0 = \phi$ as in (24). Then for any $t_0, L, p > 0$*

$$E \left[\sup_{|x| \leq L} |\tilde{u}_x(t, x)|^p + \sup_{|x| \leq L} |\tilde{u}_x(t, x)|^p \right] \leq C(L, p, t_0) < \infty \quad \text{for all } t \geq t_0.$$

Proof We need to check that centering the solutions does not spoil the control of these derivatives from Theorem 3(iii). First note that

$$\int_{\mathbb{R}} (\phi - \tilde{\phi})(x) dx = \int_{\mathbb{R}} \int_0^1 (I(y \leq \phi(x)) - I(y \leq \tilde{\phi}(x))) dy dx = \Gamma(\phi)$$

by interchanging the order of integration. Hence

$$\begin{aligned} & |\Gamma(u(t)) - \Gamma(u(t - t_0))| \\ &= \left| \int (u - \tilde{u})(t, x) dx - \int (u - \tilde{u})(t - t_0, x) dx \right| \\ &\leq \int |\tilde{u}(t, x) - \tilde{u}(t - t_0, x)| dx + \int |u(t, x) - u(t - t_0, x)| dx \\ &\leq C(a) \int u(1 - u)(t, x) + u(1 - u)(t - t_0, x) dx + \int |u(t, x) - u(t - t_0, x)| dx \end{aligned} \quad (32)$$

where $C(a) = a^{-1} + (1 - a)^{-1}$. The first two terms in (32) have first moments bounded uniformly in t by (28). By conditioning on time $t - t_0$ and using Lemma 11 (i) the third term also has a bounded first moment. This shows that $E [|\Gamma(u(t)) - \Gamma(u(t - t_0))|]$ is bounded independently of $t \geq t_0$. Then we use Chebychev's inequality to estimate

$$\begin{aligned} & P \left[\sup_{|x| \leq L} |\tilde{u}_x(t, x)| > K \right] \\ &\leq P \left[\sup_{|x| \leq L + K^p} |u_x(t, x + \Gamma(u(t - t_0)))| > K \right] + P [|\Gamma(u(t)) - \Gamma(u(t - t_0))| \geq K^p] \\ &\leq K^{-2p} E \left[\sup_{|x| \leq L + K^p} |u_x(t, x + \Gamma(u(t - t_0)))|^{2p} \right] + K^{-p} E [|\Gamma(u(t)) - \Gamma(u(t - t_0))|] \\ &\leq C(t_0, p)(1 + L + K^p)K^{-2p} + CK^{-p}. \end{aligned}$$

In the final inequality we have used the moments from Theorem 3 (iii). The desired moments for $\sup_{|x| \leq L} |\tilde{u}_x(t, x)|$ follow from these tail estimates. The second derivatives are entirely similar. \square

Remark Such estimates could be used to improve the topology of convergence in Theorem 1 - indeed they imply the convergence of $\tilde{u}(t)$ holds in $\mathcal{C}_{loc}^1(\mathbb{R})$. Convergence of higher derivatives should follow in a similar way (requiring more smoothness on f, g as necessary).

4.2 Proof of Theorem 2

Consider first the case where f_0 is of KPP type. Let u be the solution to (1) with initial condition $u(0) = \phi \in \mathcal{D}$ which satisfies (24). Write \tilde{Q}_t^ϕ for the law of $\tilde{u}(t)$. For $t > 0$ write $\tilde{Q}_{[0, t]}^\phi$ for the law

$t^{-1} \int_0^t \tilde{Q}_r^\phi dr$. Choose $t_n \uparrow \infty$. We may find a subsequence $t_{n'}$ along which the laws $\tilde{Q}_{[0, t_{n'}]}^\phi$ converge as elements of $\mathcal{M}(\mathcal{D})$ to a limit which we denote as μ (use compactness of $\mathcal{M}(\mathcal{D}_c)$ and the bound (28) to show that limit points charge only \mathcal{D}). We shall show that $\mu = \nu$. The subsequence principle then implies that $\tilde{Q}_{[0, t]}^\phi \rightarrow \nu$ as $t \rightarrow \infty$ and finishes the proof of the theorem.

We will need later to know that μ charges only \mathcal{C}^1 strictly decreasing paths. To see this we will check that μ is the law of stationary travelling wave, and many arguments are as in the proof of Theorem 1. Let $\tilde{\mu}$ be the centered measure. As before, take $F : \mathcal{D} \rightarrow R$ that is bounded, continuous and translation invariant and let F_0 be the restriction of F to D_0 . Then

$$\begin{aligned}
\int_{\mathcal{D}_0} F_0 d(\tilde{P}_s^* \tilde{\mu}) &= \int_{\mathcal{D}_0} \tilde{P}_s F_0 d\tilde{\mu} \\
&= \int_{\mathcal{D}} P_s F d\mu \quad (\text{by translation invariance of } P_s F) \\
&= \lim_{n' \rightarrow \infty} \int_{\mathcal{D}} P_s F d\tilde{Q}_{[0, t_{n'}]}^\phi \\
&= \lim_{n' \rightarrow \infty} \frac{1}{t_{n'}} \int_0^{t_{n'}} \int_{\mathcal{D}} P_s F d\tilde{Q}_r^\phi dr \\
&= \lim_{n' \rightarrow \infty} \frac{1}{t_{n'}} \int_0^{t_{n'}} \int_{\mathcal{D}} P_s F dQ_r^\phi dr \quad (\text{by translation invariance of } P_s F) \\
&= \lim_{n' \rightarrow \infty} \frac{1}{t_{n'}} \int_0^{t_{n'}} \int_{\mathcal{D}} F dQ_{r+s}^\phi dr \\
&= \lim_{n' \rightarrow \infty} \frac{1}{t_{n'}} \int_s^{t_{n'}+s} \int_{\mathcal{D}} F dQ_r^\phi dr \\
&= \lim_{n' \rightarrow \infty} \frac{1}{t_{n'}} \int_0^{t_{n'}} \int_{\mathcal{D}} F dQ_r^\phi dr \\
&= \int_{\mathcal{D}_0} F_0 d\tilde{\mu}.
\end{aligned}$$

As before, this implies that $\tilde{P}_s^* \tilde{\mu} = \tilde{\mu}$ and so $\tilde{\mu}$ is the law of a stationary travelling wave. Also as before this implies that $\tilde{\mu} = \mu$.

Next we derive an implicit formula for the expected wave-speed in terms of μ . Our solution is started from ϕ and not the Heavyside function, but the formula (30) still holds. Indeed in it's derivation (in Lemma 12) we used the moment control (28), which holds for our solutions, the decay of derivatives $\tilde{u}_x(t, x)$ as $x \rightarrow \pm\infty$ from Theorem 3 (iv), and the first moment bounds on $E[u(t, x)]$ and $E[1 - u(t, x)]$ which can again be obtained by comparing with the coupled solutions u_r and u_l . The plan, once more, is to take expectations, divide by $t_{n'}$ and then let $n' \rightarrow \infty$. The third term on the right hand side of (30) does not contribute to this limit, again by using the estimates

from (28). The convergence $\tilde{Q}_{[0,t_{n'}]}^\phi \rightarrow \mu$ implies that

$$\begin{aligned} \frac{1}{t_{n'}} E \left[\int_0^{t_{n'}} \int_R f_0(\tilde{u})(s, x) dx ds \right] &= \int_{\mathcal{D}} \int_R f_0(\psi) dx \tilde{Q}_{[0,t_{n'}]}^\phi(d\psi) \\ &\rightarrow \int_{\mathcal{D}} \int_R f_0(\psi) dx \mu(d\psi). \end{aligned}$$

Here we again use the tail estimates from (27) to allow us to approximate $\int_R f_0(\phi) dx$ by the bounded functional $\int_{-N}^N f_0(\phi) dx$. We deduce that

$$\frac{1}{t_{n'}} E[\Gamma(u(t_{n'}))] \rightarrow \int_{\mathcal{D}} \int_R f_0(\psi) dx \mu(d\psi).$$

Comparing this with the earlier formula (29) we find that

$$\int_{\mathcal{D}} \int_R f_0(\psi) dx \mu(d\psi) = \int_{\mathcal{D}} \int_R f_0(\psi) dx \nu(d\psi). \quad (33)$$

Note that $\tilde{Q}_t^\phi \succ \tilde{Q}_t^H$ for all t and hence $\tilde{Q}_{[0,t]}^\phi \succ \tilde{Q}_{[0,t]}^H$ (argue by integrating against bounded increasing $F : \mathcal{D} \rightarrow R$). By the closure property Lemma 6 (iii) we find $\mu \succ \nu$. So we may take variables U, V where $\mathcal{L}(U) = \mu$, $\mathcal{L}(V) = \nu$ and $U \succ V$ almost surely. Both U and V are \mathcal{C}^1 and strictly decreasing, almost surely. Write U^{-1} and V^{-1} for their inverse functions. Lemma 5 (iv) implies that the derivatives satisfy

$$U_z^{-1}(z) \leq V_z^{-1}(z) \text{ for all } z \in (0, 1), \text{ almost surely.} \quad (34)$$

Then

$$\int_R f_0(U)(x) dx = \int_0^1 f_0(z) U_z^{-1}(z) dz \leq \int_0^1 f_0(z) V_z^{-1}(z) dz = \int_R f_0(V)(x) dx.$$

But (33) says that all the above variables have the same expectation. This implies that

$$\int_0^1 f_0(z) U_z^{-1}(z) dz = \int_0^1 f_0(z) V_z^{-1}(z) dz \quad \text{almost surely.}$$

In the KPP case we have $f_0(z) > 0$ for $z \in (0, 1)$ and we may conclude, using (34), that $U_z^{-1} = V_z^{-1}$ and hence $\tilde{U} = \tilde{V}$, almost surely. This yields that $\mu = \nu$.

Now we consider the case where f_0 is of Nagumo type. We argue in a similar manner except that we must use the alternative wavespeed formulae from Proposition 18. For $t > t_0 > 0$ write $\tilde{Q}_{[t_0,t]}^\phi$ for the law $(t - t_0)^{-1} \int_0^t \tilde{Q}_r^\phi dr$. Choose $t_n \uparrow \infty$. The laws $\tilde{Q}_{[t_0,t]}^\phi$ are again tight in $\mathcal{M}(\mathcal{D}_c)$ and the limit points charge only \mathcal{D} (by (28)). The derivative estimates in Lemma 19 imply that the laws of $(\phi_x(x) : |x| \leq L)$ under $\tilde{Q}_{[t_0,t]}^\phi$ are also tight on $\mathcal{C}^1[-L, L]$. So we may choose a subsequence $(t_{n'})$ along which the laws $\tilde{Q}_{[t_0,t_{n'}]}^\phi$ converge as elements of $\mathcal{M}(\mathcal{D})$ to a limit μ , the law of a stochastic

travelling wave, and where $(\phi_x(x) : |x| \leq L)$ also converge in distribution on $\mathcal{C}^1[-L, L]$. We claim that

$$\begin{aligned} \frac{1}{t_{n'}} E[\Gamma(u(t_{n'}))] &\rightarrow \frac{1}{a} \int_{\mathcal{D}} \left(\int_0^\infty f_0(\phi(x)) dx - \phi_x(0) \right) \mu(d\phi) \\ &= \frac{1}{1-a} \int_{\mathcal{D}} \left(\int_{-\infty}^0 f_0(\phi(x)) dx + \phi_x(0) \right) \mu(d\phi). \end{aligned}$$

The extra ingredient to derive these formulae is that

$$t_{n'}^{-1} \int_{t_0}^{t_{n'}} E[\tilde{u}_x(s, 0)] ds = \int_{\mathcal{D}} \phi_x(0) \tilde{Q}_{[t_0, t_{n'}]}^\phi(d\phi) \rightarrow \int_{\mathcal{D}} \phi_x(0) \mu(d\phi).$$

This follows from the convergence in the space $\mathcal{C}^1[-L, L]$ and the uniform integrability of $\phi_x(0)$ under the laws $\tilde{Q}_{[t_0, t_{n'}]}^\phi$ implied by Lemma 19. Now we compare these two formulae for the expected wave speed with those from (29) to find that

$$\begin{aligned} \int_{\mathcal{D}} \left(\int_0^\infty f_0(\phi)(x) dx - \phi_x(0) \right) \mu(d\phi) &= \int_{\mathcal{D}} \left(\int_0^\infty f_0(\phi)(x) dx - \phi_x(0) \right) \nu(d\phi), \\ \int_{\mathcal{D}} \left(\int_{-\infty}^0 f_0(\phi)(x) dx + \phi_x(0) \right) \mu(d\phi) &= \int_{\mathcal{D}} \left(\int_{-\infty}^0 f_0(\phi)(x) dx + \phi_x(0) \right) \nu(d\phi). \end{aligned} \quad (35)$$

We again exploit the fact that $\mu \stackrel{s}{\succ} \nu$. As before we take variables U, V where $\mathcal{L}(U) = \mu$, $\mathcal{L}(V) = \nu$ and $U \succ V$ almost surely. Using (34) we find that

$$\begin{aligned} \int_0^\infty f_0(U)(x) dx - U_x(0) &= - \int_0^a f_0(z) U_z^{-1}(z) dz - U_x(0) \\ &\leq - \int_0^a f_0(z) V_z^{-1}(z) dz - V_x(0) \\ &= \int_0^\infty f_0(V)(x) dx - V_x(0). \end{aligned}$$

The first formula in (35) shows that the expectations of both sides are equal. Since $f_0 > 0$ on $(0, a)$, we may conclude that $U_z^{-1}(z) = V_z^{-1}(z)$ for $z \in (0, a)$. Applying the same reasoning to the second formula in (35) we find that $U_z^{-1}(z) = V_z^{-1}(z)$ for $z \in (a, 1)$. These imply that $\tilde{U} = \tilde{V}$ as before and this concludes the proof in the Nagumo case and completes this paper.

References

- [1] Bramson, Maury. Convergence of solutions of the Kolmogorov equation to travelling waves. Mem. Amer. Math. Soc. 44 (1983), no. 285, iv+190 pp. MR0705746
- [2] Fife, Paul C.; McLeod, J. B. A phase plane discussion of convergence to travelling fronts for nonlinear diffusion. Arch. Rational Mech. Anal. 75 (1980/81), no. 4, 281–314. MR0607901

- [3] Kunita, Hiroshi. Stochastic flows and stochastic differential equations. Cambridge Studies in Advanced Mathematics, 24. Cambridge University Press, Cambridge, 1990. xiv+346 pp. ISBN: 0-521-35050-6 MR1070361
- [4] Lindvall, Torgny. On Strassen's theorem on stochastic domination. Electron. Comm. Probab. 4 (1999), 51–59 (electronic). MR1711599
- [5] McKean, H. P. Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov. Comm. Pure Appl. Math. 28 (1975), no. 3, 323–331. MR0400428
- [6] Mueller, Carl. On the support of solutions to the heat equation with noise. Stochastics Stochastics Rep. 37 (1991), no. 4, 225–245. MR1149348
- [7] Murray, J. D. Mathematical biology. Second edition. Biomathematics, 19. Springer-Verlag, Berlin, 1993. xiv+767 pp. ISBN: 3-540-57204-X MR1239892
- [8] Nakayama, Toshiyuki. Support theorem for mild solutions of SDE's in Hilbert spaces. J. Math. Sci. Univ. Tokyo 11 (2004), no. 3, 245–311. MR2097527
- [9] Revuz, Daniel; Yor, Marc. Continuous martingales and Brownian motion. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 293. Springer-Verlag, Berlin, 1991. x+533 pp. ISBN: 3-540-52167-4 MR1083357
- [10] Sharpe, Michael. General theory of Markov processes. Pure and Applied Mathematics, 133. Academic Press, Inc., Boston, MA, 1988. xii+419 pp. ISBN: 0-12-639060-6 MR0958914
- [11] Shiga, Tokuzo. Two contrasting properties of solutions for one-dimensional stochastic partial differential equations. Canad. J. Math. 46 (1994), no. 2, 415–437. MR1271224
- [12] Tessitore, Gianmario; Zabczyk, Jerzy. Wong-Zakai approximations of stochastic evolution equations. J. Evol. Equ. 6 (2006), no. 4, 621–655. MR2267702
- [13] N. Woodward, University of Warwick Ph.D. Thesis, Stochastic Travelling waves, 2010.