

Research Article

On Boundaries of Parallelizable Regions of Flows of Free Mappings

Zbigniew Leśniak

Received 7 May 2007; Revised 19 August 2007; Accepted 5 September 2007

Recommended by John Michael Rassias

We are interested in the first prolongational limit set of the boundary of parallelizable regions of a given flow of the plane which has no fixed points. We prove that for every point from the boundary of a maximal parallelizable region, there exists exactly one orbit contained in this region which is a subset of the first prolongational limit set of the point. Using these uniquely determined orbits, we study the structure of maximal parallelizable regions.

Copyright © 2007 Zbigniew Leśniak. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

We assume that f is a *free mapping*, that is, an orientation preserving homeomorphism of the plane onto itself without fixed points. We consider a relation in \mathbb{R}^2 defined in the following way:

$p \sim q$ if $p = q$ or p and q are endpoints of some arc K for which $f^n(K) \rightarrow \infty$ as $n \rightarrow \pm\infty$. By an *arc* K with endpoints p and q , we mean that the image of a homeomorphism $c : [0, 1] \rightarrow c([0, 1])$ satisfying conditions $c(0) = p$, $c(1) = q$, where the topology on $c([0, 1])$ is induced by the topology of \mathbb{R}^2 . It turns out that the relation defined above is an equivalence relation (see [1]) and has the same equivalence classes as the relation defined by Andrea [2]. Moreover, each equivalence class is an invariant simply connected set (see [2, 1]).

From now on, we assume that f is embeddeable in a flow $\{f^t : t \in \mathbb{R}\}$. It follows from the Jordan theorem that each orbit C of $\{f^t : t \in \mathbb{R}\}$ divides the plane into two simply connected regions. Note that each of them is invariant under f^t for $t \in \mathbb{R}$. Thus two different orbits C_p and C_q of points p and q , respectively, divide the plane into three simply

2 Abstract and Applied Analysis

connected invariant regions, one of which contains both C_p and C_q in its boundary. We will call this region the *strip* between C_p and C_q and denote it by D_{pq} .

For any distinct orbits $C_{p_1}, C_{p_2}, C_{p_3}$ of $\{f^t : t \in \mathbb{R}\}$, one of the following two possibilities must be satisfied: exactly one of the orbits $C_{p_1}, C_{p_2}, C_{p_3}$ is contained in the strip between the other two, or each of the orbits $C_{p_1}, C_{p_2}, C_{p_3}$ is contained in the strip between the other two. In the first case, if C_{p_j} is the orbit which lies in the strip between C_{p_i} and C_{p_k} , we will write $C_{p_i}|C_{p_j}|C_{p_k}$ ($i, j, k \in \{1, 2, 3\}$ and i, j, k are different). In the second case, we will write $|C_{p_i}, C_{p_j}, C_{p_k}|$ (see [3, 4]).

Put

$$\begin{aligned} J^+(q) &:= \{p \in \mathbb{R}^2 : \text{there exist a sequence } (q_n)_{n \in \mathbb{N}} \text{ and a sequence } (t_n)_{n \in \mathbb{N}} \\ &\quad \text{such that } q_n \rightarrow q, t_n \rightarrow +\infty, f^{t_n}(q_n) \rightarrow p \text{ as } n \rightarrow +\infty\}, \\ J^-(q) &:= \{p \in \mathbb{R}^2 : \text{there exist a sequence } (q_n)_{n \in \mathbb{N}} \text{ and a sequence } (t_n)_{n \in \mathbb{N}} \\ &\quad \text{such that } q_n \rightarrow q, t_n \rightarrow -\infty, f^{t_n}(q_n) \rightarrow p \text{ as } n \rightarrow +\infty\}. \end{aligned} \quad (1.1)$$

The set $J(q) := J^+(q) \cup J^-(q)$ is called the *first prolongational limit set* of q . Let us observe that $p \in J(q)$ if and only if $q \in J(p)$ for any $p, q \in \mathbb{R}^2$. For a subset $H \subset \mathbb{R}^2$, we define

$$J(H) := \bigcup_{q \in H} J(q). \quad (1.2)$$

One can observe that for each $p \in \mathbb{R}^2$, the set $J(p)$ is invariant. In [5], it has been proved that each orbit contained in $J(\mathbb{R}^2)$ is a boundary orbit of an equivalence class. Therefore every equivalence class can contain at most two orbits from $J(\mathbb{R}^2)$ (see [6]).

An invariant region $M \subset \mathbb{R}^2$ is said to be *parallelizable* if there exists a homeomorphism φ mapping M onto \mathbb{R}^2 such that

$$f^t(x) = \varphi^{-1}(\varphi(x) + (t, 0)) \quad \text{for } x \in M. \quad (1.3)$$

It is known that a region M is parallelizable if and only if there exists a homeomorphic image K of a straight line which is a closed set in M such that K has exactly one common point with every orbit of $\{f^t : t \in \mathbb{R}\}$ contained in M (see [7, page 49], and, e.g., [6]). We will call such a set K a *section* in M .

It is known that a region M is parallelizable if and only if $J(M) \cap M = \emptyset$ (see [7, pages 46 and 49]). Hence for every parallelizable region M , we have $J(M) \subset \text{fr} M$. If M is a maximal parallelizable region (i.e., M is not contained properly in any parallelizable region), then $J(M) = \text{fr} M$ (see [8]). In [5], it has been proved that every maximal parallelizable region M is a union of equivalence classes of the relation \sim .

Now we collect the results from [5, 9] which are needed in this paper.

PROPOSITION 1.1. (see [5]) *Let M be a parallelizable region and let $p \in \text{fr} M$. Then $\text{cl} M \setminus C_p$ is contained in one of the components of $\mathbb{R}^2 \setminus C_p$.*

PROPOSITION 1.2. (see [5]) *Let M be a maximal parallelizable region and $p \in \text{fr} M$. Let G_0 be the equivalence class which contains p . Assume that G_0 does not consist of just one orbit. Then $p \notin J(q)$ for each point q belonging to the component of $\mathbb{R}^2 \setminus C_p$ that does not contain M .*

PROPOSITION 1.3. (see [9]) Let p and q belong to different equivalence classes G_1 and G_2 , respectively. Then there exists a point r lying in the strip between the orbits C_p and C_q of p and q , respectively, such that $r \notin G_1 \cup G_2$.

PROPOSITION 1.4. (see [9]) Let M be a parallelizable region. Let $G_1 \cup G_2 \subset M$ and $\text{fr} G_1 \cap \text{fr} G_2 \neq \emptyset$. Let $p \in G_1$, $q \in G_2$. Then there exists a point $z \in D_{pq}$ such that $z \in \text{fr} M$. Moreover $|C_p, C_q, C_z|$ for each $z \in D_{pq} \cap \text{fr} M$.

2. Boundary orbits of a parallelizable region

In this section, we prove some properties of boundary orbits of parallelizable regions. The main result of this section says that for every point from the boundary of a maximal parallelizable region, there exists exactly one orbit contained in this region which is a subset of the first prolongational limit set of the point.

PROPOSITION 2.1. Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$. Then $\text{fr} M$ is invariant.

Proof. Let $p \in \text{fr} M$ and $t \in \mathbb{R}$. Then on account of Proposition 1.1, M is contained in one of the components of $\mathbb{R}^2 \setminus C_p$. Denote this component by H_0 , and the other by H_1 . Fix $\varepsilon > 0$ and consider the ball $B(f^t(p), \varepsilon)$ centered at $f^t(p)$ with radius ε . By the continuity of f^t , there exists $\delta > 0$ such that $f^t(B(p, \delta)) \subset B(f^t(p), \varepsilon)$, where $B(p, \delta)$ denotes the ball centered at p with radius δ .

Since $p \in \text{fr} M$, there exists $r \in M \cap B(p, \delta)$. Thus $f^t(r) \in B(f^t(p), \varepsilon)$. Moreover, $f^t(r) \in M$ since M is invariant. Consequently, $B(f^t(p), \varepsilon)$ contains a point from M . On the other hand, $B(f^t(p), \varepsilon) \cap H_1$ does not contain any point from M . Thus $f^t(p) \in \text{fr} M$. \square

PROPOSITION 2.2. Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$. Then for all distinct orbits $C_{p_1}, C_{p_2}, C_{p_3}$ contained in $\text{fr} M$, the relation $|C_{p_1}, C_{p_2}, C_{p_3}|$ holds.

Proof. Let $C_{p_1}, C_{p_2}, C_{p_3}$ be distinct orbits which are contained in $\text{fr} M$. Suppose, on the contrary, that for these orbits the relation $\cdot | \cdot | \cdot$ holds. Without loss of generality, we can consider only the case $C_{p_1} | C_{p_2} | C_{p_3}$. Then the orbits C_{p_1} and C_{p_3} are contained in different components of $\mathbb{R}^2 \setminus C_{p_2}$. On the other hand, by Proposition 1.1, $\text{cl} M \setminus C_{p_2}$ is contained in the same component of $\mathbb{R}^2 \setminus C_{p_2}$. Hence C_{p_1} and C_{p_3} are contained in the same component of $\mathbb{R}^2 \setminus C_{p_2}$ since $C_{p_1} \cup C_{p_3} \subset \text{cl} M \setminus C_{p_2}$. Thus we get a contradiction, and consequently $|C_{p_1}, C_{p_2}, C_{p_3}|$. \square

PROPOSITION 2.3. Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$. Let $r \in M$ and let H be a component of $\mathbb{R}^2 \setminus C_r$. Then for all distinct orbits C_{p_1}, C_{p_2} contained in $\text{fr} M \cap H$, the relation $|C_{p_1}, C_{p_2}, C_r|$ holds.

Proof. By Proposition 1.1, the points r, p_1 and r, p_2 are contained in the same component of $\mathbb{R}^2 \setminus C_{p_2}$ and in the same component of $\mathbb{R}^2 \setminus C_{p_1}$, respectively. Hence, by assumption that p_1 and p_2 are contained in the same component of $\mathbb{R}^2 \setminus C_r$, we obtain $|C_{p_1}, C_{p_2}, C_r|$. \square

PROPOSITION 2.4. Let $q_1, q_2 \in J(p)$, $C_{q_1} \neq C_{q_2}$. Then $|C_{q_1}, C_{q_2}, C_r|$ for every $r \in D_{q_1, q_2} \setminus C_p$ holds (cf. Figure 2.1).




FIGURE 2.1. The first prolongational limit set of p containing two orbits.

Proof. First we show that $p \in D_{q_1, q_2}$. Suppose, on the contrary, that p belongs to the component of $\mathbb{R}^2 \setminus C_{q_1}$ which does not contain q_2 . Denote this component by H_0 . Then $J(p) \subset \text{cl}H_0 = H_0 \cup C_{q_1}$. Hence $q_2 \notin J(p)$, which is a contradiction. In the same way, we can show that p cannot belong to the component of $\mathbb{R}^2 \setminus C_{q_2}$ which does not contain q_1 . Fix a point $r \in D_{q_1, q_2} \setminus C_p$. Then either $|C_{q_1}, C_{q_2}, C_r|$ or $C_{q_1} | C_r | C_{q_2}$ holds. We show that the second possibility cannot hold. Suppose that $C_{q_1} | C_r | C_{q_2}$ holds. Then either $p \in D_{q_1, r}$ or $p \in D_{r, q_2}$ since $p \notin C_r$ and $p \in D_{q_1, q_2}$. The first case contradicts the assumption that $q_2 \in J(p)$, and the second one contradicts the assumption that $q_1 \in J(p)$ since $J(p) \subset \text{cl}H_1$, where H_1 is the component of $\mathbb{R}^2 \setminus C_r$ which contains p . Thus $|C_{q_1}, C_{q_2}, C_r|$ holds. \square

COROLLARY 2.5. *Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$. Let $p \in \text{fr}M$ and $q_1, q_2 \in M$. Assume that $q_1, q_2 \in J(p)$. Then $C_{q_1} = C_{q_2}$.*

Proof. Suppose, on the contrary, that $C_{q_1} \neq C_{q_2}$. Since $q_1, q_2 \in M$ and M is arcwise connected, there exists a point $r \in M \cap D_{q_1, q_2}$. Hence by the parallelizability of M , we get $C_{q_1} | C_r | C_{q_2}$. By Proposition 2.1, we have $C_p \subset \text{fr}M$. Hence $r \notin C_p$ since $r \in M$ and M is open. Thus on account of Proposition 2.4, we have $|C_{q_1}, C_{q_2}, C_r|$, which is a contradiction. \square

Remark 2.6. From Corollary 2.5, we get that for every parallelizable region M and every $p \in \text{fr}M$, the set $M \cap J(p)$ is either an orbit (in case $p \in J(M)$) or empty (in case $p \in \text{fr}M \setminus J(M)$). In the case where M is a maximal parallelizable region such that $M \neq \mathbb{R}^2$ (i.e., $\text{fr}M \neq \emptyset$), the existence of such an orbit for each $p \in \text{fr}M$ follows from the fact that $J(M) = \text{fr}M$ (see [8]). In this case, for each $p \in \text{fr}M$ the set $J(p)$ can contain also orbits from $\text{fr}M$ and orbits from the component of $\mathbb{R}^2 \setminus C_p$ which does not contain M . By Proposition 1.2, the last possibility can hold only if the equivalence class containing p consists of just one orbit.

3. First prolongational limit set of the boundary of a parallelizable region

In this section, we study properties of orbits contained in a parallelizable region M by using the set $J(\text{fr}M) \cap M$.

PROPOSITION 3.1. *Let $p \in J(q)$. Then $|C_p, C_q, C_r|$ for every $r \in D_{pq}$ holds.*




FIGURE 3.1. A parallelizable region with two boundary orbits.

Proof. Since $r \in D_{pq}$, the points r and q belong to the same component of $\mathbb{R}^2 \setminus C_p$ and r and p belong to the same component of $\mathbb{R}^2 \setminus C_q$. Now we prove that the points p and q are elements of the same component of $\mathbb{R}^2 \setminus C_r$. Denote by H_0 the component of $\mathbb{R}^2 \setminus C_r$ which contains q . Then, by the definition of $J(q)$, we have $J(q) \subset \text{cl}H_0$. Hence $p \in H_0$ since $p \notin C_r$. Therefore $|C_p, C_q, C_r|$ holds since each orbit of C_p, C_q, C_r divides the plane in such a way that the other two orbits are contained in the same component. \square

COROLLARY 3.2. *Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$, $p \in \text{fr}M$ and $q \in M \cap J(p)$. Let $r \in M$ be contained in the component of $\mathbb{R}^2 \setminus C_q$ which contains p . Then $|C_p, C_q, C_r|$ holds.*

Proof. On account of Proposition 1.1, the point r is contained in the component of $\mathbb{R}^2 \setminus C_p$ which contains q . Thus $r \in D_{pq}$. Hence by Proposition 3.1, we have $|C_p, C_q, C_r|$. \square

PROPOSITION 3.3. *Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$. Let $p_1, p_2 \in \text{fr}M$, $q_1, q_2 \in M$, $q_1 \in J(p_1)$, $q_2 \in J(p_2)$, and $C_{q_1} \neq C_{q_2}$. Then there exists $r \in M$ such that $C_{q_1} | C_r | C_{p_2}$, $C_{p_1} | C_r | C_{q_2}$, and $C_{p_1} | C_r | C_{p_2}$ hold (cf. Figure 3.1).*

Proof. Since $q_1, q_2 \in M$, $C_{q_1} \neq C_{q_2}$, and M is arcwise connected, there exists $r \in M \cap D_{q_1 q_2}$. Then $C_{q_1} | C_r | C_{q_2}$ holds since M is parallelizable. Denote by H_1 the component of $\mathbb{R}^2 \setminus C_r$ which contains q_1 and by H_2 the component of $\mathbb{R}^2 \setminus C_r$ which contains q_2 . Then by the definition of the first prolongational limit set, we have $p_1 \in \text{cl}H_1$ and $p_2 \in \text{cl}H_2$, since $p_1 \in J(q_1)$, $p_2 \in J(q_2)$. By the fact that $p_1, p_2 \notin M$, we have $p_1, p_2 \notin C_r$. Thus $p_1 \in H_1$ and $p_2 \in H_2$. Since each component of $\mathbb{R}^2 \setminus C_r$ is invariant, we have $C_{q_1} \subset H_1$ and $C_{q_2} \subset H_2$. Consequently $C_{q_1} | C_r | C_{p_2}$, $C_{p_1} | C_r | C_{q_2}$, and $C_{p_1} | C_r | C_{p_2}$ hold. \square

Remark 3.4. From Proposition 3.3, we get that if C_{p_1} and C_{p_2} are boundary orbits of a maximal parallelizable region M such that the only orbit C_{q_1} contained in $M \cap J(p_1)$ and the only orbit C_{q_2} contained in $M \cap J(p_2)$ are different, then there exists an orbit $C_r \subset M$ such that C_{p_1} and C_{p_2} belong to the different components of $\mathbb{R}^2 \setminus C_r$. However, in the case where the boundary orbits C_{p_1} and C_{p_2} have the same orbit C_q contained in $M \cap J(p_1)$ and in $M \cap J(p_2)$, we get from Proposition 2.4 that for every $r \in M \setminus C_q$, the orbits C_{p_1} and C_{p_2} belong to the same component of $\mathbb{R}^2 \setminus C_r$ (the assumptions of Proposition 2.4 are satisfied, since on account of Proposition 1.1 $M \subset D_{p_1, p_2}$). Moreover, by Corollary 3.2

for $i \in \{1, 2\}$, we have $|C_{p_i}, C_q, C_r|$ if p_i and r are contained in the same component of $\mathbb{R}^2 \setminus C_q$.

4. Properties of components of parallelizable regions

In this section, we will consider orbits C_r of a parallelizable region M having the property that at least one of the components of $\mathbb{R}^2 \setminus C_r$ does not contain any point from $J(\text{fr}M) \cap M$.

PROPOSITION 4.1. *Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$. Let $r \in M$ and let H be a component of $\mathbb{R}^2 \setminus C_r$. Assume that $H \cap J(\text{fr}M) \cap M = \emptyset$. Then $H \cap M$ is contained in an equivalence class.*

Proof. Suppose, on the contrary, that there exist $p, q \in H \cap M$ such that $p \in G_1$ and $q \in G_2$ for some distinct equivalence classes G_1, G_2 . On account of Proposition 1.3, there exists a point $s \in D_{pq}$ such that $s \notin G_1 \cup G_2$. Denote by G_3 the equivalence class which contains s . Now we will show that D_{pq} is contained in an equivalence class.

First we will show that $D_{pq} \subset M$. Suppose, on the contrary, that there exists $x \in D_{pq}$ such that $x \notin M$. Put $A := D_{pq} \cap M$ and $B := D_{pq} \setminus A$. Since D_{pq} is connected, A is open in D_{pq} , $A \neq \emptyset$, and $B \neq \emptyset$, there exists a point $y \in D_{pq}$ such that $y \in \text{fr}A$. Hence $y \in \text{fr}M$.

Let M_1 be a maximal parallelizable region such that $M \subset M_1$. Now we prove that $M_1 \cap \text{cl}D_{pq} = M \cap \text{cl}D_{pq}$. Let $z \in M_1 \cap D_{pq}$. Then $C_p|C_z|C_q$ holds since M_1 is parallelizable. Hence $C_z \cap M \neq \emptyset$ since M is arcwise connected and $p, q \in M$. Thus by the fact that M is invariant, we have $z \in M$.

Take a ball $B(y, \varepsilon)$ centered at y with radius $\varepsilon > 0$. Without loss of generality, we can assume that $B(y, \varepsilon) \subset D_{pq}$ (such a ball exists since D_{pq} is an open set). From the fact that $y \in \text{fr}M$, we obtain that there exist $z_1 \in B(y, \varepsilon) \cap M$ and $z_2 \in B(y, \varepsilon) \setminus M$. Then by the equality $M_1 \cap \text{cl}D_{pq} = M \cap \text{cl}D_{pq}$, we have $z_1 \in M_1$ and $z_2 \notin M_1$. Consequently $y \in \text{fr}M_1$.

Since M_1 is a maximal parallelizable region, we have $J(M_1) = \text{fr}M_1$ (see [8]). Thus $y \in J(M_1)$. Hence $J(y) \cap M_1 \neq \emptyset$. By the definition of the first prolongational limit set, we have $J(y) \subset \text{cl}D_{pq}$ since $y \in D_{pq}$. Hence $J(y) \cap M \neq \emptyset$ since $M_1 \cap \text{cl}D_{pq} = M \cap \text{cl}D_{pq}$. Thus by the fact that $\text{cl}D_{pq} \subset H$, the set $J(y) \cap M$ is contained in H , which contradicts the assumption that $H \cap J(\text{fr}M) \cap M = \emptyset$. Consequently $D_{pq} \subset M$.

Fix $p_1, q_1 \in D_{pq}$. Then $p_1, q_1 \in M$. Since M is parallelizable, there exists a homeomorphism $\varphi : M \rightarrow \mathbb{R}^2$ such that $f^t(x) = \varphi^{-1}(\varphi(x) + (t, 0))$ for $x \in M$ and $t \in \mathbb{R}$. Let K be preimage of the segment with endpoints $\varphi(p_1)$ and $\varphi(q_1)$. Then K is an arc with endpoints p_1 and q_1 . We will prove that $f^n(K) \rightarrow \infty$ as $n \rightarrow \pm\infty$.

Take a ball $B(s, \varepsilon)$ centered at a point $s \in D_{pq}$ with radius $\varepsilon > 0$. Then $\text{cl}B(s, \varepsilon) \cap \text{cl}D_{pq}$ is a compact set. Hence $\varphi(\text{cl}B(s, \varepsilon) \cap \text{cl}D_{pq})$ is compact, since φ is a homeomorphism. Using properties of the flow of translations, we get $(\varphi(K) + (n, 0)) \cap \varphi(\text{cl}B(s, \varepsilon) \cap \text{cl}D_{pq}) \neq \emptyset$ only for finitely many $n \in \mathbb{Z}$. Hence $f^n(K) \cap (\text{cl}B(s, \varepsilon) \cap \text{cl}D_{pq}) \neq \emptyset$ only for finitely many $n \in \mathbb{Z}$. Since D_{pq} is invariant and $K \subset D_{pq}$, we have $f^n(K) \cap (\text{cl}B(s, \varepsilon) \setminus \text{cl}D_{pq}) = \emptyset$ for all $n \in \mathbb{Z}$. Hence by the definition of the equivalence relation, p_1 and q_1 belong to the same class. Thus we have shown that D_{pq} is contained in an equivalence class. Since




FIGURE 4.1. A maximal parallelizable region containing two classes.

$s \in D_{pq} \cap G_3$, we have $D_{pq} \subset G_3$. Hence by the fact that $p \notin G_3$ and $q \notin G_3$, we get $D_{pq} = G_3$ since each equivalence class is connected.

From the fact that $p \in G_1$, $q \in G_2$, $D_{pq} = G_3$, it follows that $p \in \text{fr} G_1 \cap \text{fr} G_3$ and $q \in \text{fr} G_2 \cap \text{fr} G_3$. Assume without loss of generality that q is contained in the component of $\mathbb{R}^2 \setminus C_p$ which does not contain r . Then $C_r | C_p | C_q$ holds. Let $y \in D_{pq}$. Then $C_p | C_y | C_q$ holds since $\text{cl} D_{pq} \subset M$ and M is parallelizable. Hence $D_{yq} \subset D_{pq}$. On account of Proposition 1.4, there exists a point $z \in D_{yq}$ such that $z \in \text{fr} M$, since $G_2 \cup G_3 \subset M$ and $\text{fr} G_2 \cap \text{fr} G_3 \neq \emptyset$. Hence $z \in D_{pq}$ and $z \notin M$ since $D_{yq} \subset D_{pq}$ and M is an open set, respectively. But this contradicts the fact that $D_{pq} \subset M$. Thus $H \cap M$ is contained in an equivalence class. \square

COROLLARY 4.2. *Let M be a parallelizable region of $\{f^t : t \in \mathbb{R}\}$. Let $r \in M$ and let H be a component of $\mathbb{R}^2 \setminus C_r$. Assume that $H \cap \text{fr} M = \emptyset$. Then $H \subset M$ and H is contained in an equivalence class.*

Proof. Let $H' = \mathbb{R}^2 \setminus (C_r \cup H)$. From the assumption $H \cap \text{fr} M = \emptyset$, we obtain that $\text{fr} M \subset H'$ since $C_r \subset M$. Thus by the definition of the first prolongational limit set, $J(\text{fr} M) \subset \text{cl} H' = H' \cup C_r$. Hence $H \cap J(\text{fr} M) = \emptyset$. Thus on account of Proposition 4.1, $H \cap M$ is contained in an equivalence class. Put $H_1 = H \cap M$ and $H_2 = H \setminus H_1$. Then H_1 is an open set in H . Suppose, on the contrary, that $H_2 \neq \emptyset$. Then H_2 cannot be an open set in H since H is connected, $H_1 \cap H_2 = \emptyset$, and $H = H_1 \cup H_2$. Hence there exists a point $p \in H_2 \cap \text{fr} H_2$. Take a ball $B(p, \varepsilon)$ centered at p with radius $\varepsilon > 0$ such that $B(p, \varepsilon) \subset H$. Then there exist $q \notin H_2$ such that $q \in B(p, \varepsilon)$, since $p \in \text{fr} H_2$. Hence $q \in H_1$. Thus $p \in \text{fr} H_1$ and consequently $p \in \text{fr} M$, which contradicts the assumption that $H \cap \text{fr} M = \emptyset$. Hence $H_2 = \emptyset$ and consequently $H \subset M$. Thus $H \cap M = H$ and H is contained in an equivalence class. \square

Remark 4.3. From Proposition 4.1, we do not obtain that H is contained in an equivalence class. Let us consider the case where $J(\mathbb{R}^2) = C_r \cup C_q$ for some $r, q \in \mathbb{R}^2$ such that $r \notin C_q$ (cf. Figure 4.1). Let H be the component of $\mathbb{R}^2 \setminus C_r$ which contains q . Let $H' = \mathbb{R}^2 \setminus (C_r \cup H)$ and let M be a maximal parallelizable region containing r . Then $M = H' \cup C_r \cup D_{rq}$, $\text{fr} M = C_q$, and $H \cap J(\text{fr} M) \cap M = \emptyset$. The only equivalence class containing $H \cap M$ is the strip D_{rq} , and D_{rq} is a proper subset of H since $q \notin D_{rq}$.

References

- [1] Z. Leśniak, "On an equivalence relation for free mappings embeddable in a flow," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 13, no. 7, pp. 1911–1915, 2003.
- [2] S. A. Andrea, "On homeomorphisms of the plane which have no fixed points," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 30, pp. 61–74, 1967.
- [3] W. Kaplan, "Regular curve-families filling the plane—I," *Duke Mathematical Journal*, vol. 7, pp. 154–185, 1940.
- [4] W. Kaplan, "Regular curve-families filling the plane—II," *Duke Mathematical Journal*, vol. 8, pp. 11–46, 1941.
- [5] Z. Leśniak, "On maximal parallelizable regions of flows of the plane," *International Journal of Pure and Applied Mathematics*, vol. 30, no. 2, pp. 151–156, 2006.
- [6] Z. Leśniak, "On parallelizability of flows of free mappings," *Aequationes Mathematicae*, vol. 71, no. 3, pp. 280–287, 2006.
- [7] N. P. Bhatia and G. P. Szegö, *Stability Theory of Dynamical Systems*, vol. 161 of *Die Grundlehren der mathematischen Wissenschaften*, Springer, New York, NY, USA, 1970.
- [8] R. C. McCann, "Planar dynamical systems without critical points," *Funkcialaj Ekvacioj*, vol. 13, pp. 67–95, 1970.
- [9] Z. Leśniak, "On parallelizable regions of flows of the plane," *Grazer Mathematische Berichte*, vol. 350, pp. 175–183, 2006.

Zbigniew Leśniak: Institute of Mathematics, Pedagogical University, Podchorążych 2,
30-084 Kraków, Poland
Email address: zlesniak@wsp.krakow.pl