

DISCRETE DIFFERENTIAL OPERATORS IN MULTIDIMENSIONAL HAAR WAVELET SPACES

CARLO CATTANI and LUIS M. SÁNCHEZ RUIZ

Received 30 July 2003

We consider a class of discrete differential operators acting on multidimensional Haar wavelet basis with the aim of finding wavelet approximate solutions of partial differential problems. Although these operators depend on the interpolating method used for the Haar wavelets regularization and the scale dimension space, they can be easily used to define the space of approximate wavelet solutions.

2000 Mathematics Subject Classification: 35A35, 42C40, 41A15.

1. Introduction. Advantage has extensively taken upon wavelets in order to analyze a number of different applications, such as image processing, signal detection, geophysics, medicine, or turbulent flows [8, 9, 10, 13, 14, 15, 18]. More mathematically focussed differential equations and even nonlinear problems have also been studied with this cargo of possibilities that wavelets have brought out into scene [1, 2, 6, 16].

In this paper, we study a Cauchy problem with the PDE $u_t = Lu$, where the unknown function $u(t, \mathbf{x}) \in L^2([0, 1])$ for each $t \in [0, T]$, $\mathbf{x} \in [0, 1]^d$, $d \in \mathbb{N}$, and L is a (nonnecessarily linear) partial differential operator whose derivatives act on the space variable. Situations like this happen quite commonly in evolution processes, such as fluid dynamics, heat transfer, elasticity, traffic flow, shock propagation, and so on [5, 12]. According to the nature and properties of L , different approaches have been developed, and even simple difference approximations may provide good results [3, 11]. However, in the presence of discontinuities some difficulties arise.

The development of *Haar wavelets* with its extreme suitability for dealing with experimental problems having piecewise constant functions as initial conditions or functions with sharp discontinuities, such as those that appear in the Riemann problem or within smooth environments, seems to provide a natural scheme to study these problems, and wavelet (approximate) solutions have been obtained by using some regular (mostly Daubechies) bases of wavelets [11].

In the following, we propose an algorithm based upon Haar wavelet basis in order to define a discrete operator that maps piecewise constant functions into piecewise constant functions, which projects the continuous operator L therein, and enables to get wavelet approximate solutions in $L^2([0, 1])$ just working with Haar wavelets.

2. Haar wavelet basis. Haar wavelets are piecewise constant functions with compact support and finite jumps at their extremal points. The drawback of not being smooth enough has been overcome by a *smoothing* process based on spline interpolation [3].

Haar functions $\varphi_k^n(x) := 2^{n/2}\varphi(2^n x - k)$ are generated from the scaling function $\varphi(x)$ which is given by the characteristic function $\chi_{[0,1]}$. Then the Haar family of wavelets $\psi_k^n(x) := \{2^{n/2}\psi(2^n x - k)\}_{k,n \in \mathbb{Z}}$ fulfills

$$\psi_k^n(x) = \begin{cases} 2^{-n/2}, & x \in \left[\frac{k}{2^n}, \frac{k+1/2}{2^n} \right), \\ -2^{-n/2}, & x \in \left[\frac{k+1/2}{2^n}, \frac{k+1}{2^n} \right), \\ 0, & \text{elsewhere,} \end{cases} \tag{2.1}$$

$\|\psi_k^n(x)\|_2 = 1$, and the multiresolution axioms [7], as well as the recursive conditions

$$2^{n+1/2}\varphi_k^n(x) = \varphi_k^{n+1}(x) + \varphi_{k+1}^{n+1}(x), \quad 2^{n+1/2}\psi_k^n(x) = \varphi_k^{n+1}(x) - \varphi_{k+1}^{n+1}(x), \quad k, n \in \mathbb{Z}. \tag{2.2}$$

Haar wavelets form a complete orthonormal system for the (finite-energy) $L^2(\mathbb{R})$ -functions [7] so that

$$\begin{aligned} L^2(\mathbb{R}) &= \bigoplus_{n \in \mathbb{Z}} W_n = V_q \oplus \bigoplus_{j \geq q} W_j, \quad q \in \mathbb{Z}, \\ V_{n+1} &= V_n \oplus W_n. \end{aligned} \tag{2.3}$$

Here, for each fixed $n \in \mathbb{Z}$, V_n is the subspace

$$V_n := \left\{ f \in L^2(\mathbb{R}) : f = \sum_{n,k \in \mathbb{Z}} c_k^n \chi_{D_k^n}, \quad c_k^n = \text{constant} \right\}, \tag{2.4}$$

where $D_k^n = [k/2^n, (k+1)/2^n)$ and W_n are orthogonal subspaces in $L^2(\mathbb{R})$, the so-called *wavelet spaces*. As usual, we consider the scalar product

$$\langle f, g \rangle := \int_{-\infty}^{+\infty} f(x)\overline{g(x)}dx \quad (\forall f, g \in L^2(\mathbb{R})). \tag{2.5}$$

According to the first equation of (2.3), we have that $f(x) = \sum_{n,k \in \mathbb{Z}} \beta_k^n \psi_k^n(x)$. If we fix a resolution value $N = 2^M$, $M \in \mathbb{N}$, in (2.3), then an approximation of the $L^2(\mathbb{R})$ -space is obtained, for example, by $L^2(\mathbb{R}) \cong V_0 \oplus_{n=0}^N W_n$, that is,

$$f(x) \cong \pi^N f(x) := \alpha_0^0 + \sum_{n=0}^{M-1} \sum_{k=0}^{2^n-1} \beta_k^n \psi_k^n(x), \tag{2.6}$$

π^N being a projection operator into V_N so that $\pi^N : L^2(\mathbb{R}) \rightarrow V_N$. In general, the coefficients α_n^k, β_k^n are given by

$$\alpha_k^n = \langle f, \varphi_k^n \rangle, \quad \beta_k^n = \langle f, \psi_k^n \rangle. \tag{2.7}$$

The dyadic discretization is the operator $\nabla^N : L^2(\mathbb{R}) \rightarrow \mathbb{K}^N \subset \ell^2$, where \mathbb{K} stands for the complex or real field, and $\nabla^N f(x) = \mathbf{f}^N = (f_0, f_1, \dots, f_{N-1})^T \in \mathbb{K}^N$ with $f_k := f(k/(N-1))$, $0 \leq k \leq N-1$.

The above notation may be simplified if we denote $\beta_N := (\alpha_0^0, \beta_0^0, \beta_0^1, \dots, \beta_{2^{M-1}-1}^{M-1})$ and $\Psi_N(x) := (\varphi(x), \psi_0^0(x), \psi_0^1(x), \dots, \psi_{2^{M-1}-1}^{M-1}(x))$ so that

$$\pi^N f(x) = \beta_N \Psi_N^T(x), \quad \mathbf{f}^N = \beta_N \nabla^N \Psi_N^T(x). \tag{2.8}$$

If P_{2^i} denotes the permutation (or shuffle) matrix (cf. [3, 4, 7]) and I is the identity matrix, then the discrete Haar wavelet transform is the one-to-one linear operator \mathcal{H} of \mathbb{K}^N onto \mathbb{K}^N such that

$$\mathcal{H}\mathbf{f}^N = \beta_N^T = \left[\prod_{k=1}^M (P_{2^k} \oplus I_{2^{M-2^k}}) (H_{2^k} \oplus I_{2^{M-2^k}}) \right] \mathbf{f}^N. \tag{2.9}$$

The matrices H_{2^i} have size $2^i \times 2^i$ and are the direct sum of the lattice coefficients of the system (2.2),

$$H_{2^i} := \begin{pmatrix} 2^{-1/2} & 2^{-1/2} \\ 2^{-1/2} & -2^{-1/2} \end{pmatrix}_1 \oplus \dots \oplus \begin{pmatrix} 2^{-1/2} & 2^{-1/2} \\ 2^{-1/2} & -2^{-1/2} \end{pmatrix}_{2^{i-1}}, \tag{2.10}$$

where, for $1 \leq j \leq 2^i - 1$ and $O_{p,q}$ standing for the $p \times q$ null matrix,

$$\begin{pmatrix} 2^{-1/2} & 2^{-1/2} \\ 2^{-1/2} & -2^{-1/2} \end{pmatrix}_j = \begin{pmatrix} O_{2j-2,2j-2} & O_{2j-2,2} & O_{2j-2,2^i-2j} \\ O_{2,2j-2} & \begin{pmatrix} 2^{-1/2} & 2^{-1/2} \\ 2^{-1/2} & -2^{-1/2} \end{pmatrix} & O_{2,2^i-2j} \\ O_{2^i-2j,2j-2} & O_{2^i-2j,2} & O_{2^i-2j,2^i-2j} \end{pmatrix}. \tag{2.11}$$

Thus, having in mind that $V_{N+1} \simeq \mathbb{K}^N$ by identifying $\alpha_0^0 + \sum_{n=0}^{M-1} \sum_{k=0}^{2^n-1} \beta_k^n \psi_k^n(x)$ with $\{\alpha_0^0, \beta_k^n\}_{k=0, \dots, 2^M-1}^{n=0, \dots, M-1}$, the projection operator $\pi^N : L^2(\mathbb{R}) \rightarrow V_{N+1}$ may be factorized by means of the following diagram:

$$\begin{array}{ccc} L^2(\mathbb{R}) & \xrightarrow{\nabla^N} & \mathbb{K}^N \\ & \searrow \pi^N & \downarrow \mathcal{H} \\ & & V_{N+1} \simeq \mathbb{K}^N \end{array} \tag{2.12}$$

The operator π^N maps each $f \in L^2(\mathbb{R})$ into a piecewise constant function $\pi^N f \in V_{N+1}$, being easy to check $\lim_{N \rightarrow \infty} \pi^N f = f$ pointwise (cf. [7]).

Note that the inverse discrete Haar transform \mathcal{H}^{-1} of (2.9) is univocally defined [7], but the inverse $\nabla^{-N} := (\nabla^N)^{-1}$ of the operator ∇^N is not because there are infinitely many functions interpolating a finite set of values.

3. Haar series regularization. In what follows, we will consider the interpolating operator $\mathcal{S}^p : \mathbb{K}^N \rightarrow C^{p-1}(\Omega)$ which maps the vector $\mathbf{f}^N (= (f_i)_{i=0}^{N-1})$, based on the dyadic nodes of $\Omega \subseteq \mathbb{R}$, into the p -differentiable function $s(x) := \mathcal{S}^p \mathbf{f}^N$, where $C^{-1}(\Omega)$ stands for the piecewise continuous functions defined on Ω . Hence $s(x)$ is an interpolating function of order p through $(i/(N-1), f_i)$, $i = 0, \dots, N-1$.

DEFINITION 3.1 (interpolating p -space). For a given resolution N , the interpolating p -space σ^p is the space $\sigma^p \subset L^2(\Omega)$ of interpolating functions with (nonredundant) fixed conditions at the boundary

$$\sigma^p := \left\{ s(x) = \mathcal{G}^p \mathbf{f}^N, \text{ with } \mathcal{G}^{-p} := (\mathcal{G}^p)^{-1} = \nabla^N, \forall \mathbf{f}^N \in \mathbb{K}^N \right\}. \tag{3.1}$$

There follows that, fixing the resolution N and the interpolating method, each function in σ^p corresponds one-to-one to a discrete set \mathbf{f}^N , making the following diagrams commutative:

$$\begin{array}{ccc} L^2(\Omega) \supset \sigma^p & \xrightarrow{\nabla^N} & \mathbb{K}^N \\ & \searrow \pi^N & \downarrow \mathcal{H} \\ & & V_{N+1} \end{array} \qquad \begin{array}{ccc} L^2(\Omega) \supset \sigma^p & \xleftarrow{\mathcal{G}^p} & \mathbb{K}^N \\ & \swarrow \pi^{-N} & \uparrow \mathcal{H}^{-1} \\ & & V_{N+1} \end{array} \tag{3.2}$$

THEOREM 3.2. For a fixed resolution N and a given interpolating method, the inverse π^{-N} of the projection operator π^N is univocally defined when acting on the σ^p space functions by means of

$$\pi^{-N} = \mathcal{G}^p \mathcal{H}^{-1} \quad (\pi^N = \mathcal{H} \nabla^N). \tag{3.3}$$

PROOF. Note that $\nabla^N \mathcal{G}^p = \mathcal{G}^p \nabla^N = I$ and \mathcal{H}^{-1} is univocally defined [7]. □

DEFINITION 3.3. The q -derivative of a p -interpolating function, $0 \leq q \leq p$, at a given dimension N is the linear operator $\delta^{(p,q)} : \mathbb{K}^N \rightarrow V_{N+1}$ such that

$$\delta^{(p,q)} := \pi^N \frac{d^q}{dx^q} \mathcal{G}^p. \tag{3.4}$$

This definition provides an algorithm that maps piecewise constant functions into piecewise constant functions. If we look at it, we may note that firstly it smoothens piecewise constant functions with a suitable interpolation; secondly it finds out the derivative of the interpolating function, and then this derivative is transformed into a piecewise constant function by the discrete Haar wavelet transform.

From [7, page 13] and since $(d^q/dx^q) \mathcal{G}^p \nabla^N f$ belongs to $L^2(\Omega)$ for each $f \in L^2(\Omega)$, it follows that $\|(d^q/dx^q) \mathcal{G}^p \nabla^N f - \delta^{(p,q)} \nabla^N f\|_{L^2}^2 \leq 2^{-2M-1} 2^{\mu(\Omega)} K + \varepsilon$, where μ stands for the Lebesgue measure, $K = \sum_{k=0}^{M-1} ((m_{0,k,N})^2 + (m_{-1,k,N})^2)^{1/2}$, with $m_{0,k,N}$ and $m_{-1,k,N}$ being the average value of $(d^q/dx^q) \mathcal{G}^p \nabla^N f$ over the first and the second half of D_k^N , respectively, and ε may be taken as small as desired since any function in $L^2(\Omega)$ may be approximated by functions with compact support which are piecewise constant.

It is easy to see that if $f = \sum_{k=0}^{M-1} f_k \chi_{D_k}$, $g = \sum_{k=0}^{M-1} g_k \chi_{D_k} \in V_N$, where f_k, g_k are constants, then there are unique c_k such that $f_k = g_k + c_k$. Hence, there exists a matrix $A^{(p,q)}$ depending on N that makes

$$\begin{aligned} \delta^{(p,q)} \nabla^N f &= A^{(p,q)} \nabla^N f, \\ \delta^{(p,q)} \mathbf{f}^N &= A^{(p,q)} \beta_N \nabla^N \Psi_N^T(x). \end{aligned} \tag{3.5}$$

At the end of this paper, we provide an appendix with some of these matrices A for different spline interpolations.

4. Multidimensional wavelet solution. In a d -dimensional variable space $\mathbf{x} = (x_1, \dots, x_d)$, we restrict ourselves to considering the same dyadic discretization on each coordinate of \mathbf{x} so that the natural generalization of the dyadic discretization operator $\nabla^N : L^2(\mathbb{R}^d) \rightarrow \prod_{k=1}^d \mathbb{K}^N$ ($N = 2^M$) is given by

$$\nabla^N f(\mathbf{x}) := f\left(\frac{i_1}{N-1}, \frac{i_2}{N-1}, \dots, \frac{i_d}{N-1}\right), \quad 0 \leq i_j \leq N-1, \quad 1 \leq j \leq d. \tag{4.1}$$

By using tensor products, we define $\mathbf{V}_N(\mathbf{x}) := \otimes_{i=1}^d V_N(x_i)$, where the (orthonormal) wavelet basis is $\{\psi_k^n(x_1) \otimes \dots \otimes \psi_k^n(x_d)\}$, $0 \leq k \leq N-1$ (cf. [17]). Thus $\mathbf{V}_N(\mathbf{x})$ is the space of the piecewise constant functions on the d -dimensional product $D_k^N \times D_h^N$, $0 \leq h, k \leq 2^N - 1$, which can be represented as

$$\begin{aligned} \pi^N f(\mathbf{x}) := & \prod_{1 \leq i \leq d} \alpha_i \varphi(x_i) + \sum_{\substack{0 \leq k_i \leq 2^{n_i-1} \\ 0 \leq n_i \leq M \\ 1 \leq i \leq d}} \prod_{1 \leq i \leq d} \beta_{k_i}^{n_i} \psi_{k_i}^{n_i}(x_i) \\ & + \sum_{\substack{0 \leq k_i \leq 2^{n_i-1} \\ 0 \leq n_i \leq M \\ 1 \leq i \leq d}} \prod_{\substack{1 \leq i, j \leq d \\ i \neq j}} \omega_{k_i, j}^{n_i} \varphi(x_j) \psi_{k_i}^{n_i}(x_i), \end{aligned} \tag{4.2}$$

in short, $\pi^N f(\mathbf{x}) := \sum_{0 \leq i_j \leq M-1, 1 \leq j \leq d} \beta_N^{i_1 i_2 \dots i_d} \nabla^N (\psi_N(x_{i_1}) \otimes \dots \otimes \psi_N(x_{i_d}))$.

Then the discrete d -dimensional Haar wavelet transform is the one-to-one linear operator $\mathcal{H}_d = \mathcal{H} \otimes \dots \otimes \mathcal{H}$ of $(\mathbb{K}^N)^d$ onto $(\mathbb{K}^N)^d$, and the discrete derivative of $\nabla^N f(\mathbf{x})$ is the tensor product of the corresponding 1-dimensional discrete derivative. Hence, in the 2-dimensional case, this leads to a matrix $A^{(p,q)}$, which in fact is the same as the one of the 1-dimensional case, such that

$$\begin{aligned} \delta_x^{(p,q)} f\left(\frac{i_1}{N-1}, \frac{i_2}{N-1}\right) &= A^{(p,q)} f\left(\cdot, \frac{i_2}{N-1}\right), \quad 0 \leq i_2 \leq N-1, \\ \delta_y^{(p,q)} f\left(\frac{i_1}{N-1}, \frac{i_2}{N-1}\right) &= A^{(p,q)} \left(\frac{i_1}{N-1}, \cdot\right), \quad 0 \leq i_1 \leq N-1, \end{aligned} \tag{4.3}$$

proving that if $x_1 = i_1 / (N-1)$, $x_2 = i_2 / (N-1)$,

$$\delta_x^{(p,q)} \nabla^N f(x_1, x_2) = A^{(p,q)} \beta_N \nabla^N \Psi_N^T(x_1) \otimes \nabla^N \Psi_N^T(x_2), \tag{4.4}$$

and analogously for $\delta_y^{(p,q)}$. Taking this into account, we obtain the following result.

PROPOSITION 4.1. *An approximate wavelet solution $\tilde{u}(t, \mathbf{x}) = \pi^N u(t, \mathbf{x})$ of the Cauchy problem*

$$u_t = Lu, \quad u(0, \mathbf{x}) = u_0(\mathbf{x}), \quad \mathbf{x} \in [0, 1]^d, \quad t \in [0, T], \tag{4.5}$$

is obtained by taking

$$\tilde{u}(t, \mathbf{x}) := \sum_{\substack{0 \leq i_j \leq M-1 \\ 1 \leq j \leq d}} \beta_N^{i_1 i_2 \dots i_d}(t) \nabla^N (\psi_N(x_{i_1}) \otimes \dots \otimes \psi_N(x_{i_d})), \tag{4.6}$$

where the wavelet coefficients are explicit functions of t .

PROOF. From (4.5), it follows that $\pi_N u_t = \pi_N Lu$. Hence, if L_δ denotes the operator acting on piecewise functions obtained by replacing the partial derivatives that appear on L with the corresponding $\delta_{i_j}^{(p,q)} \nabla^N$, for example, if $L = L(\partial/\partial x, \partial^2/\partial x \partial y)$, then $L_\delta = L(\delta_x^{(p,1)} \nabla^N, \delta_y^{(p,1)} \delta_x^{(p,1)} \nabla^N)$, it happens that the projection in $\mathbf{V}_N(\mathbf{x})$ must verify $\tilde{u}_t(t, \mathbf{x}) = L_\delta \tilde{u}(t, \mathbf{x})$ and

$$\sum_{\substack{0 \leq i_j \leq M-1 \\ 1 \leq j \leq d}} \beta_N^{i_1 i_2 \dots i_d}(0) \nabla^N (\psi_N(x_{i_1}) \otimes \dots \otimes \psi_N(x_{i_d})) = \nabla^N f(\mathbf{x}). \tag{4.7}$$

Orthonormality of the scaling functions φ and ψ_k^n gives an easily solved first-order ordinary system in the unknown wavelet coefficients. Note that due to the definition of Haar wavelets, the boundary conditions are automatically fulfilled. □

5. Spline Haar solution of a parabolic equation. In this section, we will compare the Haar regularized solution of the following 1-dimensional parabolic problem with the exact solution and the one obtained with Crank-Nicholson’s method.

EXAMPLE 5.1. Compare at time 0.04 and 0.08 the above-mentioned methods in the heat equation, without exchanges at the boundary, and a periodic function as initial function given by

$$\begin{aligned} u_t &= u_{xx}, \quad 0 \leq x \leq 1, \quad t \geq 0, \\ u(x, 0) &= \sin(\pi x), \\ u(0, t) &= u(1, t) = 0. \end{aligned} \tag{5.1}$$

In Figure 5.1, the exact solution $u(x, t) = \sin(\pi x)e^{-\pi^2 t}$ at $t = 0.01$ and the one obtained by the Haar regularized method based on 64 nodes are shown as functions of the variable distance, x . Table 5.1 compares the set of values that we obtain (HR) at time $t = 0.04$, $t = 0.08$ with the exact ones (E) and those obtained by Crank-Nicholson’s method (CN).




FIGURE 5.1. Exact and Haar regularized solution with 64 nodes.

TABLE 5.1

x	t = 0.04			t = 0.08		
	E	HR	CN	E	HR	CN
0	0	0	0	0	0	0
0.2	0.396065	0.398	0.399	0.266878	0.269	0.271
0.4	0.640846	0.648	0.646	0.431818	0.440	0.439
0.6	0.640846	0.648	0.646	0.431818	0.440	0.439
0.08	0.396065	0.398	0.399	0.266878	0.269	0.271
1	0	0	0	0	0	0

6. Annex. Finally, we enclose a list of matrices $A^{(p,q)}$, announced at the end of Section 3, which satisfy $\delta^{(p,q)} \nabla^N f = A^{(p,q)} \nabla^N f$. We fix our attention on the case $p = 3$, $q = 1$. For this and different resolutions $N = 2^M$, $M \in \mathbb{N}$, the matrices $A^{(p,q)}$ are computed according to the following steps:

- (1) discretize the interval $[0, 1]$ at the dyadic points $k/(N - 1)$, $0 \leq k \leq N - 1$,
- (2) consider the points whose first coordinates are the dyadic points and whose second coordinates are arbitrary values a_0, a_1, \dots, a_{N-1} , that is, $(0, a_0), (1/(N - 1), a_1), \dots, (1, a_{N-1})$,
- (3) build up the cubic spline that goes through the aforementioned set of points,
- (4) compute the first derivative of this spline and evaluate it at the dyadic points,
- (5) assuming the above evaluations generate the points $(0, a'_0), (1/(N - 1), a'_1), \dots, (1, a'_{N-1})$, then $A^{(3,1)}$ is the $N \times N$ matrix that satisfies

$$\begin{pmatrix} a'_0 \\ a'_1 \\ \vdots \\ a'_{N-1} \end{pmatrix} = A^{(3,1)} \begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_{N-1} \end{pmatrix}. \tag{6.1}$$

With a bound error of 10^{-5} , routine calculations make $A^{(3,1)}$ become

$$\begin{pmatrix} -0.16667 & 0.16667 \\ -0.16667 & 0.16667 \end{pmatrix},$$

$$N = 2^1 = 2,$$

$$\begin{pmatrix} -0.97941 & 1.24412 & -0.34412 & 0.07941 \\ -2.14412 & 0.29118 & 2.40882 & -0.55588 \\ 0.55588 & -2.40882 & -0.29118 & 2.14412 \\ -0.07941 & 0.34412 & -1.24412 & 0.97941 \end{pmatrix},$$

$$N = 2^2 = 4,$$

$$\begin{pmatrix} -3.96693 & 5.02851 & -1.33162 & 0.31661 & -0.04656 & 0 & 0.00401 & -0.00401 \\ -4.56556 & 0.16762 & 5.51672 & -1.31167 & 0.19289 & 0 & -0.01662 & 0.01662 \\ 1.22919 & -5.69897 & 0.26473 & 4.93006 & -0.72501 & 0 & 0.06246 & -0.06246 \\ -0.35120 & 1.62828 & -6.57564 & 2.59141 & 2.70715 & 0 & -0.23324 & 0.23324 \\ 0 & 0 & 0 & 0 & -10.5 & 14 & -2.62951 & -0.87049 \\ 0 & 0 & 0 & 0 & -3.5 & 0 & 0.25129 & 3.24871 \\ 0 & 0 & 0 & 0 & 0 & 0 & -21 & 21 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$N = 2^3 = 8.$$

(6.2)

ACKNOWLEDGMENT. The second author is supported by Generalitat Valenciana.

REFERENCES

- [1] E. Bacry, S. Mallat, and G. Papanicolaou, *A wavelet based space-time adaptive numerical method for partial differential equations*, RAIRO Modél. Math. Anal. Numér. **26** (1992), no. 7, 793-834.
- [2] W. Cai and J. Wang, *Adaptive multiresolution collocation methods for initial-boundary value problems of nonlinear PDEs*, SIAM J. Numer. Anal. **33** (1996), no. 3, 937-970.
- [3] C. Cattani, *Haar wavelet splines*, J. Interdiscip. Math. **4** (2001), no. 1, 35-47.
- [4] ———, *Haar wavelet-based technique for sharp jumps classification*, Math. Comput. Modelling **39** (2004), no. 2-3, 255-278.
- [5] R. Courant and K. O. Friedrichs, *Supersonic Flow and Shock Waves*, Interscience Publishers, New York, 1948.
- [6] W. Dahmen, *Wavelet methods for PDEs—some recent developments*, J. Comput. Appl. Math. **128** (2001), no. 1-2, 133-185.
- [7] I. Daubechies, *Ten Lectures on Wavelets*, CBMS-NSF Regional Conference Series in Applied Mathematics, vol. 61, Society for Industrial and Applied Mathematics, Pennsylvania, 1992.
- [8] L. Debnath, *Wavelet transforms and their applications*, Proc. Indian Nat. Sci. Acad. Part A **64** (1998), no. 6, 685-713.
- [9] R. A. DeVore, B. Jawerth, and B. J. Lucier, *Image compression through wavelet transform coding*, IEEE Trans. Inform. Theory **38** (1992), no. 2, 719-746.
- [10] D. L. Donoho, *Nonlinear wavelet methods for recovery of signals, densities, and spectra from indirect and noisy data*, Different Perspectives on Wavelets (San Antonio, Tex,

- 1993), Proc. Sympos. Appl. Math., vol. 47, American Mathematical Society, Rhode Island, 1993, pp. 173-205.
- [11] S. Lazaar, P. Ponenti, J. Liandrat, and P. Tchamitchian, *Wavelet algorithms for numerical resolution of partial differential equations*, Comput. Methods Appl. Mech. Engrg. **116** (1994), no. 1-4, 309-314.
- [12] R. J. LeVeque, *Numerical Methods for Conservation Laws*, 2nd ed., Lectures in Mathematics ETH Zürich, Birkhäuser, Basel, 1992.
- [13] P. J. Oonincx, *A wavelet method for detecting S-waves in seismic data*, Comput. Geosci. **3** (1999), no. 2, 111-134.
- [14] W. Schempp, *Wavelet modelling of high resolution radar imaging and clinical magnetic resonance tomography*, Results Math. **31** (1997), no. 3-4, 195-244.
- [15] J. Segman and Y. Y. Zeevi, *Image analysis by wavelet-type transforms: group-theoretic approach*, J. Math. Imaging Vision **3** (1993), no. 1, 51-77.
- [16] O. V. Vasilyev and S. Paolucci, *A fast adaptive wavelet collocation algorithm for multidimensional PDEs*, J. Comput. Phys. **138** (1997), no. 1, 16-56.
- [17] P. Wojtaszczyk, *A Mathematical Introduction to Wavelets*, London Mathematical Society Student Texts, vol. 37, Cambridge University Press, Cambridge, 1997.
- [18] V. Zimin and F. Hussain, *Wavelet based model for small-scale turbulence*, Phys. Fluids **7** (1995), no. 12, 2925-2927.

Carlo Cattani: Dipartimento di Scienze Farmaceutiche (DiFARMA), Università degli Studi di Salerno, Via Ponte Don Melillo, Invariante 11/C, 84084 Fisciano (Salerno), Italy
E-mail address: ccattani@unisa.it

Luis M. Sánchez Ruiz: Departamento de Matemática Aplicada, Escuela Técnica Superior de Ingeniería de Diseño (ETSID), Universidad Politécnica de Valencia, 46022 Valencia, Spain
E-mail address: lmsr@mat.upv.es

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru