

SEPARABLE SUBALGEBRAS OF A CLASS OF AZUMAYA ALGEBRAS

GEORGE SZETO

Department of Mathematics
Bradley University
Peoria, Illinois 61625, U.S.A.

(Received July 15, 1996 and in revised form April 14, 1997)

ABSTRACT. Let S be a ring with 1, C the center of S , G a finite automorphism group of S of order n invertible in S , and S^G the subring of elements of S fixed under each element in G . It is shown that the skew group ring $S * G$ is a G' -Galois extension of $(S * G)^{G'}$ that is a projective separable C^G -algebra where G' is the inner automorphism group of $S * G$ induced by G if and only if S is a G -Galois extension of S^G that is a projective separable C^G -algebra. Moreover, properties of the separable subalgebras of a G -Galois H -separable extension S of S^G are given when S^G is a projective separable C^G -algebra.

KEY WORDS AND PHRASES: Azumaya algebras, Galois extensions, H -separable extensions, Skew group rings.

1991 AMS SUBJECT CLASSIFICATION CODES: 16S30; 16W20.

1. INTRODUCTION

DeMeyer [1] and Kanzaki [2] studied central Galois algebras and Galois extensions whose center is a Galois algebra with Galois group induced by and isomorphic with the group of the extension. These two types of Galois extensions were recently generalized to a bigger class of Galois Azumaya extensions [3] where S is called a G -Galois Azumaya extension of S^G if S is a G -Galois extension of S^G that is an Azumaya C^G -algebra where C is the center of S and S^G is the subring of elements fixed under each element of G . Sugano [4] investigated a G -Galois H -separable extension of S^G , and recently, Szeto [5] proved that a G -Galois H -separable extension S of S^G that is a projective separable C^G -algebra if and only if S is a C^G -Azumaya algebra. We call such an S a GHS-extension. It will be shown that the skew group ring $S * G$ is a G' HS-extension if and only if S is a G -Galois extension of S^G that is a projective separable C^G -algebra, where G' is the inner automorphism group of $S * G$ induced by G . Moreover, properties of some separable subalgebras of a GHS-extension are also given.

2. PRELIMINARIES

Throughout, S is a ring with 1, G a finite automorphism group of S of order n invertible in S , C the center of S , and S^G the subring of elements fixed under each element in G . S is called a separable extension of a subring T if there exist $\{a_i, b_i \text{ in } S / i = 1, 2, \dots, m\}$ for some integer m such that $\sum a_i b_i = 1$ and $\sum s a_i \otimes b_i = \sum a_i \otimes b_i s$ for each s in S where \otimes is over T . We call $\{a_i, b_i\}$ a separable system for S . S is called an H -separable extension of T if $S \otimes_T S$ is isomorphic with a direct summand of a finite direct sum of S as a bimodule over S . It is known that an H -separable extension is a separable extension and an Azumaya algebra is an H -separable extension. S is called a G -Galois extension of S^G , if there exist $\{c_i, d_i / i = 1, 2, \dots, k\}$ in S for some integer k such that $\sum c_i d_i = 1$ and $\sum c_i g(d_i) = 0$ for each $g \neq 1$ in G . We call $\otimes \{c_i, d_i\}$ a G -Galois system for S .

3. SKEW GROUP RINGS

In this section, we shall show that $S * G$ is a G'HS-extension if and only if S is a G-Galois extension of S^G that is a projective separable C^G -algebra, and give some properties of the separable subalgebras of an G'HS-extension skew group ring.

THEOREM 3.1. By keeping the notations of section 2, $S * G$ is a G'HS-extension if and only if S is a G-Galois extension of S^G that is a projective separable C^G -algebra, where G' is the inner automorphism group of $S * G$ induced by G .

PROOF. Let S be a G-Galois extension of S^G that is a projective separable C^G -algebra. Noting that S is a subring of $S * G$, we have that $S * G$ is also a G' -Galois extension of $(S * G)^{G'}$ with a same Galois system as S where G' is the inner automorphism group of $S * G$ induced by G such that the restriction of G' to S is G . Hence $S * G$ is an H-separable extension of $(S * G)^{G'}$ ([4], Corollary 3). Moreover, since n is a unit in S , $S * G$ is a separable extension of S . But $S * G$ is a free module over S and S is a G-Galois extension of S^G that is a projective separable C^G -algebra by hypothesis, so $S * G$ is a projective separable C^G -algebra by the transitivity of projective separable extensions. Since the order of G' is n , it is easy to see that $(S * G)^{G'}$ is a direct summand of $S * G$ as a two sided $(S * G)^{G'}$ -module. Noting that $S * G$ is finitely generated and projective module as a right $(S * G)^{G'}$ -module or a left module, we have that $(S * G)^{G'}$ is a projective separable C^G -algebra by the same argument as given in the proof of Lemma 2 in [1]. This completes the sufficiency.

For the necessity, $S * G$ is a projective separable C^G -algebra by the transitivity of projective separable extensions because $S * G$ is a G' -Galois extension of $(S * G)^{G'}$ that is a projective separable algebra of C^G . Hence $S * G$ is an Azumaya algebra of its center Z . But S is a free module over S and n is a unit in S , so S is a finitely generated and projective left $S * G$ -module by the proof of Proposition 2.3 in [6] where $gs = g(s)$ for each s in S and g in G . Thus S is a finitely generated and projective Z -module by the transitivity of finitely generated and projective modules. Noting that 1 is in C^G and that C^G is contained in Z , we have that S is a faithful Z -module. Thus S is a progenerator over Z . Since $S * G$ is an Azumaya Z -algebra, S is a progenerator over $S * G$. Therefore, S is a G-Galois extension of S^G . Moreover, since S is a direct summand of $S * G$ as a C^G -module and $S * G$ is a finitely generated and projective C^G -module, S is also a finitely generated and projective C^G -module. Now n is a unit in S , so S^G is a S^G -direct summand of S . This implies that S^G is a finitely generated and projective C^G -module. So, it suffices to show that S^G is a separable C^G -algebra. In fact, since S is a progenerator over C^G (for 1 is in C^G), $\text{Hom}_{C^G}(S, S)$ is an Azumaya C^G -algebra. But $S^G \approx \text{Hom}_{S * G}(S, S) =$ the commutator of $S * G$ in $\text{Hom}_{C^G}(S, S)$, so S^G is a separable C^G -algebra (for so is $S * G$) by the commutant theorem for Azumaya algebras ([6], Theorem 4.3).

Next we give some properties of the separable subalgebras of $S * G$.

COROLLARY 3.2. If $S * G$ is a G'HS-extension, then, for any subgroup K of G , $S * K$ is a K' -Galois extension of $(S * K)^{K'}$ that is a separable C^G -algebra where K' is the inner automorphism group of $S * K$ induced by K .

PROOF. By Theorem 3.1, S is a G-Galois extension of S^G that is a projective C^G -algebra, so S is a K -Galois extension of S^K . Hence $S * K$ is a K' -Galois of $(S * K)^{K'}$. Noting that the order of K' is a unit in S , we have that $(S * K)^{K'}$ is a direct summand of $S * K$ as a $(S * K)^{K'}$ -module. But $S * K$ is a projective separable C^G -algebra, so $(S * K)^{K'}$ is a separable C^G -algebra by the same argument as given in the proof of Lemma 2 in [1].

Let $V_S(T)$ be the commutator subring of the subring T in S , and Z the center of $S * G$. We give an expression of the commutator subring of $(S * G)^{K'}$ in $S * G$.

THEOREM 3.3. If S^*G is a GHS-extension, then (1) for any subgroup K of G , $V_{S^*G}((S^*G)^K)$ is ZK , and (2) ZK is an Azumaya algebra over its center D such that $D = D^{K'}$.

PROOF. (1) By Theorem 3.1, S^*G is a separable C^G -algebra, so S^*G is an Azumaya Z -algebra. Since n is a unit in S , the order of K is a unit in S ; and so ZK is a separable Z -algebra contained in S^*G . Noting that $V_{S^*G}(ZK) = (S^*G)^{K'}$, we have that $(S^*G)^{K'}$ is a separable Z -subalgebra of S^*G such that $ZK = V_{S^*G}((S^*G)^{K'})$ by the commutant theorem for Azumaya algebras ([6], Theorem 4.3).

(2) Since S^*G is a separable C^G -algebra, Z is a separable C^G -algebra. Hence ZK is a separable C^G -algebra (for the order of K is a unit in Z). Thus ZK is an Azumaya D -algebra. It remains to show that $D = D^{K'}$. Clearly, $D^{K'} \subset D$. Conversely, let d be an element in D . Then $gd = dg$ for each g in K , so $gdg^{-1} = d$ for each g in K . Hence d is in $D^{K'}$.

The following consequences are immediate.

COROLLARY 3.4. Let S^*G be an GHS-extension. If K is an abelian subgroup of G , then $(S^*G)^{K'}$ is an Azumaya ZK -algebra.

PROOF. By the proof of Corollary 3.3, $(S^*G)^{K'}$ and ZK are separable subalgebras of the Azumaya Z -algebra S^*G such that $V_{S^*G}(ZK) = (S^*G)^{K'}$ and $V_{S^*G}((S^*G)^{K'}) = ZK$, so ZK is contained in the center of $(S^*G)^{K'}$ and the center of $(S^*G)^{K'}$ is contained in ZK . Thus ZK is the center of $(S^*G)^{K'}$.

COROLLARY 3.5. Let S^*G be a GHS-extension. Then (1) if $(S^*G)^{K'}$ is a commutative ring, then ZK is an Azumaya $(S^*G)^{K'}$ -algebra, and (2) if $(S^*G)^{K'}$ and ZK are commutative, then ZK is a splitting ring for the Azumaya Z -algebra S^*G .

PROOF. (1) It is immediate by the same argument of Corollary 3.4-(1). (2) Since $(S^*G)^{K'}$ and ZK are separable subalgebras of the Azumaya Z -algebra S^*G such that $V_{S^*G}((S^*G)^{K'}) = ZK$ and $V_{S^*G}(ZK) = (S^*G)^{K'}$, $(S^*G)^{K'}$ and ZK are commutative) such that $V_{S^*G}(ZK) = ZK$. Hence ZK is a maximal commutative separable subalgebra of S^*G . Thus ZK is a splitting ring for the Azumaya algebra S^*G ([6], Theorem 5.5).

4. SEPARABLE ALGEBRAS

In this section, we shall give a property of a separable subalgebra of any GHS-extension similar to Theorem 3.3. Let T be a subalgebra of S over C^G . The commutator subring of T in S is denoted by T' .

THEOREM 4.1. Let S be a GHS-extension and T a separable C^G -subalgebra of S , and $K = \{g \text{ in } G / g(t) = t \text{ for each } t \text{ in } T\}$. Then T' is invariant under K and an Azumaya $D^{K''}$ -algebra, where D is the center of T' and K'' is the restriction of K to T' .

PROOF. Since $tt' = t't$ for each t in T and t' in T' , $tg(t') = g(t')t$ for each g in K . Hence T' is invariant under K .

Next, since S is a GHS-extension, $V_S(V_S(S^G)) = S^G$ ([4], Proposition 4-1). Hence $C = V_S(S) \subset V_S(V_S(S^G)) = S^G$. This implies $C = C^G$. Noting that S is a separable C^G -algebra (for S is a GHS-extension), we have that S is an Azumaya C^G -algebra. But T is a separable subalgebra of S , so T' is also a separable subalgebra of S such that $V_S(T') = T'$ by the commutant theorem for Azumaya algebras ([6], Theorem 4.3). Let D be the center of T' . Then $D \subset V_S(T') = T' \subset S^{K''}$. Thus $D = D^{K''}$ where K'' is the restriction of K to T' . The proof is complete.

COROLLARY 4.2. Let S be a GHS-extension, T a separable C^G -algebra of S , and $N = \{g \text{ in } G / g(t') = t' \text{ for each } t' \text{ in } V_S(T')\}$. Then T' is invariant under N and an Azumaya $E^{N''}$ -algebra, where E is the center of T' and N'' is the restriction of N to T' .

PROOF. By the proof of Theorem 4.1, T and T' ($= V_S(T')$) are separable subalgebras of the Azumaya C^G -algebra S such that $T = V_S(T')$, so the corollary is immediate by Theorem 4.1.

By Theorem 3.1 in [5], Corollary 4.2 implies the following consequence.

COROLLARY 4.3. By keeping the notations and hypotheses of Corollary 4.2, T is a N'' HS-extension.

We conclude the paper with two examples: (1) S is a G -Galois extension of S^G that is a projective separable C^G -algebra, but not an H -separable extension of S^G , and (2) S is a GHS-extension.

Example 1. Let Q be the rational field, $Q[\sqrt{2}]$ the G -Galois extension of Q with Galois group $G = \{1, g\}$ where $g(\sqrt{2}) = -\sqrt{2}$ and $S = M_2(Q[\sqrt{2}])$, the matrix ring of order 2 over $Q[\sqrt{2}]$.

Let $G' = \{1, g'\}$ where $g'([a_{ij}]) = [g(a_{ij})]$ for all $[a_{ij}]$ in S . Then

- (1) S is a G' -Galois extension of $S^{G'}$,
- (2) $S^{G'} = M_2(Q)$, the matrix ring of order 2 over Q ,
- (3) $S^{G'}$ is a projective separable Q -algebra,
- (4) the center C of S is $Q[\sqrt{2}]$ and $C^G = Q$, and
- (5) S is not an H -separable extension of $S^{G'}$ because $C \neq C^G$.

Example 2. Let S and G' be given by Example 1. Then S is a G' -Galois extension of $S^{G'}$ that is a projective separable C^G -algebra by properties (1) through (4) in Example 1. Then the skew group ring $S * G'$ is a GHS-extension by Theorem 3.1.

ACKNOWLEDGMENT. This work was done under the support of a Caterpillar Fellowship at Bradley University. The paper is revised under the suggestions of the referee. The author would like to thank him for his valuable suggestions.

REFERENCES

- [1] DEMEYER, F. R., Some Notes On The General Galois Theory Of Rings, *Osaka J. Math.* **2** (1965), 117-127.
- [2] KANZAKI, T., On Galois Algebra Over A Commutative Ring, *Osaka J. Math.* **2** (1965), 309-317.
- [3] ALFARO, R. and SZETO, G., Skew Group Rings Which Are Azumaya, *Comm. In Algebra* **23** (6), (1995), 2255-2261.
- [4] SUGANO, K., On A Special Type Of Galois Extensions, *Hokkaido J. Math.* **9** (1980), 123-128.
- [5] SZETO, G., On A Class Of Azumaya Algebras, *Math. Japon.* (to appear).
- [6] DEMEYER, F. R. and INGRAHAM, E., *Separable Algebras Over Commutative Rings* **181**, Springer-Verlag, Berlin, 1971.
- [7] IKEHATA, S., Note On Azumaya Algebras And H -Separable Extensions, *Math. J. Okayama Univ.* **23** (1981), 17-18.
- [8] OKAMOTO, H., On Projective H -separable Extensions Of Azumaya Algebras, *Results In Mathematics* **14** (1988), 330-332.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk