

ON ISOMORPHISMS AND HYPER-REFLEXIVITY OF CLOSED SUBSPACE LATTICES

HAN DEGUANG

Department of Mathematics
Qufu Normal University, Shandong
P.R. China

(Received March 15, 1988 and in revised form January 2, 1991)

ABSTRACT. There are some papers, such as [1], [2] and [3], in which some properties on isomorphism of closed subspace lattices of Hilbert spaces were studied. In this short paper we will point out that the hyper-reflexivity of closed subspace lattice is invariant under isomorphism of $\xi(H_1)$ on $\xi(H_2)$. We also proved that if T is in $L(H)$ such that $0 \in \pi(T)$ and \mathfrak{F} is a hyper-reflexive subspace lattice, then $\phi_T(\mathfrak{F}) \cup \{H\}$ is hyper-reflexive where ϕ_T is a homomorphism induced by T .

KEY WORDS AND PHRASES: Reflexivity, Hyper-reflexivity, Isomorphism.

1980 AMS SUBJECT CLASSIFICATION CODES. 46C10, 47D25.

1. INTRODUCTION.

Let H be a complex Hilbert space, $L(H)$ denotes the set of all bounded linear operators on H and let $\xi(H)$ be the set of all closed subspaces of H . For any subset \mathcal{A} of $L(H)$ and any family $\mathfrak{F} \subseteq \xi(H)$, let $\text{Lat } \mathcal{A}$ denote the lattice of closed subspaces invariant for \mathcal{A} (or the lattice of invariant projections for \mathcal{A}) and let $\text{Alg } \mathfrak{F}$ be the set of all operators invariant for \mathfrak{F} . \mathfrak{F} is said to be reflexive if $\text{Lat Alg } \mathfrak{F} = \mathfrak{F}$. A subalgebra \mathcal{A} of $L(H)$ is said to be reflexive if $\mathcal{A} = \text{Alg Lat } \mathcal{A}$.

Let \mathcal{A} be a reflexive algebra and let $T \in L(H)$. It is easy to see that

$$\text{dist}(T, \mathcal{A}) \geq \sup \{ \| P^\perp T P \| : P \in \text{Lat } \mathcal{A} \}.$$

\mathcal{A} is called to be hyper-reflexive (See [4]) if there exists a constant $K > 0$ such that for any $T \in L(H)$

$$\text{dist}(T, \mathcal{A}) \leq K \sup \{ \| P^\perp T P \| : P \in \text{Lat } \mathcal{A} \}.$$

For any subspace lattice $\mathfrak{F} \subseteq \xi(H)$, if \mathfrak{F} is reflexive and $\text{Alg } \mathfrak{F}$ is hyper-reflexive, then \mathfrak{F} is said to be hyper-reflexive. Let ϕ be a lattice isomorphism of $\xi(H_1)$ onto $\xi(H_2)$ (i.e., a bijection with the

property that $M = N$ iff $\phi(M) = \phi(N)$). It was proved in [1] and [2] that \mathfrak{F} is reflexive if and only if $\phi(\mathfrak{F})$ is reflexive. For an operator $A \in L(H)$, A can give rise to a map $\phi_A: \xi(H) \rightarrow \xi(H)$ given by $\phi_A(M) = \overline{AM}$, where ' - ' denotes norm closure. In the paper [3] the author proved that if the approximate point spectrum $\pi(A)$ of A does not contain 0 and \mathfrak{F} is reflexive, then $\phi_A(\mathfrak{F}) \cup \{H\}$ is also reflexive. Here we prove the following theorem:

LEMMA 1. Let H_1 and H_2 be two complex Hilbert spaces, and let ϕ be a lattice isomorphism of $\xi(H_1)$ onto $\xi(H_2)$. Then a subspace lattice \mathfrak{F} of $\xi(H_1)$ is hyper-reflexive if and only if $\phi(\mathfrak{F})$ is hyper-reflexive.

THEOREM 2. Let H be a complex Hilbert space, $A \in L(H)$ and $0 \notin \pi(A)$. If the subspace lattice \mathfrak{F} of $\xi(H)$ is hyper-reflexive, then so is $\phi_A(\mathfrak{F}) \cup \{H\}$.

2. THE PROOF OF THE THEOREMS.

Lemma 1 may be known, we give a proof by the following theorem:

THEOREM A. ([4]) Let $\mathcal{A} \subseteq L(H)$ be a σ -weakly closed unital subalgebra of $L(H)$. Then \mathcal{A} is hyper-reflexive iff every element $f \in \mathcal{A}_\perp$ has a representation

$$f = \sum_{n=1}^{\infty} f_n$$

where $\mathcal{A}_\perp = \left\{ f: f \text{ is a } \sigma\text{-weakly continuous linear functional on } L(H) \text{ and } f(\mathcal{A}) = \{0\} \right\}$, each f_n is an elementary functional in \mathcal{A}_\perp and $\sum_{n=1}^{\infty} \|f_n\| < \infty$.

REMARK. A σ -weakly continuous functional f on $L(H)$ is said to be elementary if there exist $x, y \in H$ such that $f(T) = (Tx, y)$ for any $T \in L(H)$. We write $f = (x \otimes y)$.

Let S be a conjugate linear continuous map from H_1 , into H_2 . It can be defined uniquely the adjoint S^* of S by the formula

$$(S^*x, y) = (Sy, x) = \overline{(x, Sy)}. \tag{2.1}$$

It is easy to check that $(S^*)^* = S$, and $(S^{-1})^* = (S^*)^{-1}$ when S has continuous inverse.

PROOF OF LEMMA 1. It is sufficient to prove the necessity. First $\phi(\mathfrak{F})$ is reflexive by the reflexivity of \mathfrak{F} . If $\dim H_1 < \infty$, it is easy to prove that $\text{Alg } \phi(\mathfrak{F})$ is hyper-reflexive ([5]). Now suppose that $\dim H_1 = \infty$. Then there exists a bicontinuous linear or conjugate linear bijection $S: H_1 \rightarrow H_2$ such that $\phi = \phi_S$ i.e., $\phi(M) = SM$ for every $M \in \xi(H_1)$ (see [1]).

We may suppose that S is conjugate linear. For any $f \in (\text{Alg } \phi(\mathfrak{F}))_\perp$, define $g(A) = \overline{f(SAS^{-1})}$, then $g \in (\text{Alg } \mathfrak{F})_\perp$ since $\text{Alg } \phi(\mathfrak{F}) = S(\text{Alg } \mathfrak{F})S^{-1}$.

By theorem A, there exist $x_n, y_n \in H_1$ such that $(x_n \otimes y_n) \in (\text{Alg } \mathfrak{F})_\perp$,

$$g = \sum_{n=1}^{\infty} (x_n \otimes y_n)$$

$$\sum_{n=1}^{\infty} \|x_n\| \|y_n\| < \infty.$$

For any $T \in L(H_2)$

$$f(T) = \overline{g(S^{-1}TS)} = \sum_{n=1}^{\infty} \overline{(S^{-1}TSx_n, y_n)}$$

$$\begin{aligned}
 &= \sum_{n=1}^{\infty} \overline{((S^{-1})^* y_n, TSx_n)} \\
 &= \sum_{n=1}^{\infty} (TSx_n, (S^*)^{-1} y_n).
 \end{aligned}$$

Let $u_n = Sx_n, v_n = (S^*)^{-1} y_n$, then

$$f = \sum_{n=1}^{\infty} (u_n, \otimes v_n)$$

and $(u_n \otimes v_n) \in (\text{Alg } \phi(\mathfrak{F}))_{\perp}$, $\sum_{n=1}^{\infty} \|u_n\| \|v_n\| < \infty$. And therefore $\text{Alg } \phi(\mathfrak{F})$ is hyperreflexive by

Theorem A. The proof is complete.

A PROOF OF THEOREM 2. Since $0 \bar{\in} \pi(A), R(A)$, the range of A , is a closed subspace of H . Let $H_1 = H(A)$, then ϕ_A defines a lattice isomorphism from $\xi(H)$ onto $\xi(H_1)$. From Lemma 1 we have that $\text{Alg}_H \phi_A(\mathfrak{F}) = \{T \in L(H_1) : \phi_A(\mathfrak{F}) \subseteq \text{Lat}(T)\}$ is hyper-reflexive. By the definition of hyper-reflexivity,¹ there exists $K > 0$ such that for any $T \in L(H_1)$

$$\begin{aligned}
 &\text{dist}(T, \text{Alg}_{H_1} \phi_A(\mathfrak{F})) \\
 &\leq K \sup \{ \|(I_{H_1} - \bar{P} \phi_A(M)) T \bar{P} \phi_A(M)\| : M \in \mathfrak{F} \}
 \end{aligned}$$

where $P_{\phi_A(M)}$ denotes the orthogonal projection from H_1 onto $\phi_A(M)$. For any $T \in \text{Alg}_{H_1}(\phi_A(\mathfrak{F}))$ and $S \in L(H)$, we define an operator $\tilde{T} \in L(H)$ by formula

$$\tilde{T}(x \otimes y) = Tx + Sy, \quad x \otimes y \in H = H_1 \otimes H_1^{\perp}.$$

Then $\tilde{T} \in \text{Alg}(\phi_A(\mathfrak{F}) \cup \{H\})$ since $T \in \text{Alg}_{H_1} \phi_A(\mathfrak{F})$. Denote by E the orthogonal projection from H onto H_1 , then

$$\begin{aligned}
 &\text{dist}(S, \text{Alg}(\phi_A(\mathfrak{F}) \cup \{H\})) \\
 &\leq \|S - T\| \\
 &\leq \|ES|_{H_1} - T\| + \|E^{\perp}SE\| \\
 &\leq \|ESE - T\| + \sup \{ \|P_{\phi_A(M)}^{\perp} SP_{\phi_A(M)}\| : M \in \mathfrak{F} \}
 \end{aligned}$$

and so

$$\begin{aligned}
 &\text{dist}(S, \text{Alg}(\phi_A(\mathfrak{F}) \cup \{H\})) \\
 &\leq \text{dist}(ESE, \text{Alg}_{H_1} \phi_A(\mathfrak{F})) + \sup \{ \|P_{\phi_A(M)}^{\perp} SP_{\phi_A(M)}\| : M \in \mathfrak{F} \} \\
 &\leq (K + 1) \sup \{ \|P_N^{\perp} SP_N\| : N \in \phi_A(\mathfrak{F}) \cup \{H\} \}
 \end{aligned}$$

which implies the hyper-reflexivity of $\text{Alg}_A(\mathfrak{F}) \cup \{H\}$. Together with the reflexivity of $\phi_A(\mathfrak{F}) \cup \{H\}$

(see [3]), we obtain $\phi_A(\mathcal{F}) \cup \{H\}$ is hyper-reflexivity.

REFERENCES

1. FILLMORE, P.A. and LONGSTAFF, W.E., On isomorphism of lattices of closed subspaces, Cand. J. Math., **36** (1984), 820-829.
2. HARRISON, K.J. and LONGSTAFF, W.E., Automorphic images of commutative subspace lattices, Trans. Amer. Math. Soc., **296** (1986), 217-228.
3. LONGSTAFF, W.E., Subspaces maps of operators on Hilbert spaces, Proc. Amer. Math. Soc., **84** (1982), 195-201.
4. ARVESON, W., Ten lectures on operator algebras. CBMS Regional Conference Series, No. 55, A.M.S., Providence, 1984.
5. KRAUS, J. and LARSON, D.R., Reflexivity and distance formulae, Proc. London Math. Soc., **53** (1986), 340-356.