

TOTALLY REAL SURFACES IN CP^2 WITH PARALLEL MEAN CURVATURE VECTOR

M. A. AL-GWAIZ and SHARIEF DESHMUKH

Department of Mathematics
King Saud University
P.O. Box 2455
Riyadh 11451
Saudi Arabia

(Received October 26, 1990 and in revised form May 10, 1991)

Abstract. It has been shown that a totally real surface in CP^2 with parallel mean curvature vector and constant Gaussian curvature is either flat or totally geodesic.

Key Words and Phrases: Riemannian connection, Gaussian curvature and real surfaces.

1991 Mathematics Subject Classification Codes: 53B21, 53C20

1. INTRODUCTION.

Let J be the almost complex structure on CP^2 and g be the Hermitian metric on CP^2 of constant holomorphic sectional curvature 4. If $\bar{\nabla}$ is the Riemannian connection with respect to g and \bar{R} is the curvature tensor of $\bar{\nabla}$, then

$$(\bar{\nabla}_X J)(Y) = 0, \quad (1.1)$$

$$\bar{R}(X, Y)Z = g(Y, Z)X - g(X, Z)Y + g(JY, Z)JX - g(JX, Z)JY + 2g(X, JY)JZ, \quad (1.2)$$

where X, Y, Z are vector fields on CP^2 .

Let M be a 2-dimensional totally real submanifold of CP^2 and ν be the normal bundle of M . If $\chi(M)$ is the lie-algebra of vector fields on M , then for each $X \in \chi(M)$, $JX \in \nu$. The Riemannian connection $\bar{\nabla}$ induces the Riemannian connection ∇ on M and the connection ∇^\perp in the normal bundle ν . We then have the following Gauss and Weingarten formulae

$$\bar{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad \nabla_X N = -A_N X + \nabla_X^\perp N, \quad X, Y \in \chi(M), \quad N \in \nu, \quad (1.3)$$

where $h(X, Y)$ and $A_N X$ are the second fundamental forms and are related by $g(h(X, Y), N) = g(A_N X, Y)$. The mean curvature vector H of M is given by

$$H = (1/2) \sum h(e_i, e_i),$$

where $\{e_1, e_2\}$ is a local orthonormal frame on M . If $H = 0$, then M is said to be a minimal submanifold of CP^2 . It is known that if M is a minimal totally real surface of constant Gaussian curvature in CP^2 , then either M is flat or totally geodesic (cf. [2]). The mean curvature vector H is said to be parallel if $\nabla_X^\perp H = 0$, $X \in \chi(M)$. In this paper we consider the totally real surfaces of constant Gaussian curvature with parallel mean curvature vector in CP^2 .

The Gaussian curvature K of M is given by

$$K = 1 + g(h(X,X), 2h(Y,Y)) - g(h(X,Y), h(X,Y)), \tag{1.4}$$

where $\{X, Y\}$ is an orthonormal frame on M . The Codazzi equation gives

$$(\bar{\nabla}_X h)(Y, Z) = (\bar{\nabla}_Y h)(X, Z), \quad X, Y, Z \in \chi(M). \tag{1.5}$$

For a totally real surface M , using (1.1) and (1.3), we get

$$h(X, Y) = JA_{JY}X, \quad \nabla_X^\perp JY = J\nabla_X Y, \quad X, Y \in \chi(M). \tag{1.6}$$

Using (1.6) and the symmetry of $h(X, Y)$, we have

$$g(h(Y, Z), JX) = g(h(X, Y), JZ) = g(h(X, Z), JY), \quad X, Y, Z \in \chi(M). \tag{1.7}$$

2. MAIN RESULTS

THEOREM 2.1. Let M be a connected totally real surface in CP^2 of constant Gaussian curvature c with parallel mean curvature vector. Then either M is flat or totally geodesic.

PROOF. Let $UM = \{X \in TM : \|X\| = 1\}$ be the unit tangent bundle of M . Define the function $f: UM \rightarrow R$ by $F(X) = g(h(X, X), JX)$, which is clearly a smooth function. First suppose that f is constant. Then $f(-X) = -f(X)$ gives $f(X) = 0$ and therefore $g(h(X, X), JX) = 0, X \in UM$. Now consider a local orthonormal frame $\{X, Y\}$ on M . Then we have $g(h(X, X), JX) = 0, g(h(Y, Y), JY) = 0,$

$$g\left(h\left(\frac{X+Y}{\sqrt{2}}, \frac{X+Y}{\sqrt{2}}\right), J\left(\frac{X+Y}{\sqrt{2}}\right)\right) = 0, \quad g\left(h\left(\frac{X-Y}{\sqrt{2}}, \frac{X-Y}{\sqrt{2}}\right), J\left(\frac{X-Y}{\sqrt{2}}\right)\right) = 0$$

These equations, in view of (1.7), imply that $g(h(X, X), JY) = 0, g(h(Y, Y), JX) = 0, g(h(X, Y), JX) = 0,$ and $g(h(X, Y), JY) = 0$. Since $\{JX, JY\}$ is a local orthonormal frame in the normal bundle ν , we conclude that $h(X, X) = 0, h(X, Y) = 0$ and $h(Y, Y) = 0$, which means that M is totally geodesic.

We therefore assume that f is not a constant. Since the unit tangent bundle UM is compact, f attains a maximum at some $e_1 \in UM$. It is known that $g(h(e_1, e_1), JY) = 0$ for any vector in TM which is orthogonal to e_1 (cf. [1]). Choose e_2 such that $\{e_1, e_2\}$ is an orthonormal frame on M . Then we can set

$$h(e_1, e_1) = \alpha J e_1, \quad h(e_2, e_2) = \beta J e_1 + \gamma J e_2 \quad \text{and} \quad h(e_1, e_2) = \beta J e_2, \tag{2.1}$$

where α, β and γ are smooth functions. Using the structure equations of M we have locally

$$\nabla_{e_1} e_1 = a e_2, \quad \nabla_{e_2} e_2 = b e_1, \quad \nabla_{e_1} e_2 = -a e_1, \quad \nabla_{e_2} e_1 = -b e_2, \tag{2.2}$$

where a, b are smooth functions. Inserting different combinations of the frame vectors e_1, e_2 in (1.5) and using (2.1) and (2.2) we get, upon equating components,

$$e_1 \cdot \beta = a\gamma + 2b\beta - b\alpha, \quad e_2 \cdot \alpha = a(\alpha - 2\beta), \quad e_2 \cdot \beta - e_1 \cdot \gamma = 3a\beta - b\gamma. \tag{2.3}$$

Since the mean curvature vector $H = (1/2)(h(e_1, e_1) + h(e_2, e_2))$ is parallel, we have

$$\nabla_{e_1}^\perp (h(e_1, e_1) + h(e_2, e_2)) = 0 \quad \text{and} \quad \nabla_{e_2}^\perp (h(e_1, e_1) + h(e_2, e_2)) = 0.$$

Using (1.6), (2.1) and (2.2) in the above equations we conclude, upon equating components, that

$$e_1 \cdot (\alpha + \beta) = a\gamma, \quad e_1 \cdot \gamma = -a(\alpha + \beta) \tag{2.4}$$

$$e_2 \cdot (\alpha + \beta) = -b\gamma, \quad e_2 \cdot \gamma = b(\alpha + \beta). \tag{2.5}$$

From (2.3), (2.4) and (2.5), we have

$$\begin{aligned} e_1 \cdot \alpha &= b(\alpha - 2\beta), & e_1 \cdot \beta &= a\gamma + 2b\beta - b\alpha, & e_1 \cdot \gamma &= -a(\alpha + \beta), \\ e_2 \cdot \alpha &= a(\alpha - 2\beta), & e_2 \cdot \beta &= -b\gamma + 2a\beta - a\alpha, & e_2 \cdot \gamma &= b(\alpha + \beta). \end{aligned} \tag{2.6}$$

In view of (2.1) and (1.4), the Gaussian curvature c is given by $c = 1 + \alpha\beta - \beta^2$. If we operate on this equation by e_1 and e_2 with c constant, and use (2.6), we obtain

$$(\alpha - 2\beta)(a\gamma + b(3\beta - \alpha)) = 0 \text{ and } (\alpha - 2\beta)(-b\gamma + a(3\beta - \alpha)) = 0. \tag{2.7}$$

We have two cases:

Case (i). Suppose $\alpha \neq 2\beta$, then the two equations in (2.7) give $(a^2 + b^2)\gamma = 0$ and $(a^2 + b^2)(3\beta - \alpha) = 0$. If $a^2 + b^2 = 0$, then from (2.2) it follows that M is flat (as c is constant). If $a^2 + b^2 \neq 0$, then we have $\gamma = 0$ and $3\beta - \alpha = 0$. Since a and b cannot both be zero and $\gamma = 0$ it follows from equations (2.4) and (2.5) that $\alpha + \beta = 0$. Thus we have $\gamma = 0$ and $\alpha + \beta = 0$, which implies that $H = 0$, that is, M is minimal.

Case (ii). Suppose $\alpha = 2\beta$. Then from (2.6) we get that α is constant, and consequently β is also constant. With $\alpha = 2\beta$ and β constant equations (2.6) give $a\gamma = 0$ and $b\gamma = 0$. Thus either $a = b = 0$ or $\gamma = 0$, which results in either M being flat or $\gamma = 0$. If M is not flat, that is, not both a and b are zero, and $\gamma = 0$, then from (2.4) and (2.5) we get $\alpha + \beta = 0$. This shows that $H = 0$. Hence either M is flat or minimal. But since a minimal totally real surface is constant curvature in CP^2 is either flat or totally geodesic [2], the theorem is proved.

In the following we first prove that in any submanifold of a Riemannian manifold if the second fundamental form is parallel, then the mean curvature vector is parallel. Though this is a simple observation, it does not seem to appear in the literature and is worth mentioning. As a corollary then we obtain the same result as in Section 2 for the totally real surfaces of CP^2 with parallel second fundamental form.

THEOREM 2.2. Let M be a submanifold of a Riemannian manifold \overline{M} with parallel second fundamental form. Then the mean curvature vector of M is parallel.

PROOF. Suppose $\dim M = n$. Then for a local orthonormal frame $\{e_1, e_2, \dots, e_n\}$ of M , the mean curvature H is given by

$$H = (1/n) \sum_{i=1}^n h(e_i, e_i).$$

Since the second fundamental form is parallel we have

$$(\overline{\nabla}_X h)(Y, Z) = \nabla_X^\perp h(Y, Z) - h(\nabla_X Y, Z) - h(Y, \nabla_X Z) = 0 \text{ for } X, Y, Z \in \chi(M).$$

Thus for each frame vector e_i we can write

$$\nabla_X^\perp h(e_i, e_i) = 2h(\nabla_X e_i, e_i).$$

Adding these equations we get

$$n \nabla_X^\perp H = 2 \sum_{i=1}^n h(\nabla_X e_i, e_i).$$

Let ω_j^i be the connections forms on M . Then we have

$$\nabla_X e_i = \sum_{j=1}^n \omega_j^i(x) e_j.$$

Substituting this into the above equation we get

$$n \nabla_X^\perp H = 2 \sum_{i,j=1}^n \omega_i^j(X) h(e_i, e_j).$$

Since $\omega_i^j(X) = -\omega_j^i(X)$ and $h(e_i, e_j) = h(e_j, e_i)$, we conclude that $\nabla_X^\perp H = 0$, $X \in \chi(M)$.

As a direct consequence of this theorem and the theorem in the previous section we have

COROLLARY 2.1. Let M be a connected totally real surface in CP^2 with parallel second fundamental form and constant Gaussian curvature. Then M is either flat or totally geodesic.

ACKNOWLEDGEMENT. This work has been supported by grant No. (Math/1409/05) of the Research Center, College of Science, King Saud University, Riyadh, Saudi Arabia.

REFERENCES

- [1] Ejiri, N., Totally Real Minimal Immersions of N -dimensional Real Space Forms Into N -dimensional Complex Space Forms, Proc. Amer. Math. Soc. **84** (1982), 243-246.
- [2] Houh, C. S., Some Totally Real Minimal Surfaces in CP^2 , Proc. Amer. Math. Soc. **40**(1973), 240-244.