

RESEARCH NOTES COMPLETE LIFT OF A STRUCTURE SATISFYING

$$F^K - (-)^{K+1} F = 0$$

LOVEJOY S. DAS

Department of Mathematics
Kent State University
Tuscarawas Campus
New Philadelphia, Ohio 44663

(Received August 30, 1991 and in revised form January 12, 1992)

ABSTRACT. The idea of f -structure manifold on a differentiable manifold was initiated and developed by Yano [1], Ishihara and Yano [2], Goldberg [3] and among others. The horizontal and complete lifts from a differentiable manifold M^n of class C^∞ to its cotangent bundles have been studied by Yano and Patterson [4,5]. Yano and Ishihara [6] have studied lifts of an f -structure in the tangent and cotangent bundles. The purpose of this paper is to obtain integrability conditions of a structure satisfying $F^K - (-)^{K+1}F = 0$ and $F^W - (-)^{W+1}F \neq 0$ for $1 < W < K$, in the tangent bundle.

KEY WORDS AND PHRASES. Tangent bundle, Complete lift, F -structure, Integrability, Distributions.

1991 AMS SUBJECT CLASSIFICATION CODE. 53C15.

1. INTRODUCTION.

Let F be a nonzero tensor field of the type (1,1) and of class C^∞ on an n dimensional manifold M^n such that [7]

$$F^K - (-)^{K+1} F = 0 \quad \text{and} \quad F^W - (-)^{W+1} F \neq 0 \quad \text{for} \quad 1 < W < K, \quad (1.1)$$

where K is a fixed positive integer greater than 2. The degree of the manifold being $K(K \geq 3)$. Such a structure on M^n has been called $F(K, -(-)^{K+1})$ - structure of rank r , where the rank $(F) = r$ and is constant on M^n . The case when K is odd and $K(\geq 3)$ has been considered in this paper.

Let the operators on M^n be defined as follows [7]

$$l = (-)^{K+1} F^{K+1} \quad \text{and} \quad m = I - (-)^{K+1} F^{K+1}, \quad (1.2)$$

where I denotes the identity operator on M^n . We will state the following two theorems. [7]

THEOREM (1.1). Let M^n be an $F(K, -(-)^{K+1})$ manifold then,

$$l + m = I, \quad l^2 = l \quad \text{and} \quad m^2 = m \quad (1.3)$$

For F satisfying (1.1), there exist complementary distributions L and M , corresponding to the projection operators l and m respectively. If the rank of F is constant and is equal to $r = r(F)$ then $\dim L = r$ and $\dim M = (n-r)$.

THEOREM (1.2). We have

$$a) \quad F1 = 1F = F \quad \text{and} \quad Fm = mF = 0 \quad (1.4)a$$

$$b) \quad F^{K-1}1 = 1 \quad \text{and} \quad F^{K-1}m = 0 \quad (1.4)b$$

Then $F^{\frac{K-1}{2}}$ acts on L as an almost product structure and on M as a null operator.

2. COMPLETE LIFT ON $F(K, (-)^{K+1})$ -STRUCTURE IN TANGENT BUNDLE.

Let M be an n -dimensional differentiable manifold of class C^∞ and $T_p(M^n)$ the tangent space at a point p of M_n and

$$T(M^n) = \bigcup_{p \in M^n} T_p(M^n) \text{ is the tangent bundle over the manifold } M^n.$$

Let us denote by $T_s^r(M^n)$, the set of all tensor fields of class C^∞ and of type (r, s) in M^n and $T(M^n)$ be the tangent bundle over M^n . The complete lift F^C of an element of $T_1^1(M^n)$ with local components F_i^h has components of the form [5]

$$F^C : \begin{pmatrix} F_i^h & 0 \\ \delta_i^h & F_i^h \end{pmatrix}$$

Now we obtain the following results on the complete lift of F satisfying (1.1).

THEOREM (2.1). For $F \in T_1^1(M^n)$, the complete lift F^C of F is an $F(K, (-)^{K+1})$ -structure iff it is for F also. Then F is of rank r iff F^C is of rank $2r$.

PROOF. Let $F, G \in T_1^1(M^n)$. Then we have [5]

$$(FG)^C = F^C G^C \quad (2.2)$$

Replacing G by F in (2.2) we obtain

$$\begin{aligned} (FF)^C &= F^C F^C \\ \text{or, } (F^2)^C &= (F^C)^2 \end{aligned} \quad (2.3)$$

Now putting $G = F^{K-1}$ in (2.2) since G is $(1,1)$ tensor field therefore F^{K-1} is also $(1,1)$ so we obtain $(FF^{K-1})^C = F^C (F^{K-1})^C$ which in view of (2.3) becomes

$$(F^K)^C = (F^C)^K \quad (2.4)$$

Taking complete lift on both sides of equation (1.1) we get

$$(F^K)^C - ((-)^{K+1}F)^C = 0$$

which in consequence of equation (2.4) gives

$$(F^C)^K - (-)^{K+1} F^C = 0 \quad (2.5)$$

Thus equation (1.1) and (2.5) are equivalent. The second part of the theorem follows in view of equation (2.1).

Let F satisfying (1.1) be an F -structure of rank r in M^n . Then the complete lifts 1^C of 1 and m^C of m are complementary projection tensors in $T(M^n)$. Thus there exist in $T(M^n)$ two complementary distributions L^C and M^C determined by 1^C and m^C respectively.

3. INTEGRABILITY CONDITIONS OF $F(K, (-)^{K+1})$ STRUCTURE IN TANGENT BUNDLE.

Let $F \in T_1^{-1}(M^n)$, then the Nijenhuis tensor N_F of F satisfying (1.1) is a tensor field of the type (1,2) given by [6]

$$N_F(X, Y) = [FX, FY] - F[FX, Y] - F[X, FY] + F^2[X, Y]. \tag{3.1a}$$

Let N^C be the Nijenhuis tensor of F^C in $T(M^n)$ of F in M^n , then we have

$$N^C(X^C, Y^C) = [F^C X^C, F^C Y^C] - F^C[F^C X^C, Y^C] - F^C[X^C, F^C Y^C] + (F^2)^C[X^C, Y^C] \tag{3.1b}$$

For any $X, Y \in T_0^{-1}(M^n)$ and $F \in T_1^{-1}(M^n)$ we have [5]

$$[X^C, Y^C] = [X, Y]^C \text{ and } (X+Y)^C = X^C + Y^C \tag{3.2a}$$

$$F^C X^C = (FX)^C \tag{3.2b}$$

From (1.4)a and (3.2)b we have

$$F^C m^C = (Fm)^C = 0 \tag{3.3}$$

THEOREM (3.1). The following identities hold

$$N^C(m^C X^C, m^C Y^C) = (F^C)^2 [m^C X^C, m^C Y^C], \tag{3.4}$$

$$m^C N^C(X^C, Y^C) = m^C [F^C X^C, F^C Y^C] \tag{3.5}$$

$$m^C N^C(1^C X^C, 1^C Y^C) = m^C [F^C X^C, F^C Y^C] \tag{3.6}$$

$$m^C N^C((F^C)^{K-2} X^C, (F^C)^{K-2} Y^C) = m^C [1^C X^C, 1^C Y^C] \tag{3.7}$$

PROOF. The proofs of (3.4) to (3.7) follow in view of equations (1.4), (3.1)a, and (3.3)

THEOREM (3.2). For any $X, Y \in T_0^{-1}(M^n)$, the following conditions are equivalent.

$$m^C N^C(X^C, Y^C) = 0, \tag{i}$$

$$m^C N^C(1^C X^C, 1^C Y^C) = 0, \tag{ii}$$

$$m^C N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0 \tag{iii}$$

PROOF. In consequence of equations (3.1)b and (1.4)a,b it can be easily proved that

$$N^C(1^C X^C, 1^C Y^C) = 0 \text{ iff } N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0 \text{ for all } X \text{ and } Y \in T_0^{-1}(M^n)$$

Now due to the fact that equations (3.5) and (3.6) are equal, the conditions (i), (ii) and (iii) are equivalent to each other.

THEOREM (3.3). The complete lift of M^C of the distribution M in $T(M^n)$ is integrable iff M is integrable in M^n .

PROOF. It is known that the distribution M is integrable in M^n iff [2]

$$l[mX, mY] = 0 \tag{3.8}$$

for any $X, Y \in T_0^{-1}(M^n)$.

Taking complete lift on both sides of equations (3.8) we get

$$1^C [m^C X^C, m^C Y^C] = 0 \tag{3.9}$$

where $1^C = (I - m)^C = I - m^C$, is the projection tensor complementary to m^C . Thus the

conditions (3.8) and (3.9) are equivalent.

THEOREM (3.4). For any $X, Y \in T_0^1(M^n)$, let the distribution M be integrable in M^n iff $N(mX, mY) = 0$.

Then the distribution m^C is integrable in $T(M^n)$ iff $1^C N^C(m^C X^C, m^C Y^C) = 0$ or equivalently, $N^C(m^C X^C, m^C Y^C) = 0$.

PROOF. By virtue of condition (3.4) we have

$$N^C(m^C X^C, m^C Y^C) = (F^C)^2 [m^C X^C, m^C Y^C].$$

Multiplying throughout by 1^C we get

$$1^C N^C(m^C X^C, m^C Y^C) = (F^C)^2 1^C [m^C X^C, m^C Y^C].$$

which in view of equation (3.9) becomes

$$1^C N^C(m^C X^C, m^C Y^C) = 0 \quad (3.10)$$

Making use of equation (3.3), we get

$$m^C N^C(m^C X^C, m^C Y^C) = 0 \quad (3.11)$$

Adding (3.10) and (3.11) we obtain

$$(1^C + m^C) N^C(m^C X^C, m^C Y^C) = 0$$

Since $1^C + m^C = I^C = I$ we have $N^C(m^C X^C, m^C Y^C) = 0$.

THEOREM (3.5). For any $X, Y \in T_0^1(M^n)$ let the distribution L be integrable in M^n that is $mN(X, Y) = 0$ then the distribution L^C is integrable in $T(M^n)$ iff any one of the conditions of theorem (3.2) is satisfied.

PROOF. The distribution L is integrable in M^n iff [2] holds i.e. $m[1X, 1Y] = 0$. Thus the distribution L^C is integrable in $T(M^n)$ iff $m^C[1^C X^C, 1^C Y^C] = 0$.

On making use of equation (3.7), the theorem follows.

We now define the following:

- (i) distribution L is integrable,
- (ii) an arbitrary vector field Z tangent to an integral manifold of L ,
- (iii) the operator F , such that $FZ = FZ$.

Hence by virtue of theorem (1.2) the induced structure F^* is an almost product structure on each integral manifold of L and F^* makes tangent spaces invariant of every integral manifold of L . Let us denote the vector valued 2-form $N^*(Z, W)$, the Nijenhuis tensor corresponding to the Nijenhuis tensor of the almost product structure induced form $F(K, -(-)^{K+1})$ structure, on each integral manifold of L and for any two $Z, W \in T_0^1(M^n)$ tangent to an integral manifold of L , then we have

$$N^*(Z, W) = [F^*Z, W] - F^*[F^*Z, W] - F^*[Z, F^*W] + F^{*2}[Z, W], \quad (3.12)$$

which in view of (3.1)b and (3.12) yields

$$N^C[1^C X^C, 1^C Y^C] = N^C(1^C X^C, 1^C Y^C) \quad (3.13)$$

DEFINITION (3.1). We say that $F(K, -(-)^{K+1})$ - structure is partially integrable if the distribution L is integrable and the almost product structure F induced from F^* on each integral manifold of L is also integrable.

THEOREM (3.6). For any $X, Y \in T_0^{-1}(M^n)$ let the $F(K, -(-)^{K+1})$ - structure be partially integrable in M^n i.e. $N(1X, 1Y) = 0$. Then the necessary and sufficient condition for $F(K, -(-)^{K+1})$ - structure to be partially integrable in $T(M^n)$ is that $N^C(1^C X^C, 1^C Y^C) = 0$ or equivalently $N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0$.

PROOF. In view of equation (1.4) and equation (3.1)b we can prove easily that $N^C(1^C X^C, 1^C Y^C) = 0$ iff $N^C((F^{K-2})^C X^C, (F^{K-2})^C Y^C) = 0$, for any $X, Y \in T_0^{-1}(M^n)$.

Now in view of equation (3.13) and theorem (3.5), the result follows immediately.

When both distributions L and M are integrable we can choose a local coordinate system such that all L and M are respectively represented by putting $(n - r)$ local coordinates constant and r -coordinates constant. We call such a coordinate system an adapted coordinate system. It can be supposed that in an adapted coordinate system the projection operators l and m have the components of the form

$$l = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}, \quad m = \begin{pmatrix} 0 & 0 \\ 0 & I_{n-r} \end{pmatrix}$$

respectively where I_r denotes the unit matrix of order 'r' and I_{n-r} is of order

$(n - r)$.

Since F satisfies equation (1.4)a, the tensor has components of the form

$$F = \begin{pmatrix} F_r & 0 \\ 0 & 0 \end{pmatrix}$$

in an adapted coordinate system where F_r denotes $r \times r$ square matrix.

DEFINITION (3.2). We say that on $F(K, -(-)^{K+1})$ structure is integrable if

- (i) The structure $F(K, -(-)^{K+1})$ is partially integrable.
- (ii) The distribution M is integrable i.e. $N(mX, mY) = 0$.
- (iii) The components of the $F(K, -(-)^{K+1})$ structure are independent of the coordinates which are constant along the integral manifold of L in an adapted coordinate system.

THEOREM (3.7). For any $X, Y \in T_0^{-1}(M^n)$ let $F(K, -(-)^{K+1})$ - structure be integrable in M^n iff $N(X, Y) = 0$. Then the $F(K, -(-)^{K+1})$ - structure is integrable in $T(M^n)$ iff $N^C(X^C, Y^C) = 0$.

PROOF. In view of equations (3.1)a and (3.1)b we get

$$N^C(X^C, Y^C) = (N(X, Y))^C.$$

Since $F(K, -(-)^{K+1})$ is integrable in M^n thus the theorem follows.

REFERENCES

1. YANO, K., On a structure defined by a tensor field f of type (1,1) satisfying $f^3 + f = 0$, Tensor Vol. 14 (1963), pp. 99-109.
2. ISHIHARA, S. and YANO, K., On integrability conditions of a structure f satisfying $f^3 + f = 0$, Quarterly J. Math. Vol. 15 (1964), pp. 217-222.
3. GOLDBERG, S.I. and YANO, K., Globally framed f -manifolds, Illinois J. Math. Vol 15 (1971), pp. 456-476.

4. YANO, K. and PATTERSON, E. M., Vertical and Complete lifts from a manifold to its cotangent bundles, Journal Math. Soc. Japan. Vol. 19 (1967), pp. 91-113.
5. YANO, K. and PATTERSON, E. M., Horizontal lifts from a manifold to its cotangent bundles, Journal Math. Soc. Japan. Vol. 19 (1967), pp. 185-198.
6. YANO, K. and ISHIHARA, S., Tangent and Cotangent Bundles, Marcel Dekker, Inc. New York, (1973).
7. DAS, LOVEJOY S., On integrability conditions of $F(K, -(-)^{K+1})$ - structure on a differentiable manifold, Revista Mathematica. Argentina. Vol. 26 (1977).
8. DAS, LOVEJOY S. and UPADHYAY, M. D., On F - structure manifold, Kyung Pook Mathematical Journal. Korea. Vol. 18 (1978), pp. 277-283.
9. DAS, LOVEJOY S., On differentiable manifold with $F(K, -(-)^{K+1})$ - structure of rank 'r', Revista Mathematica. Argentina. Vol. 27 (1978), pp. 277-283.
10. DAS, LOVEJOY S., Prolongations of F - structure to the tangent bundle of order 2. International J. Math & Math Sci., (to appear).
11. HELGASON, S., Differential Geometry. Lie Groups and Symmetric Spaces, Academic Press, New York, (1970).

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru