

**AROUND THE FURUTA INEQUALITY -
THE OPERATOR INEQUALITIES $(AB^2A)^{3/4} \leq ABA \leq A^3$**

DERMING WANG

*Department of Mathematics
California State University, Long Beach
Long Beach, California 90840*

(Received July 21, 1993 and in revised form November 5, 1993)

Abstract: For positive operators A and B with A invertible it is shown that $(AB^2A)^{1/2} \leq A^2$ implies $(AB^2A)^{3/4} \leq ABA$. The inequalities in the title for $0 \leq B \leq A$ are then derived as a consequence.

KEY WORDS AND PHRASES. *Positive operator, operator inequality, the Furuta inequality*
1991 MATHEMATICS SUBJECT CLASSIFICATION CODE. 47B15

1. INTRODUCTION.

In this paper operator inequalities centered around the celebrated Furuta inequality are considered. As motivation, we begin with a brief account of the origin of the inequalities in the title.

We consider (bounded, linear) operators acting on a Hilbert space. For an operator A , we write $A \geq 0$ (or $0 \leq A$) if A is a positive operator. For positive operators A and B , we write $A \geq B$ (or $B \leq A$) if $A - B \geq 0$.

It is well-known that $0 \leq B \leq A$ implies that $B^r \leq A^r$ for every real number r with $0 \leq r \leq 1$. Thus, $0 \leq B \leq A$ implies $B^{1/2} \leq A^{1/2}$. But, in general, $0 \leq B \leq A$ does not necessarily imply that $B^2 \leq A^2$. In [1], the following conjecture was raised:

$$\text{If } 0 \leq B \leq A, \text{ then } (AB^2A)^{1/2} \leq A^2. \tag{1}$$

This conjecture was answered affirmatively by Furuta [2]. Indeed, Furuta proved a more general inequality that contains inequality (1) as a special case:

THE FURUTA INEQUALITY. For $p, r \geq 0$ and $q \geq 1$ with $p + 2r \leq (1 + 2r)q$,

$$0 \leq B \leq A \text{ implies } (A^r B^p A^r)^{1/q} \leq A^{(p+2r)/q}.$$

Setting $p = q = 2$ and $r = 1$, The Furuta inequality becomes (1). Furuta also observed that setting $p = 2, r = 1$ and $q = 4/3$, a stronger inequality resulted:

$$\text{If } 0 \leq B \leq A, \text{ then } (AB^2A)^{3/4} \leq A^3. \tag{2}$$

That (2) implies (1) can readily be seen by taking the $2/3$ -power of both sides of (2).

After the appearance of Furuta's original paper [2], Kamei [3] gave a direct proof of inequalities (1) and (2) using the special notion of *operator means*. More recently, Furuta [4] constructed the operator function $G_r(p) = (A^r B^p A^r)^{(1+2r)/(p+2r)}$ for $A \geq B \geq 0$, $r \geq 0$ and $p \in [1, \infty)$, and showed that $G_r(p)$ is a decreasing function on $[1, \infty)$. In particular, $G_1(2) \leq G_1(1)$. This yielded the following improvement of (2):

$$\text{If } 0 \leq B \leq A, \text{ then } (AB^2A)^{3/4} \leq ABA \leq A^3. \quad (3)$$

The main result of this paper is to show that for positive operators A and B with A invertible, the inequality $(AB^2A)^{3/4} \leq ABA$ is a consequence of the inequality $(AB^2A)^{1/2} \leq A^2$. We also establish inequality (3) as a corollary of our result by giving a new proof of (1) which appears to be simpler than those of [3], [5] and [6]. Our proof is completely elementary. It was inspired by the work of Pedersen and Takesaki [7].

2. THE MAIN RESULT.

THEOREM. *Suppose A and B are positive operators with A invertible. Then*

$$(AB^2A)^{1/2} \leq A^2 \text{ implies } (AB^2A)^{3/4} \leq ABA.$$

PROOF. Let $T = A^{-1}(AB^2A)^{1/2}A^{-1}$. The assumptions imply that $0 \leq T \leq I$, the identity operator. Simple calculation shows that $B^2 = TA^2T$. Now

$$\begin{aligned} [A^{-1}(AB^2A)^{3/4}A^{-1}]^2 &= [A^{-1}(ATA)^{3/2}A^{-1}]^2 \\ &= A^{-1}(ATA)^{1/2}(ATA)A^{-2}(ATA)(ATA)^{1/2}A^{-1} \\ &= A^{-1}(ATA)^{1/2}(AT^2A)(ATA)^{1/2}A^{-1} \\ &\leq A^{-1}(ATA)^{1/2}(ATA)(ATA)^{1/2}A^{-1} \\ &= A^{-1}(ATA)^2A^{-1} = TA^2T = B^2 \end{aligned}$$

Taking square roots, we have $A^{-1}(AB^2A)^{3/4}A^{-1} \leq B$ and hence $(AB^2A)^{3/4} \leq ABA$, and the proof is completed.

COROLLARY 1. *If $0 \leq B \leq A$, then $(AB^2A)^{3/4} \leq ABA \leq A^3$.*

PROOF. Without loss of generality, assume that A is invertible. In view of the theorem, it suffices to establish the inequality $(AB^2A)^{1/2} \leq A^2$. Again we employ the idea of Pedersen and Takesaki. Let $S = A^{-1/2}(A^{1/2}BA^{1/2})^{1/2}A^{-1/2}$. Since $0 \leq B \leq A$, $0 \leq S \leq I$ and $B = SAS$. Thus

$$\begin{aligned} (AB^2A)^{1/2} &= (A(SAS)^2A)^{1/2} = (ASAS^2ASA)^{1/2} \\ &\leq (ASA^2SA)^{1/2} = ASA \leq A^2. \end{aligned}$$

This completes the proof.

The following improvement of (1) is a consequence of Corollary 1.

COROLLARY 2. *If $0 \leq B \leq A$, then $(AB^2A)^{1/2} \leq (ABA)^{2/3} \leq A^2$.*

COROLLARY 3. *Suppose A and B are positive operators with A invertible.*

- (a) *Then, $(AB^2A)^{3/4} = ABA$ if the operator $T = A^{-1}(AB^2A)^{1/2}A^{-1}$ is a projection.*
- (b) *If, in addition B is invertible, then $(AB^2A)^{3/4} = ABA$ if and only if $A = B$.*

PROOF. (a) If T is a projection then $T^2 = T$. In this case, the only " \leq " in the main body of the proof of the theorem becomes " $=$ ".

(b) If $(AB^2A)^{3/4} = ABA$, then the occurrence of " \leq " mentioned in (a) again becomes " $=$ ". If B is also invertible, then T is invertible. Consequently, $T^2 = T = I$ and hence $A = B$.

REFERENCES

1. CHAN, N. N. and KWONG, M. K. Hermitian Matrix Inequalities and a Conjecture, Amer. Math. Monthly, **92** (1985), 533 – 541.
2. FURUTA, T. $A \geq B \geq 0$ Assures $(B^r A^p B^r)^{1/q} \geq B^{(p+2r)/q}$ for $r \geq 0$, $p \geq 0$, $q \geq 1$ with $(1+2r)q \geq p+2r$, Proc. Amer. Math. Soc., **101** (1987), 85 – 88.
3. KAMEI, E. Furuta's Inequality via Operator Mean, Math. Japon., **33** (1988), 737 – 739.
4. FURUTA, T. Two Operator Functions with Monotone Property, Proc. Amer. Math. Soc., **111** (1991), 511 – 516.
5. FURUTA, T. $A \geq B \geq 0$ Ensures $(BA^2B)^{1/2} \geq B^2$, Amer. Math. Monthly, **95** (1988), 933 – 934.
6. WANG, D. The Polar Decomposition and a Matrix Inequality, Amer. Math. Monthly, **96** (1989), 517 – 519.
7. PEDERSEN, G. K. and TAKESAKI, M. The Operator Equation $THT = K$, Proc. Amer. Math. Soc., **36** (1972), 311 – 312.