

RIGID LEFT NOETHERIAN RINGS

O. D. ARTEMOVYCH

Received 17 January 2003

We prove that any rigid left Noetherian ring is either a domain or isomorphic to some ring \mathbb{Z}_{p^n} of integers modulo a prime power p^n .

2000 Mathematics Subject Classification: 16P40, 16W20, 16W25.

Let R be an associative ring. A map $\sigma : R \rightarrow R$ is called a ring endomorphism if $\sigma(x + y) = \sigma(x) + \sigma(y)$ and $\sigma(xy) = \sigma(x)\sigma(y)$ for all elements $a, b \in R$. A ring R is said to be rigid if it has only the trivial ring endomorphisms, that is, identity id_R and zero 0_R . Rigid left Artinian rings were described by Maxson [9] and McLean [11]. Friger [4, 6] has constructed an example of a noncommutative rigid ring R with the additive group R^+ of finite Prüfer rank. A characterization for rigid rings of finite rank was obtained by the author in [1]. Some aspects of a ring rigidity has been studied by Suppa [12, 13], Friger [5], and the author [2].

In this paper, we study rigid left Noetherian rings and prove the following theorem.

THEOREM 1. *Let R be a left Noetherian ring. Then R is a rigid ring if and only if $R \cong \mathbb{Z}_{p^t}$ (p is a prime, $t \in \mathbb{N}$) or it is a rigid domain.*

All rings are assumed to be associative and, as a rule, with an identity element. For a ring R , $N(R)$ will always denote the set of all nil elements of R , $\text{char}(R)$ the characteristic, and $\text{Ann}(I) = \{a \in R \mid aI = Ia = \{0\}\}$ the annihilator of I in R . If R is a left order in Q (or equivalently, Q is the left quotient ring of R), then we will write $Q = Q(R)$. Any unexplained terminology is standard as in [10].

We recall that a ring R is reduced if $r^2 = 0$ implies $r = 0$ for any $r \in R$. Clearly, if R is a rigid reduced ring with an identity element, then either $\text{char}(R) = 0$ or $\text{char}(R) = p$ for some prime p .

LEMMA 2. *Let R be a reduced left Goldie ring. If R is rigid, then it is a domain.*

PROOF. Let R be a reduced rigid left Goldie ring. Assume that R is not a domain. From $bx = 0$ (resp., $xb = 0$), where $b, x \in R$, it holds that $(xb)^2 = 0$ (resp., $(bx)^2 = 0$) and thus a right (resp., left) annihilator of every element b in R coincides with $\text{Ann}(b)$. Moreover, in view of [10, Lemma 2.3.2(i)], $\text{Ann}(a)$ is a maximal left annihilator for some $a \in R$.

Assume that the quotient ring $R/\text{Ann}(a)$ contains elements $\bar{x} = x + \text{Ann}(a) \neq \bar{0}$, $\bar{y} = y + \text{Ann}(a)$ such that

$$\bar{x} \bar{y} = \bar{0} \tag{1}$$

for some $x, y \in R$. Since $y \in \text{Ann}(ax)$ and $\text{Ann}(a) = \text{Ann}(ax)$, we obtain that $\overline{y} = \overline{0}$. This means that $R/\text{Ann}(a)$ is a domain.

By [10, Lemma 2.3.3], $I_a = Ra \oplus \text{Ann}(a)$ is an essential left ideal of R and so by [10, Corollary 3.1.8], $Q(I_a) = Q(R)$. Then the map $\sigma : I_a \rightarrow I_a$ given by $\sigma(ra) = ra$ ($r \in R$) and $\sigma(\text{Ann}(a)) = \{0\}$ is a nontrivial ring endomorphism of I_a . If $\overline{\sigma} : Q(R) \rightarrow Q(R)$ is an extension of σ to $Q(R)$, then

$$\overline{\sigma}(r)a = \overline{\sigma}(ra) = ra \tag{2}$$

for any $r \in R$, in which case,

$$a(\overline{\sigma}(r) - r) = 0 = (\overline{\sigma}(r) - r)a. \tag{3}$$

Since $\overline{\sigma}(r) - r = q^{-1}t$ for some regular element $q \in R$ and some $t \in R$, we see that

$$q(\overline{\sigma}(r) - r) \in \text{Ann}(a). \tag{4}$$

But $q \notin \text{Ann}(a)$ and so $\overline{\sigma}(r) - r \in \text{Ann}(a)$. This means that $\overline{\sigma}(R) \subseteq R$ and R has a nontrivial ring endomorphism, a contradiction. The lemma is proved. □

In the commutative case, we obtain that a commutative reduced rigid Noetherian ring R of finite exponent is isomorphic to some \mathbb{Z}_p .

Indeed, as it is noted above, $\text{char}(R) = p$ for some prime p . A map $\omega : R \rightarrow R$ given by the rule $\omega(x) = x^p$ ($x \in R$) is a ring endomorphism of R and so $x^p = x$ for all elements x of R . Assume that R is not a domain and then it follows that every prime ideal is maximal in R . Hence R is an Artinian ring by Krull-Akizuki theorem [14, Chapter IV, Section 2, Theorem 2] and by the theorem of [11], $R \cong \mathbb{Z}_p$, contrary to our assumption. This means that R is a domain and [9, Theorem 2.5] allows us to state that $R \cong \mathbb{Z}_p$.

REMARK 3. Maxson [9] has proved that a rigid commutative domain of prime characteristic p is isomorphic to \mathbb{Z}_p . Rigid rings of finite rank were studied in [1]. A characterization of rigid commutative domains (in particular, rigid fields) R of characteristic 0 with the additive group R^+ of infinite (Prüfer) rank is not known. As it is noted in [8], from the result of Gaifman [7], it holds that there exist rigid Peano fields of arbitrary infinite cardinality. Moreover, it was proved by Dugas and Göbel [3] that each field can be embedded into a rigid field of arbitrary large cardinality.

REMARK 4. There exist noncommutative rigid Noetherian domains of characteristic 0 (see [4, 6]).

Recall that a map $d : R \rightarrow R$ is called a derivation of R if

$$d(x + y) = d(x) + d(y), \quad d(xy) = d(x)y + xd(y) \tag{5}$$

for all elements $x, y \in R$. A ring having no nonzero derivations is called differentially trivial (see [1]). Obviously, any differentially trivial ring is commutative.

LEMMA 5. *Let R be a left Noetherian ring such that $N(R) \neq \{0\}$. If R is a rigid ring, then it is isomorphic to some \mathbb{Z}_{p^t} .*

PROOF. Suppose that R is a rigid ring such that $N = N(R) \neq \{0\}$. Then $N \subseteq Z(R)$ (see [9, page 96]). Let d be any nonzero derivation of R . If $zd(R) = \{0\}$ for all elements $z \in N$ of the nilpotency indices $i < n - 1$ and $ad(R) \neq \{0\}$ for some element $a \in N$ of the nilpotency index n , then the rule

$$\sigma(r) = r + ad(r), \quad r \in R, \tag{6}$$

determines a nontrivial ring endomorphism σ of R , a contradiction. Hence

$$N(R)d(R) = \{0\} \tag{7}$$

for every derivation d of R .

Let $K_0 = \{a \in N \mid (N \cap \text{Ann}(N^2))a = \{0\}\}$. Then $N \cap \text{Ann}(K_0) = N \cap \text{Ann}(N^2)$. Assume that $\delta : R/K_0 \rightarrow R/K_0$ is a nonzero derivation of R/K_0 and therefore for every $r \in R$, there is an element $r_1 \in R$ such that

$$\delta(r + K_0) = r_1 + K_0. \tag{8}$$

Moreover, $a_1 \notin K_0$ for some $a \in R$. Writing I for the two-sided ideal of R generated by a_1 , we see that $(N \cap \text{Ann}(N^2))(K_0 + I) \neq \{0\}$. Thus there exists an element $m_0 \in N \cap \text{Ann}(N^2)$ such that $m_0a_1 \neq 0$ and so the rule $g(r) = m_0r_1$, with $r \in R$ and r_1 as in (8), determines a nonzero derivation g of R . In view of (7) $g(r)g(t) = 0$, for any elements $r, t \in R$ and a map $\alpha : R \rightarrow R$ given by the rule $\alpha(r) = r + g(r)$, ($r \in R$) is a nontrivial ring endomorphism of R , a contradiction with hypothesis. This gives that R/K_0 is differentially trivial and consequently commutative. Since $K_0 \subseteq N$ and $N \subseteq Z(R)$, R is a Noetherian ring and, as a consequence of [10, Theorem 4.1.9] and [9, Theorem 2.2], R is an Artinian ring. Finally, by the theorem from [11], $R \cong \mathbb{Z}_{p^t}$ for some prime p and integer t . This completes the proof. □

PROOF OF THEOREM 1. It follows immediately from Lemmas 2 and 5. □

COROLLARY 6. *Any rigid simple left Goldie ring R is a field (or equivalently, any noncommutative simple left Goldie ring has a nontrivial automorphism).*

PROOF. Since $N(R) \subseteq Z(R)$, R is a semiprime ring and so according to [10, Proposition 5.1.5] and Lemma 2, it is a domain. If q is any element of $Q(R) \setminus R$ and $A = q^{-1}Rq$, then A is a left order in $Q(R)$. Moreover, $qAq^{-1} = R$ and so A and R are equivalent left orders in $Q(R)$. By [10, Proposition 5.1.2], R is a maximal left order in $Q(R)$ and thus $A \subseteq R$, which implies $R \subseteq Z(Q(R))$, as required. □

REFERENCES

- [1] O. D. Artemovych, *Differentially trivial and rigid rings of finite rank*, Period. Math. Hungar. **36** (1998), no. 1, 1-16.
- [2] ———, *On I -rigid and q -rigid rings*, Ukraïn. Mat. Zh. **50** (1998), no. 7, 989-993 (Russian).
- [3] M. Dugas and R. Göbel, *All infinite groups are Galois groups over any field*, Trans. Amer. Math. Soc. **304** (1987), no. 1, 355-384.
- [4] M. D. Friger, *Rigid torsion-free rings*, Sibirsk. Mat. Zh. **27** (1986), no. 3, 217-219 (Russian).
- [5] ———, *Strongly rigid and I -rigid rings*, Comm. Algebra **22** (1994), no. 5, 1833-1842.
- [6] ———, *Torsion-free rings: some results on automorphisms and endomorphisms*, Second International Conference on Algebra (Barnaul, 1991), Contemp. Math., vol. 184, American Mathematical Society, Rhode Island, 1995, pp. 111-115.
- [7] H. Gaifman, *Models and types of Peano's arithmetic*, Ann. Math. Logic **9** (1976), no. 3, 223-306.
- [8] C. U. Jensen and H. Lenzing, *Model-Theoretic Algebra with Particular Emphasis on Fields, Rings, Modules*, Algebra, Logic and Applications, vol. 2, Gordon and Breach Science Publishers, New York, 1989.
- [9] C. J. Maxson, *Rigid rings*, Proc. Edinburgh Math. Soc. (2) **21** (1979), no. 2, 95-101.
- [10] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian Rings*, Pure and Applied Mathematics, John Wiley & Sons, Chichester, 1987.
- [11] K. R. McLean, *Rigid Artinian rings*, Proc. Edinburgh Math. Soc. (2) **25** (1982), no. 1, 97-99.
- [12] M. A. Suppa, *Sugli anelli I -rigidi [On I -rigid rings]*, Boll. Un. Mat. Ital. D (6) **4** (1985), no. 1, 145-152 (Italian).
- [13] ———, *Sugli anelli q -rigidi*, Riv. Mat. Univ. Parma **12** (1986), 121-125 (Italian).
- [14] O. Zariski and P. Samuel, *Commutative Algebra. Vol. II*, The University Series in Higher Mathematics, D. Van Nostrand C., New Jersey, 1960.

O. D. Artemovych: Institute of Mathematics, Cracow University of Technology, Warszawska 24, Cracow 31-155, Poland

E-mail address: artemo@usk.pk.edu.pl