

## FIXED POINTS OF ROTATIONS OF $n$ -SPHERE

NAGABHUSHANA PRABHU

(Received 18 May 1992)

**ABSTRACT.** We show that every rotation of an even-dimensional sphere must have a fixed point.

**Keywords and phrases.** Fixed point, eigenvalue.

1991 Mathematics Subject Classification. 51F10, 51F25, 51M04, 15A18.

The curious “Hairy Ball Theorem” [1] states that *there are no continuous nonvanishing vector fields tangent to the  $2k$ -dimensional sphere  $S^{2k}$* . Hairy Ball Theorem, however, is false for  $S^{2k-1}$  (easy to verify), which shows that one can geometrically determine the parity of  $n$  in  $S^n$ .

Here is another geometric and simpler asymmetry between spheres of odd and even dimensions:

**THEOREM 1.** *Every rotation of  $S^{2n}$  has at least one fixed point.*

Once again, as an example below illustrates, one can construct rotations of  $S^{2n-1}$  that have no fixed point.

**PROOF.** Rotation in  $\mathbb{R}^k$  is a linear transformation that preserves distance from the origin. Thus, if  $A$  denotes the transformation matrix, then for every  $x \in \mathbb{R}^k$ ,

$$x^T x = (Ax)^T Ax = x^T A^T Ax, \quad (1)$$

which implies that  $A^T A = I$  or  $A^{-1} = A^T$  (i.e.,  $A$  is an orthogonal matrix).  $A^{-1} = A^T$  implies that  $\det(A) = \pm 1$ . But rotation is a continuous transformation and hence one can find a continuous chain of matrices  $M(t)$  such that  $M(0) = I$  and  $M(1) = A$  and each  $M(t)$ ,  $0 \leq t < 1$ , represents a rotation.  $f(t) = \det(M(t))$  is a continuous function of  $t$  with  $f(0) = 1$ . If  $f(1) = -1$ , by intermediate value theorem  $f(t') = 0$  for  $0 < t' < 1$ , which contradicts the assumption that  $M(t')$  represents a rotation and is therefore nonsingular. Hence,  $\det(A) = +1$  (orthogonal matrices with negative determinant represent reflection).  $S^{2n} \subset \mathbb{R}^{2n+1}$ . Hence, if  $A$  represents a rotation in  $\mathbb{R}^{2n+1}$ , then  $A$  is an order  $2n+1$  matrix. The characteristic polynomial  $P(x) = \det(A - xI)$  is hence of degree  $2n+1$ . Complex roots of  $P(x)$  (if any) occur in conjugate pairs. Hence,  $P(x)$  has at least one real root. Further, since the determinant of  $A$  is the product of its eigenvalues, the product of the roots of  $P(x)$  equals  $+1$ . The product of a pair of complex conjugates is always nonnegative and hence  $A$  must have an even number of negative eigenvalues (counting multiplicity). Since  $P(x)$  has  $2n+1$  roots in all (counting multiplicity), it has at least one positive eigenvalue, say  $\lambda$ ; the eigenvector  $\gamma$  of  $\lambda$

