



# A New Kind of Fibonacci-Like Sequence of Composite Numbers

Dan Ismailescu  
Department of Mathematics  
Hofstra University  
103 Hofstra University  
Hempstead, NY 11549  
USA

[dan.p.ismailescu@hofstra.edu](mailto:dan.p.ismailescu@hofstra.edu)

Jaesung Son  
Ridgewood High School  
627 East Ridgewood Ave  
Ridgewood, NJ 07450  
USA

[sonjaesung@gmail.com](mailto:sonjaesung@gmail.com)

## Abstract

An integer sequence  $(x_n)_{n \geq 0}$  is said to be *Fibonacci-like* if it satisfies the binary recurrence relation

$$x_n = x_{n-1} + x_{n-2}, \quad n \geq 2.$$

We construct a new type of Fibonacci-like sequence of composite numbers.

## 1 The problem and previous results

In this paper we consider *Fibonacci-like* sequences, that is, sequences  $(x_n)_{n=0}^{\infty}$  satisfying the binary recurrence relation

$$x_n = x_{n-1} + x_{n-2}, \quad n \geq 2. \tag{1}$$

If  $x_0 = 0$  and  $x_1 = 1$  then  $x_n = F_n$ , the classical Fibonacci sequence. Similarly, when  $x_0 = 2$  and  $x_1 = 1$  then  $x_n = L_n$ , the Lucas sequence.

Graham [3] proved that there exist relatively prime positive integers  $x_0$  and  $x_1$  such that the sequence  $(x_n)_{n=0}^\infty$  defined by the recurrence above contains no prime numbers;  $x_0$  and  $x_1$  have 33 and 34 digits, respectively. Knuth [6] improved on Graham's method and found a 17-digit pair. Soon after, Wilf [9] discovered a smaller 17-digit pair. Nicol [7] refined Knuth's idea and found a 12-digit pair. Finally, Vsemirnov [8] found a smaller pair of 12 and 11 digits:  $x_0 = 106276436867$ ,  $x_1 = 35256392432$ .

Let us describe the common idea used in proving these results. Start by looking for a finite set of quadruples  $(p_i, m_i, r_i, c_i)$ ,  $1 \leq i \leq t$  with the following properties:

- (a) each  $p_i$  is a prime;
  - (b) every  $p_i$  divides  $F_{m_i}$ , the  $m_i$ -th Fibonacci number;
  - (c) every positive integer  $n$  satisfies a congruence  $n \equiv r_i \pmod{m_i}$  for some  $i = 1, 2, \dots, t$ .
- In other words,  $\{(r_i, m_i)\}_{i=1}^t$  is a *covering system of the integers*.

Next, define  $x_0$  and  $x_1$  as follows:

$$x_0 \equiv c_i F_{m_i - r_i} \pmod{p_i} \quad \text{and} \quad x_1 \equiv c_i F_{m_i - r_i + 1} \pmod{p_i} \quad \text{for } i = 1, 2, \dots, t. \quad (2)$$

From the recurrence relation (1) it follows that in general,  $x_n \equiv c_i F_{n + m_i - r_i} \pmod{p_i}$ . The divisibility property  $F_m \mid F_{sm}$  and condition (b) imply that  $p_i \mid x_n$  if  $n \equiv r_i \pmod{m_i}$ .

Since  $x_n$  is an increasing sequence and all primes  $p_i$  are relatively small, condition (c) guarantees that  $(x_n)_{n=0}^\infty$  contains only composite numbers. The role of the parameters  $c_i$  is to minimize the solution corresponding to a given covering system.

As mentioned earlier, the current record is due to Vsemirnov whose construction is based on the following set of  $t = 17$  quadruples  $(p_i, m_i, r_i, c_i)$ :

$p_i$	3	2	5	7	17	11	47	19	61	23	107	31	1103	181	41	541	2521
$m_i$	4	3	5	8	9	10	16	18	15	24	36	30	48	90	20	90	60
$r_i$	3	1	4	5	2	6	9	14	12	17	8	0	33	80	18	62	48
$c_i$	2	1	2	3	5	6	34	14	29	6	19	21	9	58	11	185	306

Table 1: Vsemirnov's quadruples

Graham, Knuth and Wilf used similar covering systems except with primes 2207, 1087, 4481, 53, 109 and 5779 instead of 23, 1103, 107, 181 and 541. Nicol used primes 53, 109, 5779 instead of 107, 181, 541. A major factor in deciding the size of a solution is the product of the primes in the covering system:  $P = \prod_{i=1}^t p_i$ . The smaller the value of  $P$ , the greater the chance to find a smaller solution. Of all constructions mentioned above, Vsemirnov's attains the smallest  $P$ . It is not known whether a better covering system can be found.

## 2 A new construction

We note that Izotov [5] was the first to propose an alternative approach to a different problem, namely that of construction of Sierpiński numbers, for which the only known solutions involved the use of covering systems. In fact, Erdős [4, Section F13] conjectured that Sierpiński numbers could only be constructed by the use of covering systems.

Similarly, all known examples of Fibonacci-like sequences of composite numbers are based on the existence of a finite covering set of primes  $\{p_1, p_2, \dots, p_t\}$ . In other words, all examples mentioned in the previous section have the property that for every positive integer  $n$ , there exists an  $i \in \{1, 2, \dots, t\}$  with  $x_n \equiv 0 \pmod{p_i}$ .

In this paper we construct a Fibonacci-like sequence of composite numbers for which such a covering set does not appear to exist. Our approach can be summarized as follows:

On one hand, we are going to choose two relatively prime positive integers  $x_0$  and  $x_1$ , such that for every nonnegative integer  $n$ ,  $x_{2n+1}$  is equal to the product of two integers greater than 1, both of which can be written explicitly in terms of  $n$ ,  $x_0$  and  $x_1$ .

On the other hand, we are going to find a finite set of prime numbers  $\{p_1, p_2, \dots, p_t\}$  such that for every nonnegative integer  $n$ ,  $x_{2n} \equiv 0 \pmod{p_i}$  for some  $i \in \{1, 2, \dots, t\}$ .

We will thus obtain the desired Fibonacci-like sequence of composite numbers. Notice that there are different reasons why  $x_n$  is composite depending on the parity of  $n$ .

**Theorem 1.** *Consider the sequence given by  $x_0 = p^2 + q^2$ ,  $x_1 = 2pq + q^2$ ,  $x_n = x_{n-1} + x_{n-2}$  for all  $n \geq 2$ , where  $p$  and  $q$  are integers. Then for every  $n \geq 0$  we have*

$$x_{2n+1} = (pF_n + qF_{n+1})(pL_n + qL_{n+1}). \quad (3)$$

*In particular, if  $p \geq 1$  and  $q \geq 2$ , then  $x_{2n+1}$  is composite for all  $n \geq 0$ .*

*Proof.* It is known that the Fibonacci numbers can be written in matrix form as below

$$\begin{bmatrix} F_{n+1} & F_n \\ F_n & F_{n-1} \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \quad \text{for all } n \geq 1.$$

Using the fact that  $A^{m+n} = A^m A^n$  with  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  it follows that

$$\begin{bmatrix} F_{m+n+1} & F_{m+n} \\ F_{m+n} & F_{m+n-1} \end{bmatrix} = \begin{bmatrix} F_{m+1} & F_m \\ F_m & F_{m-1} \end{bmatrix} \cdot \begin{bmatrix} F_{n+1} & F_n \\ F_n & F_{n-1} \end{bmatrix} \quad \text{from which}$$

$$F_{m+n+1} = F_{m+1}F_{n+1} + F_mF_n.$$

In particular, taking  $m = n$  and  $m = n - 1$  respectively, we obtain

$$F_{2n+1} = F_{n+1}^2 + F_n^2, \quad (4)$$

$$F_{2n} = 2F_nF_{n+1} - F_n^2. \quad (5)$$

It is easy to prove by induction that  $x_n = x_0F_{n-1} + x_1F_n$ . It follows that

$$x_{2n+1} = x_0F_{2n} + x_1F_{2n+1}$$

which after using equalities (4) and (5) gives

$$x_{2n+1} = x_0(2F_nF_{n+1} - F_n^2) + x_1(F_{n+1}^2 + F_n^2) = (x_1 - x_0)F_n^2 + 2x_0F_nF_{n+1} + x_1F_{n+1}^2. \quad (6)$$

Regard the right hand term in the above equation as a quadratic form in the variables  $F_n$  and  $F_{n+1}$ . We want this quadratic form to be reducible over the integers for all  $n$ , which is equivalent to requiring the discriminant  $\Delta = x_0^2 - (x_1 - x_0)x_1$  to be a perfect square.

In other words we want to choose  $x_0$  and  $x_1$  as solutions of the diophantine equation

$$x_0^2 + x_0x_1 - x_1^2 = k^2. \quad (7)$$

It is straightforward to check that  $x_0 = p^2 + q^2$ ,  $x_1 = 2pq + q^2$  is a solution of the above equation. Indeed, with the above choices for  $x_0$  and  $x_1$  we have

$$\begin{aligned} x_0^2 + x_0x_1 - x_1^2 &= (p^2 + q^2)^2 + (p^2 + q^2)(2pq + q^2) - (2pq + q^2)^2 = \\ &= p^4 + 2p^3q - p^2q^2 - 2pq^3 + q^4 = (p^2 + pq - q^2)^2. \end{aligned}$$

This solution can be obtained by using the techniques for solving the general equation  $ax^2 + bxy + cy^2 = k^2$ ; the interested reader may consult [1, Chapter XIII, pp. 404–409]. This explains our choices for  $x_0$  and  $x_1$ . Substituting now  $x_0 = p^2 + q^2$  and  $x_1 = 2pq + q^2$  into (6) we obtain

$$\begin{aligned} x_{2n+1} &= (2pq - p^2)F_n^2 + 2(p^2 + q^2)F_nF_{n+1} + (2pq + q^2)F_{n+1}^2 = \\ &= p^2(2F_nF_{n+1} - F_n^2) + pq(2F_n^2 + 2F_{n+1}^2) + q^2(2F_nF_{n+1} + F_{n+1}^2) = \\ &= p^2F_n(2F_{n+1} - F_n) + 2pq(F_n^2 + F_{n+1}^2) + q^2F_{n+1}(2F_n + F_{n+1}). \end{aligned}$$

We use now some basic identities relating the Fibonacci and the Lucas numbers.

$$\begin{aligned} 2F_{n+1} - F_n &= F_{n+1} + (F_{n+1} - F_n) = F_{n+1} + F_{n-1} = L_n. \\ 2F_n + F_{n+1} &= F_n + (F_n + F_{n+1}) = F_n + F_{n+2} = L_{n+1}. \\ 2(F_n^2 + F_{n+1}^2) &= F_n(2F_n + F_{n+1}) + F_{n+1}(2F_{n+1} - F_n) = F_nL_{n+1} + F_{n+1}L_n. \end{aligned}$$

Substituting these into the last equality, we obtain

$$x_{2n+1} = p^2F_nL_n + pq(F_nL_{n+1} + F_{n+1}L_n) + q^2F_{n+1}L_{n+1} = (pF_n + qF_{n+1}) \cdot (pL_n + qL_{n+1}).$$

Thus, equation (3) is satisfied. If  $p \geq 1$  and  $q \geq 2$  it follows that for all nonnegative integers  $n$  we have that  $pF_n + qF_{n+1} \geq q \geq 2$  and  $pL_n + qL_{n+1} > p + q \geq 3$ . It follows that  $x_{2n+1}$  is always composite. The proof is complete.  $\square$

From Theorem 1 it follows that if one chooses  $x_0 = p^2 + q^2$  and  $x_1 = 2pq + q^2$ , then for every  $n \geq 0$  we have that  $x_{2n+1}$  is composite. It remains to ensure that for every  $n \geq 0$ ,  $x_{2n}$  is also composite. In order to achieve this, we construct a finite partial covering set as described below.

We are looking for a collection of quadruples  $\{(p_i, m_i, r_i, c_i)\}_{i=1}^{i=t}$  such that

- (a) each  $p_i$  is a prime;
- (b) every  $p_i$  divides  $F_{m_i}$ , the  $m_i$ -th Fibonacci number;
- (c) every even positive integer  $2n$  satisfies at least a congruence  $2n \equiv r_i \pmod{m_i}$  for some  $i = 1, 2, \dots, t$ . In other words,  $\{(r_i, m_i)\}_{i=0}^{i=t}$  is a partial covering system, as it covers all even integer values.

For every  $1 \leq i \leq t$ , we have  $1 \leq c_i \leq p_i - 1$ . These values are going to come into play later on, as we will require a certain system of congruences to be compatible.

Suppose we found such a set of quadruples. Choose  $x_0$  and  $x_1$  such that

$$x_0 \equiv c_i F_{m_i - r_i} \pmod{p_i}, \quad x_1 \equiv c_i F_{m_i - r_i + 1} \pmod{p_i} \quad \text{for all } 1 \leq i \leq t. \quad (8)$$

Let  $n \geq 0$  and let  $1 \leq i \leq t$  be such that  $2n \equiv r_i \pmod{m_i}$ . The existence of such  $i$  is guaranteed by condition (b). From (2), we have

$$\begin{aligned} x_{2n} &= x_0 F_{2n-1} + x_1 F_{2n} \equiv c_i (F_{m_i - r_i} \cdot F_{2n-1} + F_{m_i - r_i + 1} \cdot F_{2n}) \pmod{p_i} \\ &\equiv c_i F_{2n + m_i - r_i} \pmod{p_i} \\ &\equiv c_i F_{sm_i} \pmod{p_i} \\ &\equiv 0 \pmod{p_i}. \end{aligned}$$

Thus,  $x_{2n} \equiv 0 \pmod{p_i}$  and therefore composite. There is however one major difficulty in finding a partial covering system with the properties (a), (b) and (c) listed above, as every odd prime  $p_i$  has to satisfy the congruence  $p_i \equiv 1 \pmod{4}$ .

Let us explain why that is the case. First, we need the following simple result.

**Lemma 2.** *For any positive odd integer  $m$ , the Fibonacci number  $F_m$  has no prime factors of the form  $4l + 3$ .*

*Proof.* Let  $m$  be odd and let  $p \mid F_m$ . From Cassini's identity, we have  $F_{m+1}^2 - F_m F_{m+2} = (-1)^m$ , which after reducing modulo  $p$  gives  $F_{m+1}^2 \equiv -1 \pmod{p}$ . This implies that  $-1$  is a quadratic residue modulo  $p$ . By the quadratic reciprocity law this is true only when  $p \equiv 1 \pmod{4}$ .  $\square$

Recall that in Theorem 1 we chose  $x_0$  and  $x_1$  such that  $x_0^2 + x_0 x_1 - x_1^2 = k^2$ . On the other hand, we selected  $x_0$  and  $x_1$  as in (8). It follows that for every  $1 \leq i \leq t$  we have that

$$c_i^2 (F_{m_i - r_i}^2 + F_{m_i - r_i} \cdot F_{m_i - r_i + 1} - F_{m_i - r_i + 1}^2) \equiv k^2 \pmod{p_i} \quad (9)$$

Using Cassini's identity again, the above congruence can be simplified to  $c_i^2 \cdot (-1)^{m_i - r_i + 1} \equiv k^2 \pmod{p_i}$ . In particular,  $(-1)^{m_i - r_i + 1}$  is a quadratic residue modulo  $p_i$  for every  $1 \leq i \leq t$ .

We want that for every  $n \geq 0$ , the congruence  $2n \equiv r_i \pmod{m_i}$  holds for some  $1 \leq i \leq t$ . If  $m_i$  is even, then  $r_i$  is even as well, and therefore  $(-1)^{m_i - r_i + 1} = -1$ , which is a quadratic residue modulo  $p_i$  if and only if  $p_i \equiv 1 \pmod{4}$ . If  $m_i$  is odd, then condition (b) states that  $p_i \mid F_{m_i}$ , and hence  $p_i \equiv 1 \pmod{4}$  follows from Lemma 2.

Hence, none of the primes  $p_i$  in the partial covering system with properties (a), (b) and (c) can be of the form  $4l + 3$ . Consider the set  $\{(p_i, m_i, r_i, c_i)\}_{i=1}^{30}$  given in Table 2. It is straightforward to check that this system of quadruples has the desired properties.

$p_i$	$m_i$	$r_i$	$c_i$
2	3	1	1
5	5	1	2
13	7	1	5
17	9	3	11
29	14	2	5
41	20	4	3
61	15	2	41
181	90	8	46
241	120	14	109
281	28	4	207
421	21	3	171
541	90	38	243
1009	126	90	294
1601	80	34	1259
2161	40	10	1706

$p_i$	$m_i$	$r_i$	$c_i$
2521	60	20	636
3041	80	74	790
8641	360	18	4664
20641	120	110	1405
31249	126	42	901
103681	72	54	80856
109441	45	23	16635
141961	35	12	12156
721561	420	180	529617
1461601	252	186	970625
35239681	63	6	25860534
764940961	252	0	562105967
8288823481	105	33	83463210
10783342081	180	162	7785411056
571385160581761	504	222	49367403415248

Table 2: A finite partial covering system with no primes  $\equiv 3 \pmod{4}$

Let us notice that in order to check condition (c), it suffices to test the even numbers  $\leq 5040$ . This is indeed the case since 5040 is the least common multiple of all the  $m_i$ ,  $1 \leq i \leq 30$ .

We are now in position to state the main result of this paper.

**Theorem 3.** *Let  $p = 1$  and  $q = 12951150255508108245872399074061259209531943793351 - 2025195406541068394745828231264515958532145970461367703231950382110924410768870$ .*

*Define a sequence  $(x_n)_{n \geq 0}$  by  $x_0 = p^2 + q^2$ ,  $x_1 = 2pq + q^2$ ,  $x_n = x_{n-1} + x_{n-2}$ , for all  $n \geq 2$ . Then  $\gcd(x_0, x_1) = 1$  and  $x_n$  is composite for all  $n \geq 0$ .*

*Proof.* By our choice of  $x_0$  and  $x_1$ , Theorem 1 immediately implies that  $x_{2n+1}$  is composite for every integer  $n \geq 0$ .

We claim that for every  $n \geq 1$ ,  $x_{2n}$  has a factor in the set  $\{p_1, p_2, \dots, p_{30}\}$ , where the primes are those in Table 2. Indeed, let us first choose  $x_0$  and  $x_1$  according to (8)

$$x_0 \equiv c_i F_{m_i - r_i} \pmod{p_i}, \quad x_1 \equiv c_i F_{m_i - r_i + 1} \pmod{p_i} \quad \text{for all } 1 \leq i \leq 30,$$

where  $(p_i, m_i, r_i, c_i)$  are those given in Table 2. Since  $p = 1$ ,  $x_0 = p^2 + q^2$  and  $x_1 = 2pq + q^2$ , these congruences can be written as

$$1 + q^2 \equiv c_i F_{m_i - r_i} \pmod{p_i}, \quad 2q + q^2 \equiv c_i F_{m_i - r_i + 1} \pmod{p_i} \quad \text{for all } 1 \leq i \leq 30. \quad (10)$$

$p_i$	$q \pmod{p_i}$	$p_i$	$q \pmod{p_i}$
2	0	2521	1934
5	0	3041	455
13	0	8641	1277
17	11	20641	13565
29	20	31249	24574
41	34	103681	22094
61	55	109441	43164
181	149	141961	112001
241	134	721561	170379
281	45	1461601	442479
421	140	35239681	5419606
541	307	764940961	483887978
1009	818	8288823481	6095337569
1601	1347	10783342081	54018520
2161	799	571385160581761	504780818763137

Table 3: Set of congruences satisfied by the solution of (10)

The choices of  $c_i$  ensure that this system has solutions: in particular, a solution is given by the set of congruences listed in Table 3.

The Chinese remainder theorem guarantees that there exists a  $q$  that satisfies all the above congruences. The smallest such value is the 129-digit number mentioned in the statement of Theorem 3.

It remains to argue why we believe this particular sequence  $(x_n)_{n \geq 0}$  does not have a finite covering set of primes. Computer verifications show that there are 803 values of  $0 \leq n \leq 200000$  such that  $x_n$  has no prime factor  $\leq 2 \times 10^6$  or a prime factor among the primes  $p_1, p_2, \dots, p_{30}$  given in table 2. Moreover, any two such terms are mutually prime.

Also, it can be checked that for these choices for  $p$  and  $q$ , the numbers  $pF_{913} + qF_{914}$ ,  $pL_{913} + qL_{914}$ ,  $pF_{943} + qF_{944}$  and  $pL_{943} + qL_{944}$  are all primes of lengths 319, 320, 326 and 326, respectively. Since (3) gives that  $x_{1827} = (pF_{913} + qF_{914})(pL_{913} + qL_{914})$  and  $x_{1887} =$

$(pF_{943} + qF_{944})(pL_{943} + qL_{944})$ , it follows that  $x_{1827}$  and  $x_{1887}$  are both products of exactly two primes, with the least prime factor having lengths 319 and 326, respectively.

As a result, if a finite covering with primes would exist, it has to have at least 803 primes larger than  $2 \times 10^6$ , and among these, at least two primes of length 319 or greater. It seems difficult to prove that the least prime factor of  $x_n$  is unbounded as  $n$  tends to infinity. As already noticed by Filaseta, Finch and Kozek [2], this type of question is open even for much simpler sequences such as  $F_n$ ,  $5 \cdot 2^n + 1$  or  $11 \cdot 5^n - 1$ .

□

## References

- [1] L. E. Dickson, *History of the Theory of Numbers*, Vol. II: Diophantine Analysis, Chelsea Publishing Co., 1966.
- [2] M. Filaseta, C. Finch and M. Kozek, On powers associated with Sierpiński numbers, Riesel numbers and Polignac's conjecture, *J. Number Theory* **128** (2008), 1916–1940.
- [3] R. L. Graham, A Fibonacci-like sequence of composite numbers, *Math. Mag.* **37** (1964), 322–324.
- [4] R. K. Guy, *Unsolved Problems in Number Theory* (3rd ed.), Problem Books in Mathematics, Springer-Verlag, 2004.
- [5] A. S. Izotov, A note on Sierpiński numbers, *Fibonacci Quart.* **33** (1995), 206–207.
- [6] D. E. Knuth, A Fibonacci-like sequence of composite numbers, *Math. Mag.* **63** (1990), 21–25.
- [7] J. W. Nicol, A Fibonacci-like sequence of composite numbers, *Electron. J. Combin.* **6** (1999), Research Paper #R44.
- [8] M. Vsemirnov, A new Fibonacci-like sequence of composite numbers, *J. Integer Seq.* **7** (2004), [Article 04.3.7](#).
- [9] H. S. Wilf, Letters to the Editor, *Math. Mag.* **63** (1990), 284.

---

2010 *Mathematics Subject Classification*: Primary 11B39; Secondary 11A51, 11B37.

*Keywords*: Fibonacci-like recurrence, composite numbers, system of covering congruences.

---

(Concerned with sequences [A082411](#), [A083103](#), [A083104](#), [A083105](#), [A083216](#), and [A221286](#).)

---

Received March 24 2014; revised version received July 8 2014; July 21 2014. Published in *Journal of Integer Sequences*, July 25 2014.

---

Return to [Journal of Integer Sequences home page](#).