

Raising and Lowering Operators for Askey–Wilson Polynomials^{*}

Siddhartha SAHI

Department of Mathematics, Rutgers University, New Brunswick, NJ 08903, USA

E-mail: sahi@math.rutgers.edu

Received September 20, 2006, in final form December 27, 2006; Published online January 04, 2007

Original article is available at <http://www.emis.de/journals/SIGMA/2007/002/>

Abstract. In this paper we describe two pairs of raising/lowering operators for Askey–Wilson polynomials, which result from constructions involving very different techniques. The first technique is quite elementary, and depends only on the “classical” properties of these polynomials, viz. the q -difference equation and the three term recurrence. The second technique is less elementary, and involves the one-variable version of the double affine Hecke algebra.

Key words: orthogonal polynomials; Askey–Wilson polynomials; q -difference equation; three term recurrence; raising operators; lowering operators; root systems; double affine Hecke algebra

2000 Mathematics Subject Classification: 33D45; 33D52; 33D80

Dedicated to the memory of Vadim Kuznetsov

1 Introduction

One of the approaches to proving integrality of the (q, t) -Kostka coefficients is the idea, due to Kirillov–Noumi [9, 10] and Lapointe–Vinet [16, 17], of using raising operators for Macdonald polynomials. (See also [6, 7, 11, 21] for other approaches.) In their proof Kirillov and Noumi give an explicit construction of such raising operators for the Macdonald polynomials $J_\lambda(x; q, t)$ for the root system of type A_{n-1} . They also pose the problem of finding analogous operators for the six-parameter Koornwinder corresponding to the BC_n root system.

This question was also raised by Tom Koornwinder at the Edinburgh conference on symmetric functions organized by Vadim Kuznetsov. The case $n = 1$ corresponds to the celebrated Askey–Wilson polynomials and Koornwinder’s paper [14] from that conference contains partial results in this direction as well as a survey of earlier results.

In this paper we such construct raising/lowering operators for Askey–Wilson polynomials. In fact we describe *two* such pairs of operators, which result from constructions involving very different techniques.

The first technique is quite elementary, and depends only on the “classical” properties of these polynomials, viz. the q -difference equation and the three term recurrence. Therefore it can be applied to all the polynomials in the Askey scheme. After this work was completed, we obtained a recent preprint by T. Koornwinder [15], the main result of which is very close to this approach. Also through [15] we discovered still earlier work of G. Bangerezako [2] which obtains similar operators based on an *ad-hoc* factorization of the Askey–Wilson operator. Our approach however is more direct and quite short.

^{*}This paper is a contribution to the Vadim Kuznetsov Memorial Issue “Integrable Systems and Related Topics”. The full collection is available at <http://www.emis.de/journals/SIGMA/kuznetsov.html>

The second technique is less elementary and involves the one-variable version of the powerful Hecke algebra method as described in [18, 19, 20, 22, 23, 24]. This approach is related to a fairly remarkable mathematical object – the double affine Hecke algebra (see [3, 4, 5, 22]). The calculations, while non-trivial to carry out, are conceptually rather straightforward. The raising/lowering operators so obtained are different from those coming from the “classical” method. This method also provides a new factorization of the Askey–Wilson operator described in Lemma 3, which is much simpler than that of Bangerezako.

In subsequent work, we hope to extend these methods to construct raising operators for Koornwinder polynomials [13, 25, 22, 23].

2 The classical approach

2.1 Askey Wilson polynomials

The q -hypergeometric series is given by the formula

$${}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} \middle| q; y \right) = \sum_{k \geq 0} \frac{(a_1, \dots, a_r)_k}{(b_1, \dots, b_s)_k} (-1)^{(1+s-r)k} q^{(1+s-r)\binom{k}{2}} \frac{y^k}{(q)_k},$$

where the “ q -Pochhammer symbols” are defined by

$$\begin{aligned} (a, b, c, \dots)_k &:= (a)_k (b)_k (c)_k \cdots, \\ (a)_k &:= (1-a)(1-aq) \cdots (1-aq^{k-1}). \end{aligned}$$

The Askey–Wilson polynomials [1] are defined by the formula

$$P_n(z; a, b, c, d|q) = \frac{(ab, ac, ad)_n}{a^n (abcdq^{n-1})_n} {}_4\phi_3 \left(\begin{matrix} q^{-n}, abcdq^{n-1}, az, az^{-1} \\ ab, ac, ad \end{matrix} \middle| q; q \right).$$

Since $(q^{-n})_k$ vanishes for $k > n$, we have

$${}_4\phi_3(\cdots) = \sum_{k=0}^n \left[\frac{(abcdq^{n-1})_k}{(ab, ac, ad)_k} \right] \left[\frac{(q^{-n})_k q^k}{(q)_k} \right] (az)_k (az^{-1})_k.$$

It follows that P_n is a Laurent polynomial of degree n , which is moreover symmetric in z and z^{-1} and is of the form

$$P_n(z; a, b, c, d|q) = (z^n + z^{-n}) + \text{lower terms.}$$

It is also symmetric in $\{a, b, c, d\}$ although this is not entirely obvious from the formula above.

We have chosen to normalize P_n in order to make it monic. Of course there are several other possible normalizations, and we discuss some of these below.

First of all, we remark that formula (3.1.7) of [12] considers the polynomial

$$\frac{(ab, ac, ad)_n}{a^n} {}_4\phi_3 \left(\begin{matrix} q^{-n}, abcdq^{n-1}, az, az^{-1} \\ ab, ac, ad \end{matrix} \middle| q; q \right)$$

which is $(abcdq^{n-1})_n$ times our P_n .

Next, since the Askey–Wilson polynomial is symmetric in z, z^{-1} , it can be expressed as an (ordinary) polynomial of degree n in

$$x = (z + z^{-1})/2.$$

The function $p_n(x)$ considered in (3.1.5) of [12], is monic in x , and hence is related to our normalization P_n by the formula

$$p_n\left(\frac{z+z^{-1}}{2}\right) = 2^{-n}P_n(z).$$

Finally, the polynomials P_n are orthogonal with respect the inner product $\langle \cdot, \cdot \rangle$ defined in (3.1.2) of [12]. If we define

$$Q_n = \gamma_n P_n, \tag{1}$$

where

$$\gamma_n = \frac{(abq^n, acq^n, adq^n, bcq^n, bdq^n, cdq^n, q^{n+1})_\infty}{(abcdq^{2n})_\infty} (abcdq^{n-1})_n.$$

Then Q_n is dual to P_n in the sense that

$$\langle P_m, Q_n \rangle = \delta_{m,n}.$$

2.2 Raising and lowering operators

The main result of this section are the following raising and lowering operators for the Askey–Wilson polynomials:

Theorem 1. *For all $n > 1$, the Askey–Wilson polynomials satisfy the relations*

$$\begin{aligned} [D(z+z^{-1}) - \lambda_{n-1}(z+z^{-1}) - \alpha_n(\lambda_n - \lambda_{n-1})] P_n &= (\lambda_{n+1} - \lambda_{n-1}) P_{n+1}, \\ [D(z+z^{-1}) - \lambda_{n+1}(z+z^{-1}) - \alpha_n(\lambda_n - \lambda_{n+1})] Q_n &= (\lambda_{n-1} - \lambda_{n+1}) Q_{n-1}, \end{aligned}$$

where D , λ_n and α_n are as in (3) and (6) below.

Proof. The proof involves two key properties of the Askey–Wilson polynomials.

The first property is the ‘ q -difference equation’ from (3.1.7) of [12] which asserts that P_n is an eigenfunction for the Askey–Wilson operator, i.e.

$$DP_n(z) = \lambda_n P_n(z). \tag{2}$$

The operator and its eigenvalue are defined by

$$\begin{aligned} D &= A(z)(T_q - 1) + A(z^{-1})(T_{q^{-1}} - 1), \\ \lambda_n &= (q^{-n} - 1)(1 - abcdq^{n-1}) = (q^{-n} + abcdq^{n-1}) - (1 + abcdq^{-1}), \end{aligned} \tag{3}$$

where $A(z)$ is the rational function

$$A(z) = \frac{(az, bz, cz, dz)_1}{(z^2)_2} = \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} \tag{4}$$

and T_q is the shift operator

$$T_q f(z) = f(qz).$$

(To forestall possible confusion we emphasize that, in accordance with custom, we think of $f(z)$ as a Laurent polynomial rather than as a function of z . This means that we have $T_q(z^k) = q^k z^k$ rather than $T_q(z^k) = q^{-k} z^k$.)

The second key property of these polynomials is the ‘normalized recurrence relation’ from (3.1.5) of [12] which can be rewritten in the form

$$(z + z^{-1}) P_n = P_{n+1} + \alpha_n P_n + \frac{\gamma_{n-1}}{\gamma_n} P_{n-1} \quad \text{for } n > 1, \quad (5)$$

where

$$\alpha_n = a + 1/a - \frac{a (bcq^{n-1}, bdq^{n-1}, cdq^{n-1}, q^n)_1}{(abcdq^{2n-2})_2} - \frac{(abq^n, acq^n, adq^n, abcdq^{n-1})_1}{a (abcdq^{2n-1})_2}. \quad (6)$$

We combine these two properties as follows:

First apply the operators $D - \lambda_{n-1}$ and $D - \lambda_{n+1}$, respectively, to the recurrence relation to get

$$\begin{aligned} (D - \lambda_{n-1}) (z + z^{-1} - \alpha_n) P_n &= (\lambda_{n+1} - \lambda_{n-1}) P_{n+1}, \\ (D - \lambda_{n+1}) (z + z^{-1} - \alpha_n) P_n &= \frac{\gamma_{n-1}}{\gamma_n} (\lambda_{n-1} - \lambda_{n+1}) P_{n-1}. \end{aligned}$$

Finally simplify, using the q -difference equation (2), to get

$$\begin{aligned} [D(z + z^{-1}) - \lambda_{n-1}(z + z^{-1}) - \alpha_n(\lambda_n - \lambda_{n-1})] P_n &= (\lambda_{n+1} - \lambda_{n-1}) P_{n+1}, \\ [D(z + z^{-1}) - \lambda_{n+1}(z + z^{-1}) - \alpha_n(\lambda_n - \lambda_{n+1})] Q_n &= (\lambda_{n-1} - \lambda_{n+1}) Q_{n-1} \end{aligned}$$

as desired. ■

3 The Hecke algebra approach

In this section we provide raising/lowering operators for Askey–Wilson polynomials based on Hecke algebra considerations [22, 23]. Once again the main idea is quite straightforward, although the calculations are a little more intricate. The resulting formulas are different and perhaps slightly simpler.

3.1 The Hecke algebra

The key to this approach are the involutions s_1, s_0 which act on Laurent polynomials as follows:

$$s_1 f(z) = f(z^{-1}) \quad \text{and} \quad s_0 f(z) = f(qz^{-1}).$$

Once again we regard these operators as acting on polynomials, rather than functions, so that we have

$$s_1(z^k) = z^{-k} \quad \text{and} \quad s_0(z^k) = q^k z^{-k}.$$

These operators provide a factorization of the q -shift operators, and one has

$$s_1 s_0 = T_q \quad \text{and} \quad s_0 s_1 = T_{q^{-1}}.$$

The affine Hecke algebra [22, 23] is the algebra of operators generated by the two operators T_0 and T_1 defined as

$$T_i := t_i + r_i (s_i - 1),$$

where

$$\begin{aligned} t_0 &= -cd/q, & r_0 &= \frac{(z-c)(z-d)}{(z^2-q)}, \\ t_1 &= -ab, & r_1 &= \frac{(1-az)(1-bz)}{(1-z^2)}. \end{aligned} \quad (7)$$

Remark 1. The operator T_i as defined here is $t_i^{1/2}$ times the corresponding operator from [22, 23]. This accounts for the slight difference between the formulas here and in [23].

From the definition of T_1 it follows that a polynomial f is symmetric in z, z^{-1} , if and only if

$$T_1 f = t_1 f. \quad (8)$$

Consequently, if g is any polynomial, then the quadratic relation (9) implies that $(T_1 + 1)g$ is a symmetric polynomial. The operators T_i are deformations of s_i and satisfy a quadratic relation. For the convenience of the reader unfamiliar with the Hecke algebra, we give a proof this relation.

Lemma 1. *The operators T_i satisfy the relation*

$$(T_i - t_i)(T_i + 1) = 0. \quad (9)$$

Proof. Define $s_i(r_i) = r'_i$, then we claim that

$$r_i + r'_i = t_i + 1 \quad (10)$$

To see this, we calculate for $i = 0$,

$$\begin{aligned} r_0 + r'_0 &= \frac{(z-c)(z-d)}{(z^2-q)} + \frac{(qz^{-1}-c)(qz^{-1}-d)}{(q^2z^{-2}-q)} \\ &= \frac{(z-c)(z-d)}{(z^2-q)} - \frac{(q-cz)(q-dz)}{q(z^2-q)} \\ &= \frac{q(z^2 - cz - dz + cd) - (q^2 - qc z - qdz + cdz^2)}{q(z^2 - q)} \\ &= \frac{qz^2 + qcd - q^2 - cdz^2}{q(z^2 - q)} = \frac{(q - cd)(z^2 - q)}{q(z^2 - q)} \\ &= 1 - \frac{cd}{q} = 1 + t_0. \end{aligned}$$

The calculation for $i = 1$ is similar and simpler.

Now the quadratic relation can be proved as follows:

$$\begin{aligned} (T_1 - t_1)(T_1 + 1) &= r_i(s_i - 1)[t_i + 1 + r_i(s_i - 1)] \\ &= r_i[(t_i + 1)(s_i - 1) + (s_i r_i)(s_i - 1) - r_i(s_i - 1)] \\ &= r_i[(t_i + 1)(s_i - 1) + (r'_i s_i)(s_i - 1) - r_i(s_i - 1)] \\ &= r_i[(t_i + 1)(s_i - 1) + r'_i(1 - s_i) - r_i(s_i - 1)] \\ &= r_i(t_i + 1 - r_i - r'_i)(s_i - 1) \stackrel{\text{by (10)}}{=} 0. \quad \blacksquare \end{aligned}$$

The following result is an immediate consequence

Corollary 1. *The operators T_i are invertible, with*

$$t_i T_i^{-1} = T_i - t_i + 1. \quad (11)$$

We will also need a number of commutation results between the T_i and the multiplication operator by z . They follow directly from the definition, and we leave the (easy) proof to the reader.

Lemma 2. *The operators T_i satisfy the following commutation relations*

$$\begin{aligned} z t_0 T_0^{-1} &= q T_0 z^{-1} + c + d, \\ (T_1 + 1) z^{-1} &= t_1 z^{-1} + z T_1 + a + b, \\ (T_1 + 1) z &= z + t_1 z^{-1} T_1^{-1} - (a + b). \end{aligned}$$

3.2 Nonsymmetric Askey–Wilson polynomials

The next ingredient in the Hecke algebra method are the nonsymmetric Askey–Wilson polynomials. These are certain Laurent polynomials, E_n , $n \in \mathbb{Z}$, which can be characterized up to multiples as eigenfunctions of the operator

$$Y = T_1 T_0.$$

More precisely, one has

$$Y E_n = \mu_n E_n, \tag{12}$$

where

$$\mu_n = \begin{cases} q^n & \text{for } n < 0, \\ q^n t_1 t_0 = q^{n-1} abcd & \text{for } n \geq 0. \end{cases}$$

The symmetric Askey–Wilson polynomials $P_{|n|}$ are closely related to $E_{\pm n}$. Up to normalization, one has $P_0 = E_0 = 1$, while for $n > 0$ we have up to a scalar

$$P_{|n|} \sim (T_1 + 1) E_{\pm n} = c_n^{\pm} E_n + c_{-n}^{\pm} E_{-n}. \tag{13}$$

The explicit formula for the coefficients c_n^{\pm} and c_{-n}^{\pm} is known, but will not be needed in what follows.

We now define a slight variant of the Askey–Wilson operator, as follows:

$$D' = A(z) (T_q - s_1) + A(z^{-1}) (T_{q^{-1}} s_1 - 1). \tag{14}$$

Observe that D and D' have the same action on functions which are symmetric z and z^{-1} , thus the Askey–Wilson polynomials satisfy

$$D' P_n = D P_n = \lambda_n P_n.$$

Just as the operator s_0 and s_1 factorize the q -shift operator, it turns out that the operators T_0 and T_1 provide a factorization of D' .

Lemma 3. *The operator D' of formula (14) admits the following factorization:*

$$D' = (T_1 + 1) (T_0 - t_0).$$

Proof. To prove this, we calculate as follows

$$\begin{aligned} (T_1 + 1) (T_0 - t_0) &= (t_1 + r_1 (s_1 - 1) + 1) (r_0 (s_0 - 1)) \stackrel{\text{by (10)}}{=} (r'_1 + r_1 s_1) (r_0 s_0 - r_0) \\ &= r'_1 r_0 (s_0 - 1) + r_1 \tilde{r}_0 (s_1 s_0 - s_1) = r'_1 r_0 (T_q^{-1} s_1 - 1) + r_1 \tilde{r}_0 (T_q - s_1), \end{aligned}$$

where $r'_1 = r_1 (z^{-1})$ and $\tilde{r}_0 = r_0 (z^{-1})$.

Comparing the formulas for r_i (7) and $A(z)$ (4), we conclude that

$$A(z) = r_1 \tilde{r}_0 \quad \text{and} \quad A(z^{-1}) = r'_1 r_0$$

which completes the proof. ■

3.3 Raising and lowering operators

To state our main result we need some notation. We write

$$e_1 = a + b + c + d, \quad e_3 = abc + abd + acd + bcd. \quad (15)$$

Also recall that for $n \geq 0$, λ_n is the symmetric Askey–Wilson eigenvalue as in (3). For $n < 0$ we define

$$\lambda_n = \lambda_{|n|}$$

and for all integral n we set

$$\beta_n = \frac{\lambda_n + 1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} e_1 - \frac{1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} e_3. \quad (16)$$

Theorem 2. *The Askey–Wilson polynomials satisfy the following relations:*

$$[D'z + (1 - q^{1-n})(z + z^{-1}) + \beta_{-n}] P_n = (q^n abcd - q^{1-n}) P_{n+1}, \quad n \geq 0, \quad (17)$$

$$[D'z + (1 - q^n abcd)(z + z^{-1}) + \beta_n] Q_n = (q^{1-n} - q^n abcd) Q_{n-1}, \quad n > 0. \quad (18)$$

The key for the proof is the “affine intertwiner” for the nonsymmetric Askey–Wilson polynomials from [23]. This involves the additional parameters u_0 and u_1 , which satisfy the relations

$$a = t_1^{1/2} u_1^{1/2}, \quad b = -t_1^{1/2} u_1^{-1/2}, \quad c = q^{1/2} t_0^{1/2} u_0^{1/2}, \quad d = -q^{1/2} t_0^{-1/2} u_0^{1/2}.$$

Now from Theorem 1.2 of [23] we have, up to a multiple,

$$E_n \sim (a_n U_0 + b_n) E_{-n-1}, \quad (19)$$

where

$$a_n = (q^{\bar{n}} - q^{-\bar{n}-1}) \quad \text{and} \quad b_n = q^{\bar{n}}(u_0^{-1/2} - u_0^{1/2}) + q^{-1/2}(u_1^{-1/2} - u_1^{1/2})$$

with

$$q^{\bar{n}} = \mu_n t_1^{1/2} t_0^{1/2}$$

and U_0 is the operator

$$U_0 = q^{-1/2} t_0^{1/2} T_0^{-1} z.$$

We will derive Theorem 2 from formula (19); however some remarks are in order before we proceed:

1. There is a typo in the statement of formula (19) in Theorem 1.2 of [23], namely n and $-n-1$ have been inadvertently switched. This is easily seen by comparison with Theorem 4.1 from which Theorem 1.2 is derived.
2. The formula for U_0 here is slightly different from that in [23] because of the difference in T_0 – see Remark 1).
3. Although Theorem 1.2 in [23] is only stated (and needed) for $n \geq 0$, it is easy to see that after the correction above it holds for all integer n .

4. Finally, we note that the ideas of [22] and [23] work in the more general setting of Koornwinder polynomials, and they involve the affine intertwiner S_0 , which can also be written as

$$S_0 = [Y, z^{-1}T_1^{-1}].$$

It is expected that this operator will play a key role in the raising operators for Koornwinder polynomials.

We now give the proof of Theorem 2.

Proof. We first simplify (19) by multiplying through by $q^{1/2}t_1^{1/2}t_0$. This gives

$$\begin{aligned} E_n &\sim [(\mu_n - \mu_{-n-1})t_0T_0^{-1}z - \mu_n(c+d) - t_0(a+b)]E_{-n-1} \\ &\sim \left[t_0T_0^{-1}z - \frac{\mu_n(c+d) + t_0(a+b)}{\mu_n - \mu_{-n-1}} \right] E_{-n-1}. \end{aligned}$$

Replacing n by $n-1$, we get

$$E_{n-1} \sim \left[t_0T_0^{-1}z - \frac{\mu_{n-1}(c+d) + t_0(a+b)}{\mu_{n-1} - \mu_{-n}} \right] E_{-n}.$$

For ease in subsequent calculations, we write this as

$$E_{n-1} \sim (t_0T_0^{-1}z - \kappa_n)E_{-n}, \quad (20)$$

where

$$\kappa_n = \frac{\mu_{n-1}y + t_0x}{\mu_{n-1} - \mu_{-n}}, \quad x = a + b, \quad y = c + d. \quad (21)$$

The key idea to obtain a raising operator for P_n is as follows: By formula (13), for $n > 1$, $P_{|n|}$ is a combination of E_n and E_{-n} . We first kill off the E_n component. This can be accomplished by applying the operator $Y - \mu_n$ to P_n . However it is more convenient (and equivalent) to apply the operator

$$t_1t_0(Y^{-1} - \mu_n^{-1}) = t_1t_0T_0^{-1}T_1^{-1} - \frac{t_1t_0}{\mu_n}$$

For $n \neq 0$ we have $\frac{t_1t_0}{\mu_n} = \mu_{-n}$. Thus since P_n is symmetric, formula (8) implies that up to a non-zero multiple, one has

$$(t_0T_0^{-1} - \mu_{-n})P_n \sim E_{-n}. \quad (22)$$

Although the argument given above only applies for $n \neq 0$, it is easy to see that formula (22) is true (up to a non-zero multiple) for $n = 0$ as well! Now combining formulas (13), (20), and (22) we conclude that up to a multiple, we have

$$P_{|n-1|} \sim R_n P_{|n|},$$

where

$$R_n = (T_1 + 1)(t_0T_0^{-1}z - \kappa_n)(t_0T_0^{-1} - \mu_{-n}).$$

The main problem now is to simplify the expression of the operator R_n using properties of $P_{|n|}$.

We first calculate using Lemma 2 and (11), as follows

$$\begin{aligned}
& (t_0 T_0^{-1} z - \kappa_n) (t_0 T_0^{-1} - \mu_{-n}) \\
&= t_0 T_0^{-1} z t_0 T_0^{-1} - t_0 T_0^{-1} (\mu_{-n} z + \kappa_n) + \kappa_n \mu_{-n} \\
&= t_0 T_0^{-1} (q T_0 z^{-1} + y) - t_0 T_0^{-1} (\mu_{-n} z + \kappa_n) + \kappa_n \mu_{-n} \\
&= t_0 T_0^{-1} (y - \mu_{-n} z - \kappa_n) + q t_0 z^{-1} + \kappa_n \mu_{-n} \\
&= (T_0 - t_0 + 1) (y - \mu_{-n} z - \kappa_n) + (q t_0 z^{-1} + \kappa_n \mu_{-n}).
\end{aligned}$$

Applying $T_1 + 1$ to this, we get by Lemma 3

$$R_n = (D' + T_1 + 1) (-\mu_{-n} z + y - \kappa_n) + (T_1 + 1) (q t_0 z^{-1} + \kappa_n \mu_{-n}).$$

To simplify this further we note that the commutation relations of Lemma 2 can be rewritten as and hence imply

$$\begin{aligned}
(T_1 + 1) z^{-1} &= t_1 z^{-1} + z T_1 + x, \\
(T_1 + 1) z &= z + t_1 z^{-1} T_1^{-1} - x.
\end{aligned}$$

Also on $P_{|n|}$, D' acts by λ_n while T_1 acts by t_1 . Therefore R_n acts by the operator

$$\begin{aligned}
& -\mu_{-n} [D' z + z + z^{-1} - x] + (\lambda_n + t_1 + 1) (y - \kappa_n) \\
&+ q t_0 [t_1 (z + z^{-1}) + x] + (t_1 + 1) \kappa_n \mu_{-n}.
\end{aligned}$$

Dividing by $-\mu_{-n}$, we see that up to a multiple

$$\left[D' z + \left(1 - \frac{q t_1 t_0}{\mu_{-n}} \right) (z + z^{-1}) + \beta'_n \right] P_{|n|} \sim P_{|n-1|}, \quad (23)$$

where

$$\beta'_n = -x - \frac{q t_0}{\mu_{-n}} x - \frac{\lambda_n + t_1 + 1}{\mu_{-n}} (y - \kappa_n) - (t_1 + 1) \kappa_n.$$

We now show that β'_n is equal to β_n from formula (16). For this we simplify the expression, substituting for κ_n , using (20) above to get

$$\begin{aligned}
\beta'_n &= -x - \frac{q t_0}{\mu_{-n}} x + \frac{(\lambda_n + t_1 + 1) \mu_{-n} y + t_0 x}{\mu_{-n} \mu_{n-1} - \mu_{-n}} - (t_1 + 1) \frac{\mu_{n-1} y + t_0 x}{\mu_{n-1} - \mu_{-n}} \\
&= \frac{\lambda_n + 1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} y + \frac{1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} t_1 y + c_1 x + c_2 (q t_0 x),
\end{aligned}$$

where c_1 and c_2 are moderately complicated expressions which can be computed explicitly. However, we can save some computation by observing that since the result is *a priori* symmetric in $\{a, b, c, d\}$, c_1 and c_2 must reduce to the coefficients of y and $-t_1 y$ respectively.

It follows then that we have

$$\begin{aligned}
\beta'_n &= \frac{\lambda_n + 1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} (x + y) + \frac{1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} (q t_0 x + t_1 y) \\
&= \frac{\lambda_n + 1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} e_1 - \frac{1 - \mu_{n-1}}{\mu_{n-1} - \mu_{-n}} e_3 = \beta_n.
\end{aligned}$$

Replacing n by $-n$, formula (23) becomes

$$\left[D' z + \left(1 - \frac{q t_1 t_0}{\mu_n} \right) (z + z^{-1}) + \beta_{-n} \right] P_{|n|} \sim P_{|-n-1|}.$$

For $n \geq 0$, we have $\mu_n = q^n t_1 t_0$ and this becomes

$$[D'z + (1 - q^{1-n})(z + z^{-1}) + \beta_{-n}] P_n \sim P_{n+1} \quad (24)$$

which is (17) up to a multiple.

For $n \geq 1$, we have $\mu_{-n} = q^{-n}$ and formula (23) becomes

$$[D'z + (1 - q^n abcd)(z + z^{-1}) + \beta_n] P_n \sim P_{n-1} \quad (25)$$

which is (18) up to a multiple.

It remains then only to calculate the multiples in (24), (25).

To determine the multiple (24), it suffices to calculate the coefficient of z^{n+1} on the left. For this we divide the left side of (24) by z^{n+1} and take the limit as $z \rightarrow \infty$. This gives

$$\begin{aligned} \lim_{z \rightarrow \infty} \frac{1}{z^{n+1}} (A(z) q^{n+1} z^{n+1} - A(z^{-1}) z^{n+1} + (1 - q^{1-n}) z^{n+1}) \\ = \frac{abcd}{q} q^{n+1} - 1 + 1 - q^{1-n} = q^n abcd - q^{1-n} \end{aligned}$$

which proves formula (17).

To determine the multiple in (25), we rewrite (25) in the form

$$[D'z + (1 - q^n abcd)(z + z^{-1}) + \beta_n] P_n = c_n (q^{1-n} - q^n abcd) \frac{\gamma_{n-1}}{\gamma_n} P_{n-1}, \quad n > 0$$

for some unknown constant c_n . It then suffices to show that $c_n = 1$. Subtracting this from (17) we have

$$\begin{aligned} [(q^n abcd - q^{1-n})(z + z^{-1}) + \beta_{-n} - \beta_n] P_n \\ = (q^n abcd - q^{1-n}) \left(P_{n+1} + c_n \frac{\gamma_{n-1}}{\gamma_n} P_{n-1} \right) \end{aligned}$$

or

$$(z + z^{-1}) P_n = P_{n+1} + \frac{\beta_n - \beta_{-n}}{q^n abcd - q^{1-n}} P_n + c_n \frac{\gamma_{n-1}}{\gamma_n} P_{n-1}.$$

Comparing this with the recurrence relation (5) we deduce $c_n = 1$, which proves formula (18) and completes the proof of the theorem.

We note in passing that comparison with (5) also proves the following identity (which can be verified independently):

$$\alpha_n = \frac{\beta_n - \beta_{-n}}{q^n abcd - q^{1-n}}. \quad \blacksquare$$

Acknowledgements

We would like to thank the (anonymous) referee for several insightful suggestions which have improved the paper considerably. The referee has also pointed out that one can give an alternative proof of formulas (17) and (18) by combining Theorem 1 with the following identity relating the operators D and D' :

$$\begin{aligned} [(1 - q^2)D'z + q^2 D(z + z^{-1}) - q(z + z^{-1})D] f \\ = (1 - q) [(e_1 - e_3) - (1 - abcd)(z + z^{-1})] f, \end{aligned}$$

which holds for all symmetric Laurent polynomials f .

References

- [1] Askey R., Wilson J., Some basic hypergeometric polynomials that generalize Jacobi polynomials, *Mem. Amer. Math. Soc.* **319** (1985), 1–53.
- [2] Bangerezako G., The factorization method for the Askey–Wilson polynomials, *J. Comput. Appl. Math.* **107** (1999), 219–232.
- [3] Cherednik I., Double affine Hecke algebras, Knizhnik–Zamolodchikov equations, and Macdonald’s operators, *Int. Math. Res. Not.* (1992), no. 9, 171–180.
- [4] Cherednik I., Double affine Hecke algebras and Macdonald’s conjectures, *Ann. of Math.* **141** (1995), 191–216.
- [5] Etingof P., Oblomkov A., Rains E., Generalized double affine Hecke algebras of rank 1 and quantized del Pezzo surfaces, [math.QA/0406480](https://arxiv.org/abs/math.QA/0406480).
- [6] Garsia A., Remmel R., Plethystic formulas and positivity for q, t -Kostka coefficients, in *Mathematical Essays in Honor of Gian-Carlo Rota*, Editors B. Sagan and R. Stanley, *Progr. Math.* **161** (1998), 245–262.
- [7] Garsia A., Tesler G., Plethystic formulas for the Macdonald q, t -Kostka coefficients, *Adv. Math.* **123** (1996), 144–222.
- [8] Ion B., Sahi S., Triple groups and Cherednik algebras, *Contemp. Math.* **417** (2006), 183–206, [math.QA/0304186](https://arxiv.org/abs/math.QA/0304186).
- [9] Kirillov A., Noumi M., q -difference raising operators for Macdonald polynomials and the integrality of transition coefficients, in *Algebraic Methods and q -Special Functions*, *CRM Proceedings and Lecture Notes* **22** (1999), 227–243, [q-alg/9605005](https://arxiv.org/abs/q-alg/9605005).
- [10] Kirillov A., Noumi M., Affine Hecke algebras and raising operators for Macdonald polynomials, *Duke Math. J.* **93** (1998), 1–39, [q-alg/9605004](https://arxiv.org/abs/q-alg/9605004).
- [11] Knop F., Integrality of two variable Kostka functions, *J. Reine Angew. Math.* **482** (1997), 177–189, [q-alg/9603027](https://arxiv.org/abs/q-alg/9603027).
- [12] Koekoek R., Swarttouw R., The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue, Delft University of Technology, Department of Technical Mathematics and Informatics, Report no. 98-17 (1998), <http://aw.twi.tudelft.nl/~koekoek/askey/ch3/par1/par1.html>.
- [13] Koornwinder T., Askey–Wilson polynomials for root systems of type BC, *Contemp. Math.* **138** (1992), 189–204.
- [14] Koornwinder T., Lowering and raising operators for some special orthogonal polynomials, *Contemp. Math.* **417** (2006), 227–238.
- [15] Koornwinder T., The structure relation for Askey–Wilson polynomials, *J. Comput. Appl. Math.*, to appear, [math.CA/0601303](https://arxiv.org/abs/math.CA/0601303).
- [16] Lapointe L., Vinet L., Creation operators for the Macdonald and Jack polynomials, *Lett. Math. Phys.* **40** (1997), 269–286.
- [17] Lapointe L., Vinet L., Rodrigues formulas for the Macdonald polynomials, *Adv. Math.* **130** (1997), 261–279, [q-alg/9607025](https://arxiv.org/abs/q-alg/9607025).
- [18] Macdonald I., *Affine Hecke algebras and orthogonal polynomials*, Cambridge University Press, 2003.
- [19] Noumi M., Macdonald–Koornwinder polynomials and affine Hecke algebras, *RIMS Kokyuroku* **919** (1995), 44–55 (in Japanese).
- [20] Noumi M., Stokman J., Askey–Wilson polynomials: an affine Hecke algebra approach, in *Laredo Lectures on Orthogonal Polynomials and Special Functions*, Editors R. Alvarez-Nodarse, F. Marcellan and W. Van Assche, Nova Science Publishers, 2004, 111–144, [math.QA/0001033](https://arxiv.org/abs/math.QA/0001033).
- [21] Sahi S., Interpolation, integrality, and a generalization of Macdonald’s polynomials, *Int. Math. Res. Not.* (1996), no. 10, 457–471.
- [22] Sahi S., Nonsymmetric Koornwinder polynomials and duality, *Ann. of Math.* **150** (1999), 267–282, [q-alg/9710032](https://arxiv.org/abs/q-alg/9710032).
- [23] Sahi S., Some properties of Koornwinder polynomials, *Contemp. Math.* **254** (2000), 395–411.
- [24] Stokman J., Koornwinder polynomials and affine Hecke algebras, *Int. Math. Res. Not.* (2000), no. 19, 1005–1042, [math.QA/0002090](https://arxiv.org/abs/math.QA/0002090).
- [25] van Diejen J., Self-dual Koornwinder–Macdonald polynomials, *Invent. Math.* **126** (1996), 319–339, [q-alg/9507033](https://arxiv.org/abs/q-alg/9507033).