

## FREE OBJECTS OVER POSEMIGROUPS IN THE CATEGORY $\mathbf{PoSgr}_\vee$

SHENGWEI HAN, CHANGCHUN XIA, XIAOJUAN GU

**ABSTRACT.** As we all know, the complete lattice  $I_D(S)$  of all  $D$ -ideals of a meet-semilattice  $S$  is precisely the injective hull of  $S$  in the category of meet-semilattices. In this paper, we consider  $sm$ -ideals of posemigroups which can be regarded as a generalization of  $D$ -ideals of meet-semilattices. Unfortunately, the quantale  $\mathcal{R}(S)$  of all  $sm$ -ideals of a posemigroup  $S$  is in general not an injective hull of  $S$ . However,  $\mathcal{R}(S)$  can be seen as a new type of quantale completions of  $S$ . Further, we can see that  $\mathcal{R}(S)$  is also a free object over  $S$  in the category  $\mathbf{PoSgr}_\vee$  of posemigroups with  $sm$ -distributive join homomorphisms.

### 1. Introduction

Bruns and Lakser investigated the injective objects in the category of meet-semilattices with meet-semilattice homomorphisms, and they also gave the concrete form of injective hulls of meet-semilattices in [Bruns and Lakser (1970)]. Since partially ordered monoids (pomonoids for short) can be regarded as a generalization of meet-semilattices, it is natural to consider the injectivity in the category  $\mathbf{PoMon}$  of pomonoids with pomonoid homomorphisms. However, injective objects in  $\mathbf{PoMon}$  are trivial [Lambek et al. (2012)]. Hence, Lambek et al. changed a few of the definitions of pomonoid homomorphisms, chose submultiplicative order-preserving maps as morphisms of the category  $\mathbf{PoMon}_\leq$  of pomonoids to investigate the injectivity in  $\mathbf{PoMon}_\leq$ , where a *submultiplicative map* between pomonoids is a map  $f: (A, \cdot, \leq) \rightarrow (B, *, \leq)$  such that  $f(a) * f(a') \leq f(a \cdot a')$  for all  $a, a' \in A$ . Also, they proved that every pomonoid has an injective hull and gave the concrete form of the injective hulls. Based on the work of Lambek et al., dropping out the unit, Zhang and Laan considered the injective hulls in the category  $\mathbf{PoSgr}_\leq$  of partially ordered semigroups (posemigroups for short) and their submultiplicative order-preserving maps, and constructed the injective hulls for a certain class of posemigroups with respect to a class of order embedding. In [Xia et al. (2017)], Xia, Han and Zhao proved the existence of injective hulls and gave the concrete form of the injective hull of an arbitrary posemigroup in the category  $\mathbf{PoSgr}_\leq$ .

As we all know, injective hulls are unique up to isomorphism. In the category of meet-semilattices, the injective hull of a meet-semilattice  $S$  is precisely the complete lattice  $I_D(S)$  of all  $D$ -ideals of  $S$  with inclusion order [Bruns and Lakser (1970)]. Motivated by the con-

---

We would like to thank the National Natural Science Foundation of China (11531009), the Natural Science Program for Basic Research of Shaanxi Province, China (2015JM1020) and the Fundamental Research Funds for the Central Universities (GK201501001).

Received by the editors 2017-03-28 and, in final form, 2017-08-24.

Transmitted by Walter Tholen. Published on 2017-08-29.

2010 Mathematics Subject Classification: 06F05, 06F07, 08B30.

Key words and phrases: partially ordered semigroup, injective hull, quantale, quantale completion, free object.

© Shengwei Han, Changchun Xia, Xiaojuan Gu, 2017. Permission to copy for private use granted.

struction of the injective hulls of meet-semilattices, Lambek wanted to carry over the way of the construction of injective hulls from meet-semilattices to pomonoids. Hence, the candidate for the injective hulls of pomonoids (posemigroups) may be the object  $\mathcal{R}(S)$  consisting of all down-closed sets that were closed under distributive joins. Unfortunately, the object  $\mathcal{R}(S)$  is in general not the injective hull of a pomonoid (posemigroupp)  $S$  [Lambek et al. (2012)]. Then we naturally ask for the question whether  $\mathcal{R}(S)$  is possible to be some kind of special objects in the category of posemigroups, which is the main motivation of this paper.

In [Han and Zhao (2008)], we investigated the quantale completions of posemigroups, and proved that up to isomorphism the quantale completions of a posemigroupp  $S$  are completely determined by the topological closures on the power-set quantale  $(\mathcal{P}(S), \subseteq, \bullet)$ . In [Han and Zhao (2008), Xia et al. (2016)], we gave two classes of standard quantale completions of a posemigroupp  $S$ —the greatest quantale completion  $((\mathcal{P}(S))_{\mathcal{D}, \subseteq, \otimes})$  and the smallest quantale completion  $(\mathbf{S}^*, \subseteq, \otimes)$ . Whether there is another type of quantale completions for a posemigroupp except two classes of standard quantale completions is an unsolvable question in [Xia et al. (2016)]. In fact, we can also see that  $(\mathbf{S}^*, \subseteq, \otimes)$  is exactly the injective hull of a posemigroupp (pomonoid)  $S$  in the category  $\mathbf{PoSgr}_\leq$  ( $\mathbf{PoMon}_\leq$ ) [Xia et al. (2017)]. In this paper, we shall prove that  $\mathcal{R}(S)$  is a new type of quantale completions of a posemigroupp  $S$ . Moreover, we shall see that  $\mathcal{R}(S)$  is also the free object over  $S$  in the category  $\mathbf{PoSgr}_\vee$  of posemigroups with  $sm$ -distributive join homomorphisms.

A *partially ordered semigroup* (posemigroupp for short) is a semigroup  $(S, \cdot)$  with a partial order  $\leq$  on  $S$  which is compatible with the multiplication  $\cdot$ , that is,  $x \leq y \Rightarrow x \cdot z \leq y \cdot z$  and  $z \cdot x \leq z \cdot y$  for any  $z$  in  $S$ . A *quantale* is a complete lattice  $Q$  with an associative binary operation  $\&$  satisfying  $a \& (\bigvee_{i \in I} b_i) = \bigvee_{i \in I} (a \& b_i)$  and  $(\bigvee_{i \in I} b_i) \& a = \bigvee_{i \in I} (b_i \& a)$  for any  $a \in Q$  and  $\{b_i\}_{i \in I} \subseteq Q$ . Since  $a \& -$  and  $- \& a$  preserve arbitrary joins, they have right adjoints and we shall denote them by  $a \rightarrow -$  and  $- \leftarrow a$  respectively. Thus  $a \& c \leq b$  iff  $c \leq a \rightarrow b$  iff  $a \leq b \leftarrow c$ . A *quantale homomorphism*  $f: P \rightarrow Q$  is both a semigroup homomorphism and a complete lattice homomorphism. A *quantic nucleus*  $j$  on a quantale  $Q$  is a submultiplicative closure operator where a closure operator on  $Q$  is an order-preserving, expansive and idempotent map. It is easy to show that  $j(a \& b) = j(a \& j(b)) = j(j(a) \& b) = j(j(a) \& j(b))$  for all  $a, b \in Q$ . Furthermore, we can also prove that the set  $Q_j$  of fixed points of  $j$  is a quantale with  $a \otimes b = j(a \& b)$  and

$\bigvee_{i \in I} a_i = j(\bigvee_{i \in I} a_i)$ . A subset  $S$  of  $Q$  is called a *quantic quotient* of  $Q$  if there exists some quantic nucleus  $j$  such that  $S = Q_j$ . Let  $j, k$  be two quantic nuclei on  $Q$ . Then we have that  $j \leq k \iff Q_k \subseteq Q_j$ . Clearly, every quantale is necessarily a posemigroupp. For a given semigroup  $(S, \cdot)$ , we denote by  $\mathcal{P}(S)$  the set of all subsets of  $S$ . Clearly,  $(\mathcal{P}(S), \bullet)$  forms a quantale under the inclusion order, where  $A \bullet B = \{a \cdot b : a \in A, b \in B\}$ . When  $A = \{a\}$ , we write  $A \bullet B$  and  $B \bullet A$  simply as  $a \cdot B$  and  $B \cdot a$ . Then  $A \bullet B \subseteq C \iff A \subseteq C \leftarrow B$  and  $B \bullet A \subseteq C \iff A \subseteq B \rightarrow C$ , where  $C \leftarrow B = \{x \in S : x \cdot B \subseteq C\}$  and  $B \rightarrow C = \{x \in S : B \cdot x \subseteq C\}$ .

For notions and concepts concerned, but not explained here, please refer to [Adámek et al. (2004)], [Lambek et al. (2012)] and [Rosenthal (1990)].

## 2. Quantale completions of posemigroups

In order to describe the revised semantic for intuitionistic linear logic, Larchey-Wendling and Galmiche introduced the concept of quantale completions of pomonoids in [Larchey-Wendling and Galmiche (2000)]. However, quantales mentioned in [Larchey-Wendling and Galmiche (2000)] are called *right semiquantales* in [Rosenthal (1990)], which are not the real quantales we usually speak of. Similar to the work of Larchey-Wendling and Galmiche, Han and Zhao considered the concept of quantale completions of posemigroups, and proved that up to isomorphism the quantale completions of a posemigroup  $S$  are completely determined by the topological closures on the power-set quantale  $\mathcal{P}(S)$  [Han and Zhao (2008)].

2.1. DEFINITION. [Han and Zhao (2008)] A pair  $(i, Q)$  is called a *quantale completion* of a posemigroup  $S$  if  $i: S \rightarrow Q$  is a posemigroup embedding into a quantale  $Q$  and  $i(S)$  is join-dense in  $Q$ , that is, for any  $a \in Q$ , there exists  $S_a \subseteq S$  such that  $a = \bigvee_{s \in S_a} i(s)$ .

For convenience of the readers, we shall list some basic concepts and results on quantale completions of posemigroups.

2.2. DEFINITION. [Han and Zhao (2008)] Let  $S$  be a posemigroup. A *topological closure* on the power-set quantale  $\mathcal{P}(S)$  is a quantic nucleus  $j$  on  $\mathcal{P}(S)$  such that  $j(\{x\}) = x \downarrow$  for all  $x \in S$ .

2.3. PROPOSITION. [Han and Zhao (2008)] Let  $S$  be a posemigroup. Then  $(i_Q, Q)$  is a quantale completion of  $S$  if and only if there exists a topological closure  $j$  on  $\mathcal{P}(S)$  and a quantale isomorphism  $f: (\mathcal{P}(S))_j \rightarrow Q$  such that  $i_Q = f \circ i_{(\mathcal{P}(S))_j}$ , where  $i_{(\mathcal{P}(S))_j}(s) = j(\{s\})$ .

2.4. COROLLARY. [Han and Zhao (2008)] Let  $S$  be a posemigroup,  $j$  be a topological closure on  $\mathcal{P}(S)$ . Then  $(i_{(\mathcal{P}(S))_j}, (\mathcal{P}(S))_j)$  is a quantale completion of  $S$ .

In what follows we shall give two classes of standard topological closures—the smallest topological closure  $\mathcal{D}$  and the greatest topological closure  $(\cdot)^*$  (see [Han and Zhao (2008), Xia et al. (2016)]).

$$\forall X \in \mathcal{P}(S), \quad \mathcal{D}(X) = X \downarrow, \quad X^* =: X^{\text{ul}} \cap X^L \cap X^R \cap X^T,$$

where

$$\begin{aligned} X^{\text{ul}} &=: \{s \in S : \forall b \in S, X \subseteq b \downarrow \Rightarrow s \leq b\} \\ X^L &=: \{s \in S : \forall a, b \in S, X \cdot a \subseteq b \downarrow \Rightarrow s \cdot a \leq b\} \\ X^R &=: \{s \in S : \forall a, b \in S, a \cdot X \subseteq b \downarrow \Rightarrow a \cdot s \leq b\} \\ X^T &=: \{s \in S : \forall a, b, c \in S, a \cdot X \cdot c \subseteq b \downarrow \Rightarrow a \cdot s \cdot c \leq b\}. \end{aligned}$$

In this paper, we denote by  $\mathbf{S}^*$  the set of fixed points of  $(\cdot)^*$ , that is,  $\mathbf{S}^* = \{A \in \mathcal{P}(S) : A^* = A\}$ . Let  $(i_1, Q_1)$  and  $(i_2, Q_2)$  be quantale completions of a posemigroup  $S$ . Then there exist two topological closures  $j_1, j_2$  such that  $(\mathcal{P}(S))_{j_1} \cong Q_1$  and  $(\mathcal{P}(S))_{j_2} \cong Q_2$ . If  $j_1 \leq j_2$ , then we say that  $Q_2$  is smaller than  $Q_1$  (or  $Q_1$  is greater than  $Q_2$ ), written as  $Q_2 \sqsubseteq Q_1$ . In this sense, if  $Q_2 \sqsubseteq Q_1$  and  $Q_1 \sqsubseteq Q_2$ , then  $Q_2 \cong Q_1$ . Hence, up to isomorphism  $(\mathcal{P}(S))_{\mathcal{D}}$  is the greatest quantale completion of  $S$  and  $\mathbf{S}^*$  is the smallest quantale completion of  $S$ . The greatest quantale

completion and the smallest quantale completion play an important role in the study of free objects and injective hulls [Kruml and Paseka (2008), Xia et al. (2017)]. Naturally, we consider a question whether there exists another type of quantale completions of posemigroups except two classes of standard quantale completions, which is another motivation of this paper.

### 3. A new type of quantale completions

In this section, we shall prove that  $\mathcal{R}(S)$  can be seen as a new type of quantale completions of a posemigroup  $S$ . Furthermore, we can see that  $\mathbf{S}^* \subseteq \mathcal{R}(S) \subseteq (\mathcal{P}(S))_{\mathcal{Q}}$ . In general,  $\mathbf{S}^*$ ,  $\mathcal{R}(S)$  and  $(\mathcal{P}(S))_{\mathcal{Q}}$  are not the same (see Example 3.13). First, we shall introduce the concept of *sm*-ideals of posemigroups which can be regarded as a generalization of *D*-deals of meet-lattices [Bruns and Lakser (1970)].

3.1. DEFINITION. Let  $(S, \cdot, \leq)$  be a posemigroup,  $A \subseteq S$ . Then  $A$  is called *left (right) m-distributive* if  $\bigvee A$  exists and  $A$  satisfies the left (right) distributive law, that is,  $(\bigvee A) \cdot s = \bigvee (A \cdot s)$ ,  $(s \cdot (\bigvee A) = \bigvee (s \cdot A))$  for all  $s \in S$ .  $A$  is called *m-distributive* if  $A$  is both left *m-distributive* and right *m-distributive*.

3.2. REMARK. Let  $(S, \cdot, \leq)$  be a posemigroup. Then

1. If  $A$  is a left (right) *m-distributive* subset of  $S$ , then  $A \cdot s$  ( $s \cdot A$ ) is left (right) *m-distributive* for all  $s \in S$ .
2. If  $A$  is a *m-distributive* subset of a posemigroup  $S$ , then  $A \cdot s$  and  $s \cdot A$  are not necessarily *m-distributive* for all  $s \in S$ .

3.3. DEFINITION. Let  $A$  be a *m-distributive* subset of a posemigroup  $S$ . Then  $A$  is called *strong m-distributive (sm-distributive for short)* provided that  $A \cdot s$  is right *m-distributive* for all  $s \in S$ .

3.4. LEMMA. Let  $A$  be a *m-distributive* subset of a posemigroup  $S$ . Then  $A$  is *sm-distributive* if and only if  $s \cdot A$  is left *m-distributive* for all  $s \in S$ .

PROOF. Let  $A$  be *sm-distributive*. Then for all  $x, s \in S$ , we have that  $(\bigvee (s \cdot A)) \cdot x = (s \cdot (\bigvee A)) \cdot x = s \cdot ((\bigvee A) \cdot x) = s \cdot (\bigvee (A \cdot x)) = \bigvee (s \cdot (A \cdot x)) = \bigvee ((s \cdot A) \cdot x)$ , which implies that  $s \cdot A$  is left *m-distributive*. By symmetry, we can prove the inverse. ■

3.5. REMARK. Let  $(S, \cdot, \leq)$  be a posemigroup. Then

1.  $\{x\}, x \downarrow$  are *sm-distributive* for all  $x \in S$ .
2. If  $X$  is *sm-distributive*, then  $X \cdot y$  and  $y \cdot X$  are also *sm-distributive* for all  $y \in S$ .
3. If  $(S, \leq)$  is a complete lattice, then every subset of  $S$  is *m-distributive* if and only if  $S$  is a quantale. Furthermore, every subset of a quantale is *sm-distributive*.

3.6. DEFINITION. Let  $(S, \cdot, \leq)$  be a posemigroup. A lower set  $X$  of  $S$  is called a *sm-ideal* if it is closed under *sm-distributive* joins, that is, for any subset  $D$  of  $X$ ,  $\bigvee D \in X$  whenever  $D$  is *sm-distributive* in  $(S, \cdot, \leq)$ . We denote by  $\mathcal{R}(S)$  the set of all *sm-ideals* of  $S$ .

3.7. REMARK. Let  $(S, \cdot, \leq)$  be a posemigroup. Then

1.  $S \in \mathcal{R}(S)$ .
2. For all  $x \in S$ ,  $x \downarrow \in \mathcal{R}(S)$ .
3. If  $(S, \cdot, \leq)$  is a meet-semilattice and  $\cdot = \wedge$ , then  $A$  is a  $D$ -ideal of  $S$  if and only if  $S$  is a  $sm$ -ideal of  $S$ .
4. If  $(S, \cdot, \leq)$  is a quantale, then for any  $A \in \mathcal{R}(S)$ , there exists an element  $s \in S$  such that  $A = s \downarrow$ .

3.8. LEMMA. [Rosenthal (1990)] *If  $Q$  is a quantale and  $S \subseteq Q$ , then  $S$  is a quantic quotient if and only if  $S$  is closed under infs, and  $a \rightarrow s, s \leftarrow a \in S$ , whenever  $a \in Q$  and  $s \in S$ .*

3.9. PROPOSITION. *Let  $(S, \cdot, \leq)$  be a posemigroup. Then  $\mathcal{R}(S)$  is a quantic quotient of the power-set quantale  $(\mathcal{P}(S), \subseteq, \bullet)$ .*

PROOF. It is easy to show that  $\mathcal{R}(S)$  is closed under arbitrary meets, that is,  $\bigcap_{i \in I} A_i \in \mathcal{R}(S)$  for all  $A_i \in \mathcal{R}(S)$ . Let  $C \in \mathcal{R}(S)$ ,  $A \in \mathcal{P}(S)$ . Then we have that  $C \leftarrow A$  and  $A \rightarrow C$  are in  $\mathcal{R}(S)$ . Indeed: (1) Let  $y \leq x \in C \leftarrow A$ . Then  $x \cdot a \in C$  for all  $a \in A$ . Since  $C$  is a lower set, one can see that  $y \cdot a \leq x \cdot a \in C$  for all  $a \in A$ , which implies that  $y \in C \leftarrow A$ , that is,  $C \leftarrow A$  is a lower set of  $S$ . (2) Let  $D \subseteq C \leftarrow A$  be a  $sm$ -distributive subset. Then  $D \bullet A \subseteq C$ , which implies that  $D \cdot a \subseteq C$  for all  $a \in A$ . Since  $D \cdot a$  is  $sm$ -distributive for all  $a \in A$ , we have  $\vee(D \cdot a) \in C$ , that is,  $(\vee D) \cdot a \in C$  for all  $a \in A$ . Thus,  $\vee D \in C \leftarrow A$ . By (1) and (2), we have  $C \leftarrow A \in \mathcal{R}(S)$ . Similarly, we have  $A \rightarrow C \in \mathcal{R}(S)$ . Therefore, by Lemma 3.8 we have that  $\mathcal{R}(S)$  is a quantic quotient of  $\mathcal{P}(S)$ . ■

3.10. REMARK. Let  $(S, \cdot, \leq)$  be a posemigroup. Then there exists a quantic nucleus  $\rho_R$  such that  $\mathcal{R}(S) = (\mathcal{P}(S))_{\rho_R}$  and  $\rho_R(\{x\}) = x \downarrow$ , that is,  $\rho_R$  is a topological closure on  $\mathcal{P}(S)$ .

By Corollary 2.4 and Remark 3.10, we have the following theorem.

3.11. THEOREM. *Let  $(S, \cdot, \leq)$  be a posemigroup. Then  $(\eta_S, \mathcal{R}(S))$  is a quantale completion of  $S$ , where  $\eta_S(s) = \rho_R(\{s\})$ .*

3.12. REMARK. If  $(S, \cdot, \leq)$  be a quantale, then  $\eta_S$  is a quantale isomorphism.

By Theorem 3.11, we see that  $(\eta_S, \mathcal{R}(S))$  is a quantale completion of a posemigroup  $S$ . According to the description in Section 2, we have that  $\mathbf{S}^* \subseteq \mathcal{R}(S) \subseteq (\mathcal{P}(S))_{\mathcal{Q}}$ . The following example will indicate that  $\mathcal{R}(S)$  lies strictly between  $\mathbf{S}^*$  and  $(\mathcal{P}(S))_{\mathcal{Q}}$ .

3.13. EXAMPLE. Let  $S$  be a posemigroup with the partial order  $\leq$  determined by Figure 3.1 and the binary multiplication  $\cdot$  defined by (3.1)




Figure 3.1

$$\forall x, y \in S, x \cdot y = \begin{cases} 0, & \text{if } x = 0 \text{ or } y = 0, \\ d, & \text{otherwise.} \end{cases} \tag{3.1}$$

One can easily verify that  $M$  is a  $sm$ -distributive subset of  $S$  if and only if  $\vee M$  exists in  $S$ . Therefore, we have

$$S^* = \{\{0\}, \{a, 0\}, \{b, 0\}, \{c, 0\}, \{a, b, d, 0\}, S\},$$

$$\mathcal{R}(S) = \{\{0\}, \{a, 0\}, \{b, 0\}, \{c, 0\}, \{a, c, 0\}, \{b, c, 0\}, \{a, b, d, 0\}, S\},$$

$$(\mathcal{P}(S))_{\mathcal{D}} = \{\emptyset, \{0\}, \{a, 0\}, \{b, 0\}, \{c, 0\}, \{a, b, 0\}, \{a, c, 0\}, \{b, c, 0\}, \{a, b, c, 0\}, \{a, b, d, 0\}, S\}.$$

#### 4. Free objects over posemigroups in the category $\mathbf{PoSgr}_\vee$

In this section, we shall show that  $\mathcal{R}(S)$  is in fact a free object over  $S$  in the category  $\mathbf{PoSgr}_\vee$  of posemigroups with  $sm$ -distributive join homomorphisms. First, we shall introduce the concept of  $sm$ -distributive join homomorphisms.

4.1. DEFINITION. Let  $(S, \cdot, \leq)$  and  $(T, *, \leq)$  be posemigroups. Then a map  $f: S \rightarrow T$  is called a  $sm$ -distributive join homomorphism if for any  $sm$ -distributive subset  $X$  of  $S$ ,  $f(X)$  is  $sm$ -distributive with  $f(\vee X) = \vee f(X)$  and  $f$  is a semigroup homomorphism.

Let  $\mathbf{PoSgr}_\vee$  the category of posemigroups with  $sm$ -distributive join homomorphisms. By Remark 3.5(3), we have that the  $sm$ -distributive join homomorphisms between quantales are exactly the quantale homomorphisms, which implies that the category  $\mathbf{Quant}$  of quantales with quantale homomorphisms is a full subcategory of  $\mathbf{PoSgr}_\vee$ .

4.2. LEMMA. Let  $(S, \cdot, \leq)$  be a posemigroup. Then for all  $a \in S, X \in \mathcal{R}(S)$ , we have that  $(-\cdot a)^{-1}(X), (a \cdot -)^{-1}(X) \in \mathcal{R}(S)$ .

PROOF. The proof is straightforward. ■

In the following, we shall give a characterization for  $sm$ -distributive join homomorphisms between posemigroups.

4.3. LEMMA. Let  $f: (S, \leq, \cdot) \rightarrow (T, \leq, *)$  be a semigroup homomorphism between posemigroups. Then  $f$  is a  $sm$ -distributive join homomorphism if and only if  $f^{-1}(X) \in \mathcal{R}(S)$  for all  $X \in \mathcal{R}(T)$ .

PROOF. Assume that  $f$  is a  $sm$ -distributive join homomorphism and  $X$  is a  $sm$ -ideal of  $T$ , that is,  $X \in \mathcal{R}(T)$ . One can easily prove that  $f$  is order-preserving, which implies that  $f^{-1}(X)$  is a lower set of  $S$ . For any  $sm$ -distributive subset  $A$  of  $f^{-1}(X)$ , we have that  $f(A) \subseteq X$  and  $f(\bigvee A) = \bigvee f(A)$ . Since  $X$  is a  $sm$ -ideal and  $f(A)$  is a  $sm$ -distributive subset of  $X$ , we have  $f(\bigvee A) \in X$ , which implies  $\bigvee A \in f^{-1}(X)$ . Thus,  $f^{-1}(X)$  is a  $sm$ -ideal of  $S$ , that is,  $f^{-1}(X) \in \mathcal{R}(S)$ .

Conversely, it suffices to prove that  $f(X)$  is  $sm$ -distributive with  $f(\bigvee X) = \bigvee f(X)$  for any  $sm$ -distributive set  $X$  of  $S$ .

(1)  $f$  is order-preserving. Let  $x, y \in S$  with  $x \leq y$ . Then from Remark 3.7 it follows that  $f^{-1}(f(y) \downarrow) \in \mathcal{R}(S)$ . Since  $y \in f^{-1}(f(y) \downarrow)$ , we have  $x \in f^{-1}(f(y) \downarrow)$ , which implies that  $f(x) \leq f(y)$ .

(2)  $f(\bigvee X) = \bigvee f(X)$ . Obviously,  $f(\bigvee X)$  is an upper bound of  $f(X)$ . Assume that  $t \in T$  is an arbitrary upper bound of  $f(X)$ , then  $X \subseteq f^{-1}(t \downarrow)$ . Since  $f^{-1}(t \downarrow)$  is a  $sm$ -ideal and  $\bigvee X \in f^{-1}(t \downarrow)$ , we have  $f(\bigvee X) \leq t$ . Thus,  $f(\bigvee X) = \bigvee f(X)$ .

(3)  $f(X)$  is  $sm$ -distributive. For any  $x \in T$ ,  $x * (\bigvee f(X))$  is clearly an upper bound of  $x * f(X)$ . Assume that  $t \in T$  is an arbitrary upper bound of  $x * f(X)$ , then  $x * f(X) \subseteq t \downarrow$ . From Lemma 4.2, it follows that  $f(X) \subseteq (x * -)^{-1}(t \downarrow) \in \mathcal{R}(T)$ . By the assumption of  $f$ , we have that  $X \subseteq f^{-1}((x * -)^{-1}(t \downarrow))$  and  $\bigvee X \in f^{-1}((x * -)^{-1}(t \downarrow))$ , which implies that  $x * f(\bigvee X) \leq t$ . Thus,  $x * (\bigvee f(X)) = \bigvee (x * f(X))$ . Similarly,  $(\bigvee f(X)) * x = \bigvee (f(X) * x)$ . So, we have that  $f(X)$  is  $m$ -distributive. In the following we shall prove that for any  $x \in T$ ,  $f(X) * x$  is right  $sm$ -distributive. For any  $y \in T$ ,  $y * (\bigvee (f(X) * x))$  is clearly an upper bound of  $y * (f(X) * x)$ . Suppose that  $t \in S$  is an arbitrary upper bound of  $y * (f(X) * x)$ . Then we have  $y * (f(X) * x) \subseteq t \downarrow$  and  $X \subseteq f^{-1}((y * -)^{-1}((- * x)^{-1}(t \downarrow)))$ . Since  $X$  is  $sm$ -distributive, by Lemma 4.2 and the assumption of  $f$  we have that  $\bigvee X \in f^{-1}((y * -)^{-1}((- * x)^{-1}(t \downarrow)))$  and  $(y * f(\bigvee X)) * x \leq t$ , which implies  $\bigvee (y * (f(X) * x)) = y * (\bigvee (f(X) * x))$ . Thus,  $f(X)$  is  $sm$ -distributive. ■

4.4. LEMMA. Let  $f: (S, \cdot, \leq) \rightarrow (T, *, \leq)$  be a  $sm$ -distributive join homomorphism between posemigroups. Then

1.  $\rho_R(f(X)) = \rho_R(f(\rho_R(X)))$  for all  $X \in \mathcal{P}(S)$ .
2.  $F_f: \mathcal{R}(S) \rightarrow \mathcal{R}(T)$  is a quantale homomorphism, where  $F_f(X) = \rho_R(f(X))$ .

PROOF. (1) It is obvious to see that  $\rho_R(f(X)) \subseteq \rho_R(f(\rho_R(X)))$ . From Lemma 4.3, it follows that  $f(X) \subseteq \rho_R(f(X)) \implies X \subseteq f^{-1}(\rho_R(f(X))) \implies \rho_R(X) \subseteq f^{-1}(\rho_R(f(X))) \implies f(\rho_R(X)) \subseteq \rho_R(f(X)) \implies \rho_R(f(\rho_R(X))) \subseteq \rho_R(f(X))$ . Thus,  $\rho_R(f(X)) = \rho_R(f(\rho_R(X)))$ .

(2) Define a map  $G: \mathcal{R}(T) \rightarrow \mathcal{R}(S)$  as follows  $G(Y) = f^{-1}(Y)$ . From Lemma 4.3, it follows that  $G$  is well defined. Since  $F_f(X) \subseteq Y \Leftrightarrow \rho_R(f(X)) \subseteq Y \Leftrightarrow f(X) \subseteq Y \Leftrightarrow X \subseteq f^{-1}(Y) \Leftrightarrow X \subseteq G(Y)$  for all  $X \in \mathcal{R}(S), Y \in \mathcal{R}(T)$ , we have that  $F_f$  is the left adjoint of  $G$ , which implies that  $F_f$  preserves arbitrary joins. For all  $A, B \in \mathcal{R}(S)$ , by (1) we have  $F_f(A \otimes B) = \rho_R(f(A \otimes B)) = \rho_R(f(\rho_R(A \bullet B))) = \rho_R(f(A \bullet B)) = \rho_R(f(A) \bullet f(B)) = \rho_R(f(A)) \otimes \rho_R(f(B)) = F_f(A) \otimes F_f(B)$ . Thus,  $F_f$  is a quantale homomorphism. ■

4.5. THEOREM.  $\mathcal{R}(S)$  is a free object over a posemigroup  $S$  in the category  $\mathbf{Posgr}_\vee$ .

PROOF. As we have seen above, **Quant** is a full subcategory of  $\mathbf{Posgr}_\vee$ . Let  $S$  be a posemigroup. Then by Proposition 3.9 we see that  $\mathcal{R}(S)$  is a quantale. Further, by Theorem 3.11 we have that  $\eta_S : S \rightarrow \mathcal{R}(S)$  is a semigroup homomorphism which is order-preserving. By Remark 3.5(3) and Remark 3.7, we have that for all  $A \in \mathcal{P}(S)$ ,  $\eta_S(A)$  is *sm*-distributive in  $\mathcal{R}(S)$ . For any *sm*-distributive subset  $A$  of  $S$ , we claim that  $\eta_S(\bigvee A) = \bigvee_{a \in A} \eta_S(a)$ . It is easy to see that  $\eta_S(a) \subseteq \eta_S(\bigvee A)$  for all  $a \in A$ . Let  $B \in \mathcal{R}(S)$  with  $\eta_S(a) \subseteq B$  for all  $a \in A$ . Then  $A \subseteq B$ . Since  $B \in \mathcal{R}(S)$  and  $A$  is *sm*-distributive, we have that  $\bigvee A \in B$  and  $\eta_S(\bigvee A) \subseteq B$ , which implies  $\eta_S(\bigvee A) = \bigvee_{a \in A} \eta_S(a)$ . Thus,  $\eta_S$  is a *sm*-distributive join homomorphism. In what follows we shall prove that  $\eta_S : S \rightarrow \mathcal{R}(S)$  has the universal property.

Let  $f : S \rightarrow Q$  be a *sm*-distributive join homomorphism into a quantale  $Q$ . We let  $\bar{f} = (\eta_Q)^{-1} \circ F_f$ . First, we shall prove  $\bar{f} \circ \eta_S = f$ . By Remark 3.12 and Lemma 4.4, we have that  $\bar{f}$  is a quantale homomorphism. For all  $x \in S$ , we have  $\bar{f} \circ \eta_S(x) = (\eta_Q)^{-1} \circ F_f(x \downarrow) = (\eta_Q)^{-1}(f(x) \downarrow) = f(x)$ . Next, we shall prove the uniqueness of  $\bar{f}$ . Suppose that  $g : \mathcal{R}(S) \rightarrow Q$  is another quantale homomorphism such that  $g \circ \eta_S = f$ , then  $g(x \downarrow) = \bar{f}(x \downarrow)$  for any  $x \in S$ . By Lemma 4.4(2), we have that for any  $X \in \mathcal{R}(S)$ ,  $\bar{f}(X) = \bar{f}(\bigvee_{x \in X} x \downarrow) = \bigvee_{x \in X} \bar{f}(x \downarrow) = \bigvee_{x \in X} g(x \downarrow) = g(\bigvee_{x \in X} x \downarrow) = g(X)$ , which implies that  $g = \bar{f}$ . ■

4.6. COROLLARY. **Quant** is a full reflective subcategory of  $\mathbf{Posgr}_\vee$ .

## 5. Acknowledgement

We would like to thank the referee for some of his or her helpful comments and suggestions for the improvement of this paper.

## References

- J. Adámek, H. Herrlich and G.E. Strecker (2004), *Abstract and Concrete Categories: The joy of cats*, Wiley, New York.
- G. Bruns, H. Lakser (1970), Injective hulls of semilattices. *Canad. Math. Bull.* **13**, 115–118.
- B. Eckmann and A. Schopf (1953), Über injektive Moduln. *Archiv der Mathematik* **4**, 75–78.
- S.W. Han and B. Zhao (2008), Quantale completion of ordered semigroups. *Atac Math. Sinica Chin. Ser.* **51**, 1081–1088.
- D. Kruml and J. Paseka (2008), Algebraic and categorical aspects of quantales. *Handbook of algebra*, **5**, 1–43.

- J. Lambek, M. Barr, J.F. Kennison and R. Raphael (2012), Injective hulls of partially ordered monoids. *Theory Appl. Categ.* **26**, 338–348.
- D. Larchey-Wendling and D. Galmiche (2000), Quantales as completions of ordered monoids: Revised semantics for Intuitionistic Linear Logic. *Electron. Notes Theor. Comput. Sci.* **35**, 94–108.
- K.I. Rosenthal (1990), *Quantales and Their Applications*. Longman Scientific and Technical, New York.
- C.C. Xia, B. Zhao and S.W. Han (2016), Some further results of quantale completions of ordered semigroup. Submitted to *Acta Mathematica Sinica, Chinese Series*.
- C.C. Xia, S.W. Han and B. Zhao (2017), A note on injective hulls of posemigroups. *Theory Appl. Categ.* **7**, 254–257.
- X. Zhang, V. Laan (2014), Injective hulls for posemigroups. *Proc. Est. Acad. Sci.* **63**, 372–378.

*Department of Mathematics, Shaanxi Normal University,  
Xi'an 710119, China*

Email: hansw@snnu.edu.cn  
xiachangchun@snnu.edu.cn  
gxj\_2886@163.com

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available from the journal's server at <http://www.tac.mta.ca/tac/>. It is archived electronically and in printed paper format.

**SUBSCRIPTION INFORMATION** Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to [tac@mta.ca](mailto:tac@mta.ca) including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca).

**INFORMATION FOR AUTHORS** The typesetting language of the journal is  $\text{T}_{\text{E}}\text{X}$ , and  $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\text{e}$  is required. Articles in PDF format may be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

**MANAGING EDITOR.** Robert Rosebrugh, Mount Allison University: [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca)

**T<sub>E</sub>XNICAL EDITOR.** Michael Barr, McGill University: [barr@math.mcgill.ca](mailto:barr@math.mcgill.ca)

**ASSISTANT T<sub>E</sub>X EDITOR.** Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: [gavin\\_seal@fastmail.fm](mailto:gavin_seal@fastmail.fm)

**TRANSMITTING EDITORS.**

Clemens Berger, Université de Nice-Sophia Antipolis: [cberger@math.unice.fr](mailto:cberger@math.unice.fr)

Richard Blute, Université d' Ottawa: [rblute@uottawa.ca](mailto:rblute@uottawa.ca)

Lawrence Breen, Université de Paris 13: [breen@math.univ-paris13.fr](mailto:breen@math.univ-paris13.fr)

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva, Nuance Communications Inc: [valeria.depaiva@gmail.com](mailto:valeria.depaiva@gmail.com)

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: [kathryn.hess@epfl.ch](mailto:kathryn.hess@epfl.ch)

Martin Hyland, University of Cambridge: [M.Hyland@dpms.cam.ac.uk](mailto:M.Hyland@dpms.cam.ac.uk)

Anders Kock, University of Aarhus: [kock@imf.au.dk](mailto:kock@imf.au.dk)

Stephen Lack, Macquarie University: [steve.lack@mq.edu.au](mailto:steve.lack@mq.edu.au)

F. William Lawvere, State University of New York at Buffalo: [wlawvere@buffalo.edu](mailto:wlawvere@buffalo.edu)

Tom Leinster, University of Edinburgh: [Tom.Leinster@ed.ac.uk](mailto:Tom.Leinster@ed.ac.uk)

Ieke Moerdijk, Utrecht University: [i.moerdijk@uu.nl](mailto:i.moerdijk@uu.nl)

Susan Niefield, Union College: [niefiels@union.edu](mailto:niefiels@union.edu)

Robert Paré, Dalhousie University: [pare@mathstat.dal.ca](mailto:pare@mathstat.dal.ca)

Jiri Rosicky, Masaryk University: [rosicky@math.muni.cz](mailto:rosicky@math.muni.cz)

Giuseppe Rosolini, Università di Genova: [rosolini@disi.unige.it](mailto:rosolini@disi.unige.it)

Alex Simpson, University of Ljubljana: [Alex.Simpson@fmf.uni-lj.si](mailto:Alex.Simpson@fmf.uni-lj.si)

James Stasheff, University of North Carolina: [jds@math.upenn.edu](mailto:jds@math.upenn.edu)

Ross Street, Macquarie University: [ross.street@mq.edu.au](mailto:ross.street@mq.edu.au)

Walter Tholen, York University: [tholen@mathstat.yorku.ca](mailto:tholen@mathstat.yorku.ca)

Myles Tierney, Université du Québec à Montréal: [tierney.myles4@gmail.com](mailto:tierney.myles4@gmail.com)

R. J. Wood, Dalhousie University: [rjwood@mathstat.dal.ca](mailto:rjwood@mathstat.dal.ca)