

ON SOME PROPERTIES OF THE PICARD OPERATORS

LUCYNA REMPULSKA AND KAROLINA TOMCZAK

ABSTRACT. We consider the Picard operators \mathcal{P}_n and $\mathcal{P}_{n;r}$ in exponential weighted spaces. We give some elementary and approximation properties of these operators.

1. INTRODUCTION

1.1. The Picard operators

$$(1) \quad \mathcal{P}_n(f; x) := \frac{n}{2} \int_{\mathbb{R}} f(x-t)e^{-n|t|} dt = \frac{n}{2} \int_{\mathbb{R}} f(x+t)e^{-n|t|} dt,$$

$x \in \mathbb{R}$ and $n \in \mathbb{N}$, ($\mathbb{N} = \{1, 2, \dots\}$, $\mathbb{R} = (-\infty, +\infty)$) are investigated for functions $f: \mathbb{R} \rightarrow \mathbb{R}$ from various classes in many monographs and papers (e.g. [2]–[8] [10, 11]).

G. H. Kirov in the paper [9] introduced the generalized Bernstein polynomials $\mathcal{B}_{n;r}$ for r -times differentiable functions $f \in C^r([0, 1])$ and he showed that $\mathcal{B}_{n;r}$ have better approximation properties than classical Bernstein polynomials \mathcal{B}_n .

The Kirov method was used in [12] to the generalized Picard operators

$$(2) \quad \mathcal{P}_{n;r}(f; x) := \mathcal{P}_n(F_r(t, x); x), \quad x \in \mathbb{R}, n \in \mathbb{N}, r \in \mathbb{N}_0,$$

$$(3) \quad F_r(t, x) := \sum_{j=0}^r \frac{f^{(j)}(t)}{j!} (x-t)^j,$$

($\mathbb{N}_0 = \mathbb{N} \cup \{0\}$) of r -times differentiable functions $f: \mathbb{R} \rightarrow \mathbb{R}$. Obviously $\mathcal{P}_{n;0}(f) \equiv \mathcal{P}_n(f)$.

In this paper we examine the Picard operators \mathcal{P}_n (in Section 2) and $\mathcal{P}_{n;r}$ (in Section 3) for functions f belonging to the exponential weighted spaces $L_q^p(\mathbb{R})$ and $L_q^{p;r}(\mathbb{R})$ which definition is given below. We present some elementary properties, the orders of approximation and the Voronovskaya – type theorems for these operators.

2000 *Mathematics Subject Classification*: primary 41A25.

Key words and phrases: Picard operator, exponential weighted space, degree of approximation, Voronovskaya type theorem.

Received August 27, 2008, revised March 2009. Editor O. Došlý.

1.2. Let $q > 0$ and $1 \leq p \leq \infty$ be fixed,

$$(4) \quad v_q(x) := e^{-q|x|} \quad \text{for } x \in \mathbb{R},$$

and let $L_q^p \equiv L_q^p(\mathbb{R})$ be the space of all functions $f: \mathbb{R} \rightarrow \mathbb{R}$ for which $v_q f$ is Lebesgue integrable with p -th power over \mathbb{R} if $1 \leq p < \infty$ and uniformly continuous and bounded on \mathbb{R} if $p = \infty$. The norm in L_q^p is defined by

$$(5) \quad \|f\|_{p,q} \equiv \|f(\cdot)\|_{p,q} := \begin{cases} \left(\int_{\mathbb{R}} |v_q(x)f(x)|^p dx \right)^{1/p} & \text{if } 1 \leq p < \infty, \\ \sup_{x \in \mathbb{R}} v_q(x)|f(x)| & \text{if } p = \infty. \end{cases}$$

Moreover, let $r \in \mathbb{N}_0$ and $L_q^{p,r} \equiv L_q^{p,r}(\mathbb{R})$ be the class of all r -times differentiable functions $f \in L_q^p$ having the derivatives $f^{(k)} \in L_q^p$, $1 \leq k \leq r$. The norm in $L_q^{p,r}$ is given by (5). ($L_q^{p,0} \equiv L_q^p$). The spaces L_q^p and $L_q^{p,r}$ are called exponential weighted spaces ([1]).

As usual, for $f \in L_q^p$ and $k \in \mathbb{N}$ we define the k -th modulus of smoothness:

$$(6) \quad \omega_k(f; L_q^p; t) := \sup_{|h| \leq t} \|\Delta_h^k f(\cdot)\|_{p,q} \quad \text{for } t \geq 0,$$

$$(7) \quad \Delta_h^k f(x) := \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} f(x + jh).$$

The above ω_k has the following properties:

$$(8) \quad \omega_k(f; L_q^p; t_1) \leq \omega_k(f; L_q^p; t_2) \quad \text{for } 0 \leq t_1 < t_2,$$

$$(9) \quad \omega_k(f; L_q^p; \lambda t) \leq (1 + \lambda)^k e^{kq\lambda t} \omega_k(f; L_q^p; t) \quad \text{for } \lambda, t \geq 0,$$

$$(10) \quad \lim_{t \rightarrow 0^+} \omega_k(f; L_q^p; t) = 0,$$

for every $f \in L_q^p$ and $k \in \mathbb{N}$ (see [6, Chapter 6] and [13, Chapter 3]).

By ω_k we define the Lipschitz class

$$(11) \quad \text{Lip}_M^k(L_q^p; \alpha) := \{f \in L_q^p : \omega_k(f; L_q^p; t) \leq Mt^\alpha \quad \text{for } t \geq 0\}$$

for fixed numbers: $1 \leq p \leq \infty$, $q > 0$, $k \in \mathbb{N}$, $M > 0$ and $0 < \alpha \leq k$.

2. SOME PROPERTIES OF \mathcal{P}_n

2.1. By elementary calculations can be obtained the following two lemmas.

Lemma 1. *The equality*

$$(12) \quad \int_0^\infty t^r e^{-st} dt = \frac{r!}{s^{r+1}}$$

there holds for every $r \in \mathbb{N}_0$ and $s > 0$.

Lemma 2. *Let $e_0(x) = 1$, $e_1(x) = x$ and let $\varphi_x(t) = t - x$ for $x, t \in \mathbb{R}$. Then*

$$(13) \quad \mathcal{P}_n(e_i; x) = e_i(x) \quad \text{for } x \in \mathbb{R}, n \in \mathbb{N}, i = 0, 1,$$

and

$$(14) \quad \mathcal{P}_n (\varphi_x^k(t); x) = \frac{(1 + (-1)^k) k!}{2n^k},$$

$$(15) \quad \mathcal{P}_n \left(|\varphi_x(t)|^k \exp (q |\varphi_x(t)|); x \right) = \frac{k!n}{(n - q)^{k+1}},$$

for $x \in \mathbb{R}$, $n \geq q + 1$ and $k \in \mathbb{N}_0$.

Using the above results and arguing analogously to the proof of Lemma 2 in [10] we can obtain the following basic lemma.

Lemma 3. *Let $f \in L_q^p$ with fixed $1 \leq p \leq \infty$ and $q > 0$. Then*

$$(16) \quad \|\mathcal{P}_n(f)\|_{p,q} \leq (1 + q)\|f\|_{p,q} \quad \text{for } n \geq q + 1.$$

The formula (1) and (16) show that \mathcal{P}_n , $n \geq q + 1$, is a positive linear operator acting from the space L_q^p to L_q^p .

2.2. By (6), (7), (11) and (16) can be derived the following geometric properties of \mathcal{P}_n given by (1).

Theorem 1. *Let $f \in L_q^p$ with fixed $1 \leq p \leq \infty$ and $q > 0$ and let $q + 1 \leq n \in \mathbb{N}$. Then*

- (i) *if f is non-decreasing (non-increasing) on \mathbb{R} , then $\mathcal{P}_n(f)$ is also non-decreasing (non-increasing) on \mathbb{R} ,*
- (ii) *if f is convex (concave) on \mathbb{R} , then $\mathcal{P}_n(f)$ is also convex (concave) on \mathbb{R} ,*
- (iii) *for every $k \in \mathbb{N}$ there holds the inequality*

$$\omega_k (\mathcal{P}_n(f); L_q^p; t) \leq (1 + q)\omega_k (f; L_q^p; t), \quad t \geq 0,$$

- (iv) *if $f \in \text{Lip}_M^k (L_q^p; \alpha)$ with fixed $k \in \mathbb{N}$, $0 < \alpha \leq k$ and $M > 0$, then also $\mathcal{P}_n(f) \in \text{Lip}_{M^*}^k (L_q^p; \alpha)$ with the same k and α and $M^* = (1 + q)M$,*
- (v) *If $f \in L_q^{\infty,r}$ with a fixed $r \in \mathbb{N}$, then $\mathcal{P}_n(f) \in L_q^{\infty,r}$ and for derivatives of $\mathcal{P}_n(f)$ there holds*

$$\|\mathcal{P}_n^{(k)}(f)\|_{\infty,q} = \|\mathcal{P}_n(f^{(k)})\|_{\infty,q} \leq (1 + q)\|f^{(k)}\|_{\infty,q}$$

Proof. For example we prove (iii). From the formulas (1) and (7) there results that

$$\Delta_h^k \mathcal{P}_n(f; x) = \mathcal{P}_n (\Delta_h^k f; x) \quad \text{for } x, h \in \mathbb{R}, k \in \mathbb{N}.$$

Next, by (5) and (16), we have

$$\|\Delta_h^k \mathcal{P}_n(f; \cdot)\|_{p,q} = \|\mathcal{P}_n (\Delta_h^k f, \cdot)\|_{p,q} \leq (1 + q) \|\Delta_h^k f(\cdot)\|_{p,q}$$

for $h \in \mathbb{R}$ and $n \geq q + 1$, and using (6), we get the statement (iii). □

2.3. Arguing similarly to [5] and [10] and applying (6)–(9), (12) and (16) we can prove the following approximation theorem.

Theorem 2. *Suppose that $f \in L_q^p$ with fixed $1 \leq p \leq \infty$ and $q > 0$. Then*

$$\|\mathcal{P}_n(f) - f\|_{p,q} \leq \frac{5}{2}(1 + 3q)^3 \omega_2\left(f; L_q^p; \frac{1}{n}\right)$$

for every $n \geq 3q + 1$.

From Theorem 2 and (8), (10) and (11) there results the following

Corollary 1. *If $f \in L_q^p$, $1 \leq p \leq \infty$, $q > 0$, then*

$$(17) \quad \lim_{n \rightarrow \infty} \|\mathcal{P}_n(f) - f\|_{p,q} = 0.$$

In particular, if $f \in \text{Lip}_M^2(L_q^p; \alpha)$ with fixed $0 < \alpha \leq 2$ and $M > 0$, then

$$\|\mathcal{P}_n(f) - f\|_{p,q} = O(n^{-\alpha}) \quad \text{as } n \rightarrow \infty.$$

Applying Corollary 1, we shall prove the Voronovskaya-type theorem for \mathcal{P}_n .

Theorem 3. *Let $f \in L_q^{\infty,2}$ with a fixed $q > 0$. Then*

$$(18) \quad \lim_{n \rightarrow \infty} n^2 [\mathcal{P}_n(f; x) - f(x)] = f''(x)$$

for every $x \in \mathbb{R}$.

Proof. Choose $f \in L_q^{\infty,2}$ and $x \in \mathbb{R}$. Then, by the Taylor formula, we have

$$f(t) = f(x) + f'(x)(t-x) + \frac{1}{2}f''(x)(t-x)^2 + \psi(t; x)(t-x)^2 \quad \text{for } t \in \mathbb{R},$$

where $\psi(t) \equiv \psi(t, x)$ is a function belonging to L_q^∞ and $\lim_{t \rightarrow x} \psi(t; x) = \psi(x) = 0$. Using operator \mathcal{P}_n , $n \geq 2q + 1$, and (13) and (14), we get

$$(19) \quad \mathcal{P}_n(f(t); x) = f(x) + n^{-2}f''(x) + \mathcal{P}_n(\psi(t)\varphi_x^2(t); x)$$

and by the Hölder inequality and (14):

$$\begin{aligned} |\mathcal{P}_n(\psi(t)\varphi_x^2(t); x)| &\leq (\mathcal{P}_n(\psi^2(t); x) \mathcal{P}_n(\varphi_x^4(t); x))^{1/2} \\ &= n^{-2} (24\mathcal{P}_n(\psi^2(t); x))^{1/2}. \end{aligned}$$

From properties of ψ and (17) there results that $\lim_{n \rightarrow \infty} \mathcal{P}_n(\psi^2(t); x) = \psi^2(x) = 0$. Consequently,

$$(20) \quad \lim_{n \rightarrow \infty} n^2 \mathcal{P}_n(\psi(t)\varphi_x^2(t); x) = 0$$

and by (19) and (20) follows (18). □

Now we estimate the rate of convergence given by (18).

Theorem 4. *Let $f \in L_q^{\infty,2}$ with a fixed $q > 0$. Then*

$$(21) \quad \|n^2 [\mathcal{P}_n(f) - f] - f''\|_{\infty,q} \leq 4(1 + q)^4 \omega_1\left(f''; L_q^\infty; \frac{1}{n}\right)$$

for $n \geq q + 1$.

Proof. For $f \in L_q^{\infty,2}$ and $x, t \in \mathbb{R}$ there holds the Taylor-type formula

$$f(t) = f(x) + f'(x)(t-x) + \frac{1}{2}f''(x)(t-x)^2 + (t-x)^2 I(t, x),$$

where

$$(22) \quad I(t, x) := \int_0^1 (1-u) [f''(x+u(t-x)) - f''(x)] du.$$

Using operator \mathcal{P}_n , $n \geq q+1$, and (13)-(15), we get

$$\mathcal{P}_n(f(t); x) = f(x) + n^{-2}f''(x) + \mathcal{P}_n(\varphi_x^2(t)I(t, x); x),$$

which implies that

$$(23) \quad |n^2[\mathcal{P}_n(f; x) - f(x)] - f''(x)| \leq n^2 \mathcal{P}_n(\varphi_x^2(t)|I(t, x)|; x),$$

for $x \in \mathbb{R}$ and $n \geq q+1$. Now, applying (6), (8) and (9), we get from (22):

$$\begin{aligned} |I(t, x)| &\leq \int_0^1 (1-u)\omega_1(f''; L_q^\infty; u|t-x|) e^{q|u|} du \\ &\leq \frac{1}{2}\omega_1(f''; L_q^\infty; |t-x|) e^{q|x|} \\ &\leq \frac{1}{2}\omega_1\left(f''; L_q^\infty; \frac{1}{n}\right) (1+n|t-x|) e^{q|x|+q|t-x|}. \end{aligned}$$

and next by (4) and (15) we can write

$$\begin{aligned} n^2 v_q(x) \mathcal{P}_n(\varphi_x^2(t)|I(t, x)|; x) &\leq \frac{n^2}{2}\omega_1\left(f''; L_q^\infty; \frac{1}{n}\right) \\ &\quad \times \left\{ \mathcal{P}_n((t-x)^2 e^{q|t-x|}; x) + n \mathcal{P}_n(|t-x|^3 e^{q|t-x|}; x) \right\} \\ &= \omega_1\left(f''; L_q^\infty; \frac{1}{n}\right) \left(\frac{n^3}{(n-q)^3} + \frac{3n^4}{(n-q)^4} \right) \\ &\leq 4(1+q)^4 \omega_1\left(f''; L_q^\infty; \frac{1}{n}\right) \quad \text{for } x \in \mathbb{R}, n \geq q+1. \end{aligned}$$

Now the estimate (21) is obvious by (23), the last inequality and (5). □

Theorem 5. Suppose that $f \in L_q^{\infty,r}$ with fixed $q > 0$ and $r \in \mathbb{N}$. Then

$$(24) \quad \|\mathcal{P}_n^{(r)}(f) - f^{(r)}\|_{\infty,q} \leq \frac{5}{2}(1+3q)^3 \omega_2\left(f^{(r)}; L_q^\infty; \frac{1}{n}\right)$$

for $n \geq 3q+1$.

Proof. If $f \in L_q^{\infty,r}$, then for the r -th derivative of $\mathcal{P}_n(f)$ we have by Theorem 1, (13) and (7):

$$\begin{aligned} \mathcal{P}_n^{(r)}(f; x) - f^{(r)}(x) &= \frac{n}{2} \int_{\mathbb{R}} [f^{(r)}(x+t) - f^{(r)}(x)] e^{-n|t|} dt \\ &= \frac{n}{2} \int_0^\infty [\Delta_t^2 f^{(r)}(x-t)] e^{-nt} dt. \end{aligned}$$

From this and by (6), (9) and (12) we deduce that

$$\begin{aligned} \|\mathcal{P}_n^{(r)}(f) - f^{(r)}\|_{\infty, q} &\leq \frac{n}{2} \int_0^\infty \omega_2\left(f^{(r)}; L_q^\infty; t\right) e^{-(n-q)t} dt \\ &\leq \omega_2\left(f^{(r)}; L_q^\infty; \frac{1}{n}\right) \frac{n}{2} \int_0^\infty (1+nt)^2 e^{-(n-3q)t} dt \\ &= \omega_2\left(f^{(r)}; L_q^\infty; \frac{1}{n}\right) \left\{ \frac{n}{2(n-3q)} + \frac{n^2}{(n-3q)^2} + \frac{n^3}{(n-3q)^3} \right\} \end{aligned}$$

for $n \geq 3q + 1$, which yields the estimate (24). \square

3. SOME PROPERTIES OF $\mathcal{P}_{n;r}$

3.1. The formulas (1)–(3) show that the operators $\mathcal{P}_{n;r}$, $r \in \mathbb{N}_0$, generalize \mathcal{P}_n and $\mathcal{P}_{n;0}(f) \equiv \mathcal{P}_n(f)$ for $f \in L_q^{p,0}$. By this fact and Section 1, we shall consider $\mathcal{P}_{n;r}$ for $r \in \mathbb{N}$ only.

Lemma 4. *Let $1 \leq p \leq \infty$, $q > 0$ and $k \in \mathbb{N}$ be fixed numbers. Then for every $f \in L_q^{p,r}$ and $n \geq q + 1$ there holds*

$$(25) \quad \|\mathcal{P}_{n;r}(f)\|_{p,q} \leq (1+q) \sum_{j=0}^r \|f^{(j)}\|_{p,q}.$$

The formulas (1)–(3) and the inequality (24) show that $\mathcal{P}_{n;r}$, $n \geq q + 1$, is a linear operator acting from $L_q^{p,r}$ to L_q^p .

Proof. Let $1 \leq p < \infty$. Then, by (1)–(3), the Minkowski inequality and (12), we get for $f \in L_q^{p,r}$ and $n \geq q + 1$:

$$\begin{aligned} \|\mathcal{P}_{n;r}(f)\|_{p,q} &\leq \sum_{j=0}^r \frac{1}{j!} \|\mathcal{P}_n\left(f^{(j)}(t)\varphi_x^j(t); \cdot\right)\|_{p,q} \\ &\leq \sum_{j=0}^r \frac{1}{j!} \left(\int_{\mathbb{R}} \left| e^{-q|x|} \frac{n}{2} \int_{\mathbb{R}} t^j f^{(j)}(x+t) e^{-n|t|} dt \right|^p dx \right)^{1/p} \\ &\leq \sum_{j=0}^r \frac{n}{2j!} \int_{\mathbb{R}} |t|^j e^{-n|t|} \left(\int_{\mathbb{R}} \left| e^{-q|x|} f^{(j)}(x+t) \right|^p dx \right)^{1/p} dt \\ &\leq \sum_{j=0}^r \frac{n}{2j!} \|f^{(j)}\|_{p,q} \int_{\mathbb{R}} |t|^j e^{-(n-q)|t|} dt \\ &= \sum_{j=0}^r \|f^{(j)}\|_{p,q} \frac{n}{(n-q)^{j+1}} \leq (1+q) \sum_{j=0}^r \|f^{(j)}\|_{p,q}. \end{aligned}$$

The proof of (25) for $p = \infty$ is similar. \square

3.2. First we shall prove an analogy of Theorem 2.

Theorem 6. *Suppose that $f \in L_q^{p,r}$ with fixed $1 \leq p \leq \infty$, $q > 0$ and $r \in \mathbb{N}$. Then*

$$(26) \quad \|\mathcal{P}_{n;r}(f) - f\|_{p,q} \leq M_1 n^{-r} \omega_1\left(f^{(r)}; L_q^p; \frac{1}{n}\right)$$

for every $n \geq q + 1$, where $M_1 = (r + 2)(1 + 2q)^{r+2}$.

Proof. For every $f \in L_q^{p,r}$ and $x, t \in \mathbb{R}$ there holds the following Taylor-type formula:

$$(27) \quad f(x) = \sum_{j=0}^r \frac{f^{(j)}(t)}{j!} (x - t)^j + \frac{(x - t)^r}{(r - 1)!} I_r(t, x),$$

where

$$(28) \quad I_r(t, x) := \int_0^1 (1 - u)^{r-1} \left[f^{(r)}(t + u(x - t)) - f^{(r)}(t) \right] du.$$

From (27), (28) and (3) there results that

$$F_r(t, x) = f(x) - \frac{(x - t)^r}{(r - 1)!} I_r(t, x),$$

and next by (2), (13) and (7) it follows that

$$(29) \quad \begin{aligned} \mathcal{P}_{n;r}(f; x) - f(x) &= \frac{(-1)^{r+1}}{(r - 1)!} \mathcal{P}_n((t - x)^r I_r(t, x); x) \\ &= \frac{(-1)^{r+1} n}{2(r - 1)!} \int_{\mathbb{R}} \left(t^r \int_0^1 (1 - u)^{r-1} \Delta_{-ut}^1 f^{(r)}(x + t) du \right) e^{-n|t|} dt \end{aligned}$$

for $x \in \mathbb{R}$ and $n \geq 2q + 1$.

If $1 \leq p < \infty$, then using the Minkowski inequality and (5)–(9) and (12), we get from (29):

$$\begin{aligned} &\|\mathcal{P}_{n;r}(f) - f\|_{p,q} \\ &= \frac{n}{2(r - 1)!} \left(\int_{\mathbb{R}} |e^{-q|x}| \int_{\mathbb{R}} t^r e^{-n|t|} \left(\int_0^1 (1 - u)^{r-1} \Delta_{-ut}^1 f^{(r)}(x + t) du \right) dt \right)^{1/p} \\ &\leq \frac{n}{2(r - 1)!} \int_{\mathbb{R}} |t|^r e^{-(n-q)|t|} \left(\int_0^1 (1 - u)^{r-1} \|\Delta_{-ut}^1 f^{(r)}(\cdot)\|_{p,q} du \right) dt \\ &\leq \frac{n}{2(r - 1)!} \int_{\mathbb{R}} |t|^r e^{-(n-q)|t|} \left(\int_0^1 (1 - u)^{r-1} \omega_1\left(f^{(r)}; L_q^p; u|t|\right) du \right) dt \\ &\leq \frac{n}{2r!} \int_{\mathbb{R}} |t|^r e^{-(n-q)|t|} \omega_1\left(f^{(r)}; L_q^p; |t|\right) dt \\ &\leq \frac{n}{r!} \omega_1\left(f^{(r)}; L_q^p; \frac{1}{n}\right) \int_0^\infty t^r (1 + nt) e^{-(n-2q)t} dt \\ &= \omega_1\left(f^{(r)}; L_q^p; \frac{1}{n}\right) \left(\frac{n}{(n - 2q)^{r+1}} + \frac{(1 + r)n^2}{(n - 2q)^{r+2}} \right) \end{aligned}$$

for $n \geq 2q + 1$, which implies (26) for $1 \leq p < \infty$.

The proof of (26) for $f \in L_q^{\infty,r}$ is analogous. □

From Theorem 6 we can derive the following

Corollary 2. *If $f \in L_q^{p,r}$, $1 \leq p \leq \infty$, $q > 0$ and $r \in \mathbb{N}$, then*

$$\lim_{n \rightarrow \infty} n^r \|\mathcal{P}_{n;r}(f) - f\|_{p,q} = 0.$$

Moreover, if $f^{(r)} \in \text{Lip}_M^1(L_q^p; \alpha)$ with some $0 < \alpha \leq 1$ and $M > 0$ then

$$\|\mathcal{P}_{n;r}(f) - f\|_{p,q} = O(n^{-r-\alpha}) \quad \text{as } n \rightarrow \infty.$$

Arguing analogously to the proof of Theorem 2 given in paper [12] and applying Corollary 1, we can obtain the following Voronovskaya-type theorem for operators $\mathcal{P}_{n;r}$.

Theorem 7. *Let $f \in L_q^{\alpha,r}$ with fixed $r \in \mathbb{N}$ and $q > 0$. Then*

$$\begin{aligned} \mathcal{P}_{n;r}(f; x) - f(x) &= \frac{(-1)^r - 1}{2n^{r+1}} f^{(r+1)}(x) \\ &+ \frac{(r+1)[1 + (-1)^r]}{2n^{r+2}} f^{(r+2)}(x) + o(n^{-r-2}) \quad \text{as } n \rightarrow \infty, \end{aligned}$$

at every $x \in \mathbb{R}$. In particular, if r is even number, then

$$(30) \quad \lim_{n \rightarrow \infty} n^{r+2} [\mathcal{P}_{n;r}(f; x) - f(x)] = (r+1)f^{(r+2)}(x)$$

at every $x \in \mathbb{R}$.

Similarly to Theorem 4 now we shall estimate the rate of convergence given by (30).

Theorem 8. *Let $q > 0$ and even number $r \in \mathbb{N}$ be fixed. Then for every $f \in L_q^{\infty,r+2}$ and $n \geq 2q + 1$ there holds*

$$(31) \quad \|n^{r+2} [\mathcal{P}_{n;r}(f) - f] - (r+1)f^{(r+2)}\|_{p,q} \leq M_2 \omega_1\left(f^{(r+2)}; L_q^\infty; \frac{1}{n}\right),$$

where $M_2 = (1 + 2q)^{r+4}(r+4)^2$.

Proof. Similarly to the proof of Theorem 6 we use the Taylor-type formula of $f \in L_q^{\infty,r+2}$:

$$(32) \quad f(x) = \sum_{j=0}^{r+2} \frac{f^{(j)}(t)}{j!} (x-t)^j + \frac{(x-t)^{r+2}}{(r+1)!} I_1(t, x),$$

for $x, t \in \mathbb{R}$, where

$$(33) \quad I_1(t, x) := \int_0^1 (1-u)^{r+1} \left[f^{(r+2)}(t+u(x-t)) - f^{(r+2)}(t) \right] du.$$

Analogously for $f^{(r+1)} \in L_q^{\infty,1}$ and $x, t \in \mathbb{R}$ we have

$$(34) \quad f^{(r+1)}(t) = f^{(r+1)}(x) + f^{(r+2)}(x)(t-x) + (t-x)I_2(t, x)$$

with

$$(35) \quad I_2(t, x) := \int_0^1 \left[f^{(r+2)}(x + u(t-x)) - f^{(r+2)}(x) \right] du.$$

By (3) and (34) the formula (32) can be rewritten in the form:

$$(36) \quad \begin{aligned} f(x) &= F_r(t, x) + \frac{(x-t)^{r+1}}{(r+1)!} f^{(r+1)}(x) \\ &\quad + \left(\frac{1}{(r+2)!} - \frac{1}{(r+1)!} \right) f^{(r+2)}(x) (x-t)^{r+2} \\ &\quad - \frac{(x-t)^{r+2}}{(r+1)!} I_2(t, x) + \frac{(x-t)^{r+2}}{(r+2)!} \left[f^{(r+2)}(t) - f^{(r+2)}(x) \right] \\ &\quad + \frac{(x-t)^{r+2}}{(r+1)!} I_1(t, x) \quad \text{for } x, t \in \mathbb{R}. \end{aligned}$$

Let now $x \in \mathbb{R}$ be a fixed point. Using operator \mathcal{P}_n and (1)–(3) and (13)–(15), we get from (36):

$$f(x) = \mathcal{P}_{n,r}(f; x) - \frac{r+1}{n^{r+2}} f^{(r+2)}(x) + \sum_{i=1}^3 T_i(x) \quad \text{for } n \geq 2q+1,$$

where

$$\begin{aligned} T_1(x) &:= \frac{1}{(r+1)!} \mathcal{P}_n \left((t-x)^{r+2} I_2(t, x); x \right), \\ T_2(x) &:= \frac{1}{(r+2)!} \mathcal{P}_n \left((t-x)^{r+2} \left[f^{(r+2)}(t) - f^{(r+2)}(x) \right]; x \right), \\ T_3(x) &:= \frac{1}{(r+1)!} \mathcal{P}_n \left((t-x)^{r+2} I_1(t, x); x \right). \end{aligned}$$

Consequently we have

$$(37) \quad \|n^{r+2} [\mathcal{P}_{n,r}(f) - f] - (r+1)f^{(r+2)}\|_{\infty, q} \leq n^{r+2} \sum_{i=1}^3 \|T_i\|_{\infty, q}.$$

From (35) and (6)–(9) it follows that

$$\begin{aligned} v_q(x) |T_1(x)| &\leq \frac{e^{-q|x|}}{(r+1)!} \mathcal{P}_n \left(|t-x|^{r+2} |I_2(t, x)|; x \right) \\ &\leq \frac{1}{(r+1)!} \mathcal{P}_n \left(|t-x|^{r+2} \omega_1 \left(f^{(r+2)}; L_q^\infty; |t-x| \right); x \right) \\ &\leq \frac{1}{(r+1)!} \omega_1 \left(f^{(r+2)}; L_q^\infty; \frac{1}{n} \right) \\ &\quad \times \left[\mathcal{P}_n \left(|t-x|^{r+1} e^{q|t-x|}; x \right) + n \mathcal{P}_n \left(|t-x|^{r+2} e^{q|t-x|}; x \right) \right] \end{aligned}$$

and further by (15) we have

$$(38) \quad \|T_1\|_{\infty,q} \leq \frac{1}{(r+1)!} \omega_1\left(f^{(r+2)}; L_q^\infty; \frac{1}{n}\right) \left[\frac{n(r+2)!}{(n-q)^{r+3}} + \frac{n^2(r+3)!}{(n-q)^{r+4}} \right].$$

Analogously, by (6)–(9) there results that

$$\begin{aligned} |f^{(r+2)}(t) - f^{(r+2)}(x)| &\leq \omega_1\left(f^{(r+2)}; L_q^\infty; |t-x|\right) e^{q|x|} \\ &\leq e^{q|x|+q|t-x|} (1+n|t-x|) \omega_1\left(f^{(r+2)}; L_q^\infty; \frac{1}{n}\right) \end{aligned}$$

and from (33):

$$\begin{aligned} |I_1(t,x)| &\leq \int_0^1 (1-u)^{r+1} \omega_1\left(f^{(r+2)}; L_q^\infty; u|t-x|\right) e^{q|t|} du \\ &\leq e^{q|t|} \omega_1\left(f^{(r+2)}; L_q^\infty; |t-x|\right) \int_0^1 (1-u)^{r+1} du \\ &\leq \frac{1}{r+2} e^{q|t|+q|t-x|} (1+n|t-x|) \omega_1\left(f^{(r+2)}; L_q^\infty; \frac{1}{n}\right). \end{aligned}$$

Using the above inequalities and (15), we deduce that

$$(39) \quad \|T_2\|_{\infty,q} \leq \omega_1\left(f^{(r+2)}; L_q^\infty; \frac{1}{n}\right) \left[\frac{n}{(n-q)^{r+3}} + \frac{(r+3)n^2}{(n-q)^{r+4}} \right]$$

and

$$(40) \quad \|T_3\|_{\infty,q} \leq \omega_1\left(f^{(r+2)}; L_q^\infty; \frac{1}{n}\right) \left[\frac{n}{(n-2q)^{r+3}} + \frac{(r+3)n^2}{(n-2q)^{r+4}} \right],$$

for $n \geq 2q+1$. Summarizing (37)–(40), we immediately obtain the desired inequality (31). \square

Remarks 1. Theorem 6 shows that the order of approximation of function $f \in L_q^{p,r}$ by $\mathcal{P}_{n,r}(f)$ is dependent on r and it improves if r grows. Moreover, Theorem 6 and Theorem 2 show that the operators $\mathcal{P}_{n,r}$ with $r \geq 2$ have better approximation properties than \mathcal{P}_n for $f \in L_q^{p,r}$.

We mention also that the similar theorems can be obtained for the Gauss-Weierstrass operators

$$W_n(f; x) := \sqrt{n/\pi} \int_{\mathbb{R}} f(x-t) e^{-nt^2} dt, \quad x \in \mathbb{R}, n \in \mathbb{N},$$

defined in exponential weighted spaces $L_q^p(\mathbb{R})$ with the weighted function $v_q(x) = e^{-qx^2}$, $q > 0$.

REFERENCES

- [1] Becker, M., Kucharski, D., Nessel, R. J., *Global approximation theorems for the Szász-Mirakyan operators in exponential weight spaces*, in *Linear Spaces and Approximation*, Proc. Conf. Oberwolfach, 1977, Birkhäuser Verlag, Basel, 1978, pp. 319–333.
- [2] Brown, B. M., Elliott, D., Paget, D. F., *Lipschitz constants for the Bernstein polynomials of Lipschitz continuous function*, *J. Approx. Theory* **49** (1987), 196–199.
- [3] Butzer, P. L., Nessel, R. J., *Fourier Analysis and Approximation*, vol. 1, Birkhäuser Verlag, Basel und Stuttgart, 1971.
- [4] De Vore, R. A., Lorentz, G. G., *Constructive Approximation*, Springer-Verlag, Berlin, 1993.
- [5] Deeba, E., Mohapatra, R. N., Rodriguez, R. S., *On the degree of approximation of some singular integrals*, *Rend. Mat. Appl. (7)* (1988), 345–355.
- [6] Ditzian, Z., Totik, V., *Moduli of Smoothness*, Springer-Verlag, New-York, 1987.
- [7] Khan, A., *On the degree of approximation of K. Picard and E. Poisson - Cauchy singular integrals*, *Rend. Mat. Appl. (7)* (1982), 123–128.
- [8] Khan, M. K., Vecchia, B. D., Fassih, A., *On the monotonicity of positive linear operators*, *J. Approx. Theory* **92** (1998), 22–37.
- [9] Kirov, G. H., *A generalization of the Bernstein polynomials*, *Math. Balkanica (N.S.)* **2** (2) (1992), 147–153.
- [10] Leśniewicz, A., Rempulska, L., Wasiak, J., *Approximation properties of the Picard singular integral in exponential weight spaces*, *Publ. Mat.* **40** (2) (1996), 233–242.
- [11] Mohapatra, R. N., Rodriguez, R. S., *On the rate of convergence of singular integrals for Hölder continuous functions*, *Math. Nachr.* **149** (1990), 117–124.
- [12] Rempulska, L., Walczak, Z., *On modified Picard and Gauss - Weierstrass singular integrals*, *Ukrain. Math. Zh.* **57** (11) (2005), 1577–1584.
- [13] Timan, A. F., *Theory of Approximation of Functions of a Real Variable*, Moscow, 1960, (Russian).

INSTITUTE OF MATHEMATICS, POZNAN UNIVERSITY OF TECHNOLOGY
UL. PIOTROWO 3A, 60-965 POZNAŃ, POLAND
E-mail: lucyna.rempulska@math.put.poznan.pl