

New generalized sets in N -topological spaces

M. Lellis Thivagar and M. Arockia Dasan

Abstract. By introducing the topological generalized closed sets into N -topological space, this paper establishes that the union of τ_i -generalized closed sets need not be $N\tau$ -generalized closed set. Further, the collection of $N\tau\tilde{g}$ closed sets form a topology. Apart from this, $N\tau\tilde{g}$ closed sets are characterized by means of $N\tau^\#gs$ -kernel.

M.S.C. 2010: 54A05, 54A10.

Key words: N -topology; $N\tau$ - g closed; $N\tau\hat{g}$ closed; $N\tau^*g$ closed; $N\tau\#gs$ closed.

1 Introduction

The concept of semi-open sets and generalized closed sets were initiated by Norman Levine [5,6] and also established their fundamental properties. O.Njastad [9] developed α -open sets and investigated its relationship with other open sets. Mashhour et al. [7,8] characterized pre-open sets and α -open sets with their continuous functions. Abd El-Monsef et al. [1] defined β -open sets with the properties of β -continuous mappings. J.Dontchev [2] evolved the concept of generalized semi-pre closed sets and derived their properties. P. Sundaram et al. [10] characterized the semi-generalized closed sets and their mappings. Lellis Thivagar et al. [3] discovered a geometrical structure of N -topological space with the N -topological open sets. Further Lellis Thivagar and Arockia Dasan [4] established some weak forms of open sets in N -topological space along with their mappings. In this paper, we discuss various kinds of generalized closed sets in N -topological spaces and establish their relationship. We also find that the $N\tau\tilde{g}$ closed sets forms a topology which places between $N\tau$ -closed and $N\tau$ -generalized closed sets.

2 Preliminaries

In this section we recall some known definitions and results of N -topological space and weak open sets that will be used in the following sections. By the space $(X, N\tau)$, we mean, N -topological space with N -topology on X with no separation axioms are assumed unless specifically stated.

Definition 2.1 (3). Let X be a non empty set, $\tau_1, \tau_2, \dots, \tau_N$ be N -arbitrary topologies defined on X . Then the collection $N\tau = \{S \subseteq X : S = (\bigcup_{i=1}^N A_i) \cup (\bigcap_{i=1}^N B_i), A_i, B_i \in \tau_i\}$, is said to be N -topology if it satisfying the following axioms:

- (i) $X, \emptyset \in N\tau$.
- (ii) $\bigcup_{i=1}^{\infty} S_i \in N\tau$ for all $\{S_i\}_{i=1}^{\infty} \in N\tau$.
- (iii) $\bigcap_{i=1}^n S_i \in N\tau$ for all $\{S_i\}_{i=1}^n \in N\tau$.

Then the ordered pair $(X, N\tau)$ is called an N -topological space on X and the elements of the collection $N\tau$ are known as $N\tau$ -open sets on X . A subset A of X is said to be $N\tau$ -closed on X if the complement of A is $N\tau$ -open on X . The set of all $N\tau$ -open sets on X and the set of all $N\tau$ -closed sets on X are respectively denoted by $N\tau O(X)$ and $N\tau C(X)$.

Definition 2.2 (3). Let $(X, N\tau)$ be an N -topological space and S be a subset of X . Then

- (i) the $N\tau$ -interior of S is defined as $N\tau\text{-int}(S) = \cup\{G : G \subseteq S \text{ and } G \text{ is } N\tau\text{-open}\}$.
- (ii) the $N\tau$ -closure of S is defined as $N\tau\text{-cl}(S) = \cap\{F : S \subseteq F \text{ and } F \text{ is } N\tau\text{-closed}\}$.

Definition 2.3 (4). A subset A of N -topological space $(X, N\tau)$ is said to be

- (i) $N\tau\alpha$ -open if $A \subseteq N\tau\text{-int}(N\tau\text{-cl}(N\tau\text{-int}(A)))$.
- (ii) $N\tau$ semi-open if $A \subseteq N\tau\text{-cl}(N\tau\text{-int}(A))$.
- (iii) $N\tau$ pre-open if $A \subseteq N\tau\text{-int}(N\tau\text{-cl}(A))$.
- (iv) $N\tau\beta$ -open if $A \subseteq N\tau\text{-cl}(N\tau\text{-int}(N\tau\text{-cl}(A)))$.

The complement of above open sets are called respective closed sets. The family of $N\tau\alpha$ (resp. $N\tau$ -semi, $N\tau$ -pre and $N\tau\beta$) open sets is denoted by $N\tau\alpha O(X)$ (resp. $N\tau SO(X)$, $N\tau PO(X)$ and $N\tau\beta O(X)$).

Theorem 2.1 (4). *In an N -topological space $(X, N\tau)$, the following are true:*

- (i) every $N\tau$ -open set is $N\tau\alpha$ open.
- (ii) every $N\tau\alpha$ open set is both $N\tau$ -semi and $N\tau$ -pre open, vice versa.
- (iii) every $N\tau$ -semi open set is $N\tau\beta$ open.
- (iv) every $N\tau$ -pre open set is $N\tau\beta$ open.

3 Generalized closed sets in N -topological spaces

This section introduce the classical generalized closed sets into N -topological space. We also state that the union of τ_i -generalized closed sets need not be $N\tau$ -generalized closed set and establish their relationships.

Definition 3.1. A subset A of *N*-topological space $(X, N\tau)$ is said to be

- (i) $N\tau$ generalized-closed (briefly $N\tau g$ -closed) if $N\tau-cl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau$ -open in $(X, N\tau)$.
- (ii) $N\tau\alpha$ generalized-closed (briefly $N\tau\alpha g$ -closed) if $N\tau-\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau$ -open in $(X, N\tau)$.
- (iii) $N\tau$ generalized α -closed (briefly $N\tau g\alpha$ -closed) if $N\tau-\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau\alpha$ -open in $(X, N\tau)$.
- (iv) $N\tau$ generalized semi-closed (briefly $N\tau gs$ -closed) if $N\tau-scl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau$ -open in $(X, N\tau)$.
- (v) $N\tau$ semi generalized-closed (briefly $N\tau sg$ -closed) if $N\tau-scl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau$ semi-open in $(X, N\tau)$.
- (vi) $N\tau\hat{g}$ -closed if $N\tau-cl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau$ semi-open in $(X, N\tau)$.
- (vii) $N\tau^*g$ -closed if $N\tau-cl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau\hat{g}$ -open in $(X, N\tau)$.
- (viii) $N\tau^\#g$ -semi closed (briefly $N\tau^\#gs$ -closed) if $N\tau-scl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau^*g$ -open in $(X, N\tau)$.
- (ix) $N\tau\tilde{g}$ -closed if $N\tau-cl(A) \subseteq U$ whenever $A \subseteq U$ and U is $N\tau^\#gs$ -open in $(X, N\tau)$.

The complement of above *N*-topological generalized closed set is called respective generalized open sets. The set of all $N\tau g$ -closed (resp. $N\tau\alpha g$ -closed, $N\tau g\alpha$ -closed, $N\tau gs$ -closed, $N\tau sg$ -closed, $N\tau\hat{g}$ -closed, $N\tau^*g$ -closed, $N\tau^\#gs$ -closed, $N\tau\tilde{g}$ -closed) sets of $(X, N\tau)$ is denoted by $N\tau GC(X)$ (resp. $N\tau\alpha GC(X)$, $N\tau G\alpha C(X)$, $N\tau GSC(X)$, $N\tau SGC(X)$, $N\tau\hat{G}C(X)$, $N\tau^*GC(X)$, $N\tau^\#GSC(X)$, $N\tau\tilde{G}C(X)$).

The following example illustrates the uniqueness of this paper namely the union of generalized closed sets of topological spaces $(X, \tau_1), (X, \tau_2), \dots, (X, \tau_N)$ need not be a generalized closed set of *N*-topological space.

Example 3.2. If $N = 2$, $X = \{a, b, c\}$, consider $\tau_1 = \{\emptyset, X, \{a\}\}$ and $\tau_2 = \{\emptyset, X, \{a, b\}\}$, then $2\tau O(X) = \{\emptyset, X, \{a\}, \{a, b\}\}$, $\tau_1 GC(X) = \tau_1\alpha GC(X) = \tau_1 GSC(X) = \{\emptyset, X, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}\}$, $\tau_2 GC(X) = \tau_2\alpha GC(X) = \tau_2 GSC(X) = \{\emptyset, X, \{c\}, \{a, c\}, \{b, c\}\}$, $2\tau GC(X) = \{\emptyset, X, \{c\}, \{a, c\}, \{b, c\}\}$, $\tau_1^*GC(X) = \{\emptyset, X, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}\}$, $\tau_2^*GC(X) = \{\emptyset, X, \{c\}, \{a, c\}, \{b, c\}\}$, $\tau_1\tilde{G}C(X) = \tau_1\hat{G}C(X) = \{\emptyset, X, \{b, c\}\}$, $\tau_2\tilde{G}C(X) = \tau_2\hat{G}C(X) = \{\emptyset, X, \{c\}, \{a, c\}, \{b, c\}\}$, $2\tau\tilde{G}C(X) = \{\emptyset, X, \{c\}, \{a, c\}, \{b, c\}\}$, $2\tau\alpha GC(X) = 2\tau GSC(X) = \{\emptyset, X, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}$, $2\tau\hat{G}C(X) = 2\tau\tilde{G}C(X) = \{\emptyset, X, \{c\}, \{b, c\}\}$ and $2\tau^*GC(X) = \{\emptyset, X, \{c\}, \{a, c\}, \{b, c\}\}$. Hence we observe that $\tau_1\tilde{G}C(X) \cup \tau_2\tilde{G}C(X) \neq 2\tau\tilde{G}C(X)$, $\tau_1 GC(X) \cup \tau_2 GC(X) \neq 2\tau GC(X)$, $\tau_1\alpha GC(X) \cup \tau_2\alpha GC(X) \neq 2\tau\alpha GC(X)$ and $\tau_1^*GC(X) \cup \tau_2^*GC(X) \neq 2\tau^*GC(X)$.

Example 3.3. If $N = 2$, $X = \{a, b, c\}$, consider $\tau_1 = \{\emptyset, X, \{a\}, \{a, b\}\}$ and $\tau_2 = \{\emptyset, X, \{a, c\}\}$, then $2\tau O(X) = \{\emptyset, X, \{a\}, \{a, b\}, \{a, c\}\}$, $\tau_1\#GSC(X) = \{\emptyset, X, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}$, $\tau_2\#GSC(X) = \{\emptyset, X, \{b\}, \{a, b\}, \{b, c\}\}$, $2\tau\#GSC(X) = \{\emptyset, X, \{b\}, \{c\}, \{b, c\}\}$. Hence we find that $\tau_1\#GSC(X) \cup \tau_2\#GSC(X) \neq 2\tau\#GSC(X)$.

By the following lemma we over come the above hurdles under certain conditions.

Lemma 3.1. (i) *If every $N\tau$ -open set is τ_i -open, then every τ_i -g closed set is $N\tau$ -g closed for $i = 1, 2, \dots, N$.*

(ii) *If every $N\tau$ -open set is τ_i -open and $N\tau\text{-scl}(A) \subseteq \tau_i\text{-scl}(A)$, then every τ_i -gs closed set is $N\tau$ -gs closed for $i = 1, 2, \dots, N$.*

(iii) *If $\tau_1 SO(X) = \tau_2 SO(X) = \dots = N\tau SO(X)$, then every τ_i -sg closed set is $N\tau$ -sg closed for $i = 1, 2, \dots, N$.*

(iv) *If every $N\tau$ -open set is τ_i -open and $N\tau\text{-}\alpha\text{cl}(A) \subseteq \tau_i\text{-}\alpha\text{cl}(A)$, then every τ_i - α g closed set is $N\tau$ - α g closed for $i = 1, 2, \dots, N$.*

(v) *If $\tau_1\alpha O(X) = \tau_2\alpha O(X) = \dots = N\tau\alpha O(X)$, then every τ_i -g α closed set is $N\tau$ -g α closed for $i = 1, 2, \dots, N$.*

(vi) *If every $N\tau$ - \hat{g} open set is τ_i - \hat{g} open, then every τ_i -*g closed set is $N\tau$ -*g closed.*

(vii) *If every $N\tau$ -*g open set is τ_i -*g open and $N\tau\text{-scl}(A) \subseteq \tau_i\text{-scl}(A)$, then every τ_i -#gs closed set is $N\tau$ -#gs closed.*

(viii) *If every $N\tau$ -#gs open set is τ_i -#gs open, then every τ_i - \tilde{g} closed set is $N\tau$ - \tilde{g} .*

Proof. Here we shall prove parts (i), (iii), (iv) and (viii). The remaining parts can be proved similarly.

(i) *Let A be a τ_i -g closed set and U be a $N\tau$ -open set containing A , then by hypothesis, $\tau_i\text{-cl}(A) \subseteq U$ implies $N\tau\text{-cl}(A) \subseteq U$. Therefore A is $N\tau$ -g closed.*

(iii) *Let A be a τ_i -sg closed set and U be a $N\tau$ -semi open set containing A , then $\tau_i\text{-scl}(A) \subseteq U$ implies $N\tau\text{-scl}(A) \subseteq U$. Therefore A is $N\tau$ -sg closed.*

(iv) *Let A be a τ_i - α g closed set and U be a $N\tau$ -open set containing A , then $\tau_i\text{-}\alpha\text{cl}(A) \subseteq U$ implies $N\tau\text{-}\alpha\text{cl}(A) \subseteq U$. Therefore A is $N\tau$ - α g closed.*

(viii) *Let A be a τ_i - \tilde{g} closed set and U be a $N\tau$ -#gs-open set containing A , then $\tau_i\text{-cl}(A) \subseteq U$ implies $N\tau\text{-cl}(A) \subseteq U$. Therefore A is $N\tau$ - \tilde{g} closed.*

The following proposition is sibling of classical topological results.

Proposition 3.2. *Let $(X, N\tau)$ be an N -topological space, then every*

(i) *$N\tau$ -closed set is $N\tau$ -g closed.*

(ii) *$N\tau$ -semi closed set is $N\tau$ -#gs closed.*

(iii) *$N\tau$ - α closed set is $N\tau$ -#gs closed.*

(iv) $N\tau$ - g closed set is $N\tau$ - αg closed.

(v) $N\tau$ - g closed set is $N\tau$ - gs closed.

(vi) $N\tau$ - sg closed set is $N\tau$ - β closed.

(vii) $N\tau$ - $g\alpha$ closed set is $N\tau$ -pre closed.

The following examples illustrate that the converse of the above proposition need not be true.

Example 3.4. If $N = 6$, $X = \{a, b, c, d\}$, consider $\tau_1 = \{\emptyset, X, \{a\}\}$, $\tau_2 = \{\emptyset, X, \{c, d\}\}$, $\tau_3 = \{\emptyset, X, \{a, c, d\}\}$, $\tau_4 = \{\emptyset, X, \{b, c, d\}\}$, $\tau_5 = \{\emptyset, X, \{a\}, \{b, c, d\}\}$ and $\tau_6 = \{\emptyset, X, \{a\}, \{c, d\}, \{a, c, d\}\}$, then $6\tau O(X) = \{\emptyset, X, \{a\}, \{c, d\}, \{a, c, d\}, \{b, c, d\}\}$, $6\tau C(X) = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{b, c, d\}\}$ and $6\tau\tilde{G}C(X) = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{b, c\}, \{b, d\}, \{a, b, c\}, \{a, b, d\}, \{b, c, d\}\}$. Here the set $\{c\}$ is 6τ -pre closed and 6τ - β closed but not 6τ - sg closed and not 6τ - $g\alpha$ closed.

Example 3.5. If $N = 2$, $X = \{a, b, c\}$, consider $\tau_1 = \{\emptyset, X, \{a\}\}$ and $\tau_2 = \{\emptyset, X, \{a, b\}\}$, then $2\tau O(X) = \{\emptyset, X, \{a\}, \{a, b\}\}$. Here the set $\{b\}$ is 2τ - αg closed and 2τ - gs closed but not 2τ - g closed. Also the set $\{a, c\}$ is 2τ - g closed but not 2τ -closed.

Example 3.6. If $N = 2$, $X = \{a, b, c\}$, consider $\tau_1 = \{\emptyset, X, \{a\}\}$ and $\tau_2 = \{\emptyset, X, \{b, c\}\}$, then $2\tau O(X) = \{\emptyset, X, \{a\}, \{b, c\}\}$. Here the set $\{b\}$ is 2τ - $\#gs$ closed but not 2τ - α closed and not 2τ -semi closed.

Proposition 3.3. Let $(X, N\tau)$ be an N -topological space, then every

(i) $N\tau$ -closed set is $N\tau$ - \tilde{g} closed.

(ii) $N\tau$ - \tilde{g} closed set is $N\tau$ - \hat{g} closed.

(iii) $N\tau$ - \tilde{g} closed set is $N\tau$ - g closed.

(iv) $N\tau$ - \tilde{g} closed set is $N\tau$ - αg closed.

(v) $N\tau$ - \tilde{g} closed set is $N\tau$ - sg closed.

(vi) $N\tau$ - \tilde{g} closed set is $N\tau$ - β closed.

(vii) $N\tau$ - \tilde{g} closed set is $N\tau$ - $g\alpha$ closed.

(viii) $N\tau$ - \tilde{g} closed set is $N\tau$ -pre closed.

(ix) $N\tau$ - \tilde{g} closed set is $N\tau$ - gs closed.

Proof. Here we shall prove part (i) and (iii) only. The proof of the remaining parts are similar.

(i) If A is any $N\tau$ -closed set in $(X, N\tau)$ and U is any $N\tau$ - $\#gs$ open set containing A . Then $N\tau\text{-cl}(A) = A \subseteq U$ implies A is a $N\tau$ - \tilde{g} closed.




Figure 1: The relationship between N -topological generalized closed sets.

(iii) If A is any $N\tau$ - \tilde{g} closed set in $(X, N\tau)$ and U is any $N\tau$ -open set containing A . Since every $N\tau$ -closed set is $N\tau$ -semi closed and every $N\tau$ -semi closed set is $N\tau$ - $\#$ gs closed, then U is $N\tau$ - $\#$ gs open set containing A and so $N\tau\text{-cl}(A) \subseteq U$. Therefore A is a $N\tau$ - g closed.

Example 3.7. The converse of the above proposition need not be true. For $N = 1$, $X = \{a, b, c\}$, $\tau = \tau_1 = \{\emptyset, X, \{a\}\}$. Then the set $\{b\}$ is $g\alpha$ -closed, pre closed and gs -closed but not \tilde{g} -closed set. From example 3.4, the set $\{b, c\}$ is 6τ - \tilde{g} closed but not 6τ -closed. Also from example 3.5, the set $\{b\}$ is 2τ - β closed and 2τ - sg closed but not 2τ - \tilde{g} closed. By example 3.6 we know that the set $\{b\}$ is 2τ - g closed, 2τ - αg closed and 2τ - \hat{g} closed but not 2τ - \tilde{g} closed.

Remark 3.8. From the proposition 3.3, we observe that the set of all $N\tau$ - \tilde{g} closed sets placed between the set of all $N\tau$ -closed and $N\tau$ - g closed sets. It also placed between the set of all $N\tau$ -closed and $N\tau$ - \hat{g} closed sets.

Remark 3.9. The following example states that the $N\tau$ - \tilde{g} closed set is independent of $N\tau$ - α closed as well as $N\tau$ -semi closed set.

Example 3.10. If $N = 2$, $X = \{a, b, c\}$, consider $\tau_1 = \{\emptyset, X, \{a, b\}\}$ and $\tau_2 = \{\emptyset, X\}$, then $2\tau O(X) = \{\emptyset, X, \{a, b\}\}$. Here the set $\{a, c\}$ is 2τ - \tilde{g} closed but not 2τ - α closed and not 2τ -semi closed. From example 3.7, the set $\{b\}$ is α closed and semi closed but not \tilde{g} -closed.

Remark 3.11. The concept of generalized closed sets in N -topological space can be described in the diagram below where the reversible implication is not possible.

4 Characterization of $N\tau$ - \tilde{g} closed sets

In this section, we introduce $N\tau^{\#}gs\text{-ker}(A)$ and discuss the essential conditions for $N\tau$ - \tilde{g} closed sets in terms of $N\tau^{\#}gs\text{-ker}(A)$.

Definition 4.1. Let A be a subset of N -topological space $(X, N\tau)$, then $N\tau^{\#}gs\text{-ker}(A)$ is defined as the intersection of all $N\tau$ - $\#g$ semi open subsets of X containing A .

Lemma 4.1. A subset A of an N -topological space $(X, N\tau)$ is $N\tau$ - \tilde{g} closed if and only if $N\tau\text{-cl}(A) \subseteq N\tau^{\#}gs\text{-ker}(A)$.

Proof. Let A is $N\tau$ - \tilde{g} closed in $(X, N\tau)$, then for every $N\tau$ - $\#g$ open set U containing A , $N\tau\text{-cl}(A) \subseteq U$. Assume $x \in N\tau\text{-cl}(A)$ and if $x \notin N\tau^{\#}gs\text{-ker}(A)$, then there exist a $N\tau$ - $\#g$ open set U containing A such that $x \notin U$ implies $x \notin N\tau\text{-cl}(A)$. This is a contradiction to our assumption. Conversely, if $N\tau\text{-cl}(A) \subseteq N\tau^{\#}gs\text{-ker}(A)$ and U is a $N\tau$ - $\#g$ open set containing A , then $N\tau\text{-cl}(A) \subseteq N\tau^{\#}gs\text{-ker}(A) \subseteq U$. Therefore A is $N\tau$ - \tilde{g} closed. \square The following remarks are similar to the topological results of Dontchev [2].

Remark 4.2. Let x be a point of $(X, N\tau)$. Then $\{x\}$ is either $N\tau$ -nowhere dense or $N\tau$ -pre open.

Remark 4.3. In remark 4.2, the decomposition of an N -topological space $(X, N\tau)$, $X = X_1 \cup X_2$, where $X_1 = \{x \in X : \{x\} \text{ is } N\tau\text{-nowhere dense}\}$ and $X_2 = \{x \in X : \{x\} \text{ is } N\tau\text{-pre open}\}$.

Theorem 4.2. For every subset A of an N -topological space $(X, N\tau)$, $X_2 \cap N\tau\text{-cl}(A) \subseteq N\tau^{\#}gs\text{-ker}(A)$.

Proof. Let $x \in X_2 \cap N\tau\text{-cl}(A)$, if $x \notin N\tau^{\#}gs\text{-ker}(A)$, then there exist a $N\tau$ - $\#g$ open set U containing A such that $x \notin U$ implies $X \setminus U$ is a $N\tau$ - $\#g$ closed set containing x . Since $x \in X_2 \cap N\tau\text{-cl}(A)$, $N\tau\text{-int}(N\tau\text{-cl}(\{x\})) \cap A \neq \emptyset$. Thus there is a point $y \in N\tau\text{-int}(N\tau\text{-cl}(\{x\})) \cap A \subseteq (X \setminus U) \cap A \subseteq (X \setminus U) \cap U$, which is a contradiction.

Theorem 4.3. A subset A of an N -topological space $(X, N\tau)$ is $N\tau$ - \tilde{g} closed if and only if $X_1 \cap N\tau\text{-cl}(A) \subseteq A$.

Proof. Let A be a $N\tau$ - \tilde{g} closed set, if $x \in X_1 \cap N\tau\text{-cl}(A)$ implies $x \in X_1$ and $x \in N\tau\text{-cl}(A)$. Since $x \in X_1$, $N\tau\text{-int}(N\tau\text{-cl}(\{x\})) = \emptyset$, $\{x\}$ is $N\tau$ -semi closed. Since every $N\tau$ -semi closed set is $N\tau$ - $\#g$ closed, $\{x\}$ is $N\tau$ - $\#g$ closed. If $x \notin A$ and $U = X \setminus \{x\}$, U is a $N\tau$ - $\#g$ open set containing A and $N\tau\text{-cl}(A) \subseteq U$. This is a contradiction. Conversely, assume $X_1 \cap N\tau\text{-cl}(A) \subseteq A$ and $A \subseteq N\tau^{\#}gs\text{-ker}(A)$, then $X_1 \cap N\tau\text{-cl}(A) \subseteq N\tau^{\#}gs\text{-ker}(A)$. Now $N\tau\text{-cl}(A) = X \cap N\tau\text{-cl}(A) = (X_1 \cup X_2) \cap N\tau\text{-cl}(A) = (X_1 \cap N\tau\text{-cl}(A)) \cup (X_2 \cap N\tau\text{-cl}(A)) \subseteq N\tau^{\#}gs\text{-ker}(A)$. Thus by lemma 4.1, A is $N\tau$ - \tilde{g} closed.

The proof of the following theorems are obvious from the above theorems.

Theorem 4.4. The arbitrary intersection of $N\tau$ - \tilde{g} closed sets is $N\tau$ - \tilde{g} closed.

Theorem 4.5. *If A is $N\tau$ - \tilde{g} closed set and B is $N\tau$ -closed set, then $A \cap B$ is $N\tau$ - \tilde{g} closed.*

Theorem 4.6. *If A and B are two $N\tau$ - \tilde{g} closed sets, then $A \cup B$ is $N\tau$ - \tilde{g} closed set.*

Theorem 4.7. *If a set A is $N\tau$ - \tilde{g} closed, then $N\tau\text{-cl}(A) \setminus A$ contains no non empty $N\tau$ -closed set.*

Proof. Suppose A is $N\tau$ - \tilde{g} closed in $(X, N\tau)$ and F be a $N\tau$ -closed subset of $N\tau\text{-cl}(A) \setminus A$, then F^c is $N\tau$ - $\#gs$ open and $A \subseteq F^c$. From the definition of $N\tau$ - \tilde{g} closed set it follows that $N\tau\text{-cl}(A) \subseteq F^c$ and $F \subseteq (N\tau\text{-cl}(A))^c$. Therefore $F \subseteq N\tau\text{-cl}(A) \cap (N\tau\text{-cl}(A))^c = \emptyset$ and so F is an empty set. \square

Remark 4.4. The converse of the above theorem need not be true. By example 3.7, if $A = \{b\}$, then $cl(A) \setminus A = \{c\}$ does not contain non empty closed set also A is not \tilde{g} -closed in (X, τ) .

Theorem 4.8. *A set A is $N\tau$ - \tilde{g} closed if and only if $N\tau\text{-cl}(A) \setminus A$ contains no non empty $N\tau$ - $\#gs$ closed set.*

Proof. Assume A is $N\tau$ - \tilde{g} closed in $(X, N\tau)$ and F is a $N\tau$ - $\#gs$ closed subset of $N\tau\text{-cl}(A) \setminus A$, then F^c is $N\tau$ - $\#gs$ open containing A and $N\tau\text{-cl}(A) \subseteq F^c$. That is, $F \subseteq (N\tau\text{-cl}(A))^c$. Therefore $F \subseteq N\tau\text{-cl}(A) \cap (N\tau\text{-cl}(A))^c = \emptyset$ and F is an empty set. Conversely, let $N\tau\text{-cl}(A) \setminus A$ contains no non empty $N\tau$ - $\#gs$ closed set and S be $N\tau$ - $\#gs$ open set containing A in $(X, N\tau)$. Suppose $N\tau\text{-cl}(A)$ is not a subset of S , then $N\tau\text{-cl}(A) \cap S^c$ is a non empty $N\tau$ - $\#gs$ closed subset of $N\tau\text{-cl}(A) \setminus A$, which is a contradiction. Therefore A is $N\tau$ - \tilde{g} closed in $(X, N\tau)$. \square

Theorem 4.9. *If A is a $N\tau$ - \tilde{g} closed set and $A \subseteq B \subseteq N\tau\text{-cl}(A)$, then B is also $N\tau$ - \tilde{g} closed.*

Proof: Let A be a $N\tau$ - \tilde{g} closed set and U be any $N\tau$ - $\#gs$ open set containing B , then $N\tau\text{-cl}(B) \subseteq N\tau\text{-cl}(A) \subseteq U$. Therefore $N\tau\text{-cl}(B) \subseteq U$ and hence B is $N\tau$ - \tilde{g} closed.

Theorem 4.10. *If A is $N\tau$ - $\#gs$ open and $N\tau$ - \tilde{g} closed, then A is $N\tau$ -closed.*

Proof. Since A is $N\tau$ - $\#gs$ open, $N\tau\text{-cl}(A) \subseteq A$. Therefore $N\tau\text{-cl}(A) = A$ and hence A is $N\tau$ -closed.

Theorem 4.11. *For each $x \in X$, either $\{x\}$ is $N\tau$ - $\#gs$ closed, or $\{x\}^c$ is $N\tau$ - \tilde{g} closed.*

Proof: If $\{x\}$ is not $N\tau$ - $\#gs$ closed in $(X, N\tau)$, then $\{x\}^c$ is not $N\tau$ - $\#gs$ open. The only $N\tau$ - $\#gs$ open set containing $\{x\}^c$ is the set X itself. Therefore $N\tau\text{-cl}(\{x\}^c) \subseteq X$ and so $\{x\}^c$ is $N\tau$ - \tilde{g} closed in $(X, N\tau)$.

5 Conclusion and future work

A N -topological space is a new space containing N -topologies defined on a non-empty set X . In this paper, we have attempted to define and establish N -topological generalized closed sets in N -topological space. Moreover, we have proved that the

collection of $N\tau\tilde{g}$ closed sets form a topology on X . This can be extended to real-life applications such as fuzzy topology, intuitionistic topology, nano topology, soft topology, etc. It also can open up doors to research areas like supra topology, digital topology, and so on.

Acknowledgements. The author thanks to the the referees for the valuable advice.

References

- [1] M. E. Abd. El-Monsef, S. N. El Deeb, R.A. Mahmoud, *β -open sets and β -continuous mappings*, Bull. Fac. Sci. Assiut Univ. 12(1983), 77-90.
- [2] J. Dontchev, *On generalizing semi-pre open sets*, Mem. Fac.Sci.Kochi Univ.Ser.A.Math. 16(1995) 35-48.
- [3] M. Lellis Thivagar, V. Ramesh, M. Arockia Dasan, *On new structure of N -topology*, Cogent Mathematics, 3(2016), 1204104.
- [4] M. Lellis Thivagar, M. Arockia Dasan, *New topologies via weak N -topological open sets and mappings*, Journal of New Theory, 29 (2019), 49-57.
- [5] N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer Math. Monthly. 70(1963), 36-41.
- [6] N. Levine, *Generalized closed sets in topology*, Rend. Circ. Mat. Palermo. 19(1970), 89-96.
- [7] A. S. Mashhour, I. A. Hasanien, S. N. El-Deeb, *On precontinuous and weak precontinuous mappings*, Proc. Math. Phys. Soc. Egypt. 53(1982), 47-53.
- [8] A. S. Mashhour, I. A. Hasanien, S. N. El-Deeb, (1983). *α -continuous and α -open mappings*, Acta. Math. Hung. 41(3-4)(1983), 213-218.
- [9] O. Njastad, *On some classes of nearly open sets*, Pacific J.Math. 15(1965), 961-970.
- [10] P. Sundaram, H. Maki, K. Balachandran, *Semi-generalized continuous maps and semi- $T_{1/2}$ spaces*, Bull. Fukuoka Univ. Edu. Part III, 40(1991), 33-40.

Author's address:

Maria Antony Lellis Thivagar
 Department of Mathematics, Madurai Kamaraj University,
 Madurai-625021, Tamil Nadu, India.
 E-mail: mlthivagar@yahoo.co.in

Muthayyan Arockia Dasan (*corresponding author*)
 Department of Mathematics, St. Jude's College,
 Thoothoor, Kanyakumari-629176, Tamil Nadu, India.
 E-mail: dassfredy@gmail.com