

The Skew and Relative Derangements of Type B

William Y.C. Chen¹ and Jessica C.Y. Zhang²

Center for Combinatorics, LPMC-TJKLC

Nankai University,

Tianjin 300071, P. R. China

¹chen@nankai.edu.cn, ²zhangcy@mail.nankai.edu.cn

Submitted: Aug 27, 2007; Accepted: Oct 28, 2007; Published: Nov 5, 2007

Mathematics Subject Classification: 05E15, 05A05

Abstract

By introducing the notion of relative derangements of type B , also called signed relative derangements, which are defined in terms of signed permutations, we obtain a type B analogue of the well-known relation between the relative derangements and the classical derangements. While this fact can be proved by using the principle of inclusion and exclusion, we present a combinatorial interpretation with the aid of the intermediate structure of signed skew derangements.

1 Introduction

A derangement on a set $[n] = \{1, 2, \dots, n\}$ is a permutation $\pi = \pi_1\pi_2 \cdots \pi_n$ such that $\pi_i \neq i$ for all $i \in [n]$. A relative derangement $\pi_1\pi_2 \cdots \pi_n$ on $[n]$ is a permutation such that $\pi_{i+1} \neq \pi_i + 1$ for $1 \leq i \leq n - 1$. Let Q_n denote the number of relative derangements on $[n]$, and let D_n denote the number of the derangements on $[n]$. The following relation is well-known, see Brualdi [2, Theorem 6.5.1], or Andreescu and Feng [1, Example 6.11]:

$$Q_n = D_n + D_{n-1}. \quad (1.1)$$

A combinatorial interpretation of (1.1) has been obtained by Chen [3] based on the intermediate structure of skew derangements, which are equivalent to the generalized derangements as studied by Hanson, Seyffarth and Weston [6] and Wang [8]. Clarke, Han and Zeng [4] gave a similar construction based on variant of Foata's first fundamental transformation. The main objective of this paper is to present a type B analogue of

(1.1). This goal is achieved by introducing the notion of signed relative derangements, or relative derangements of type B . The concept of derangements of type B is introduced by Chow [5]. A *signed permutation* π on $[n]$ can be viewed as a bijection on the set $\{\bar{1}, \dots, \bar{n}, 1, \dots, n\}$ such that $\pi(\bar{i}) = \overline{\pi(i)}$. Intuitively, a signed permutation on $[n]$ is just an ordinary permutation $\pi_1\pi_2 \cdots \pi_n$ with some elements associated with a bar $-$. For example, $3\bar{2}\bar{5}1\bar{4}$ is a signed permutation on $\{1, 2, 3, 4, 5\}$. The set of signed permutations on $[n]$ is often denoted by B_n . The following order relation is often imposed on the elements of signed permutations for B_n :

$$\bar{1} < \bar{2} < \cdots < \bar{n} < 1 < 2 < \cdots < n. \quad (1.2)$$

According to the above ordering, for the above signed permutation $3\bar{2}\bar{5}1\bar{4}$, 3 is the largest element and $\bar{2}$ is the smallest. We recall the following definition of derangements of type B .

Definition 1.1. A *derangement of type B on $[n]$* is a signed permutation $\pi_1\pi_2 \cdots \pi_n$ such that $\pi_i \neq i$, for all $i \in [n]$.

For example, $3\bar{2}\bar{5}1\bar{4}$ is a derangement in B_5 , whereas $3\bar{2}\bar{4}1\bar{5}$ has a fixed point 2 . Let D_n^B denote the number of derangements of type B on $[n]$. It is not hard to derive the following formula by using the principle of inclusion-exclusion [4, Chapter 2]:

$$D_n^B = n! \sum_{k=0}^n \frac{(-1)^k \cdot 2^{n-k}}{k!} \quad (1.3)$$

In fact, it is also a consequence of the q -analogue given by Chow [5].

We now give the definition of relative derangements of type B on $[n]$, or signed relative derangements, for short.

Definition 1.2. A *relative derangement of type B on $[n]$* is a signed permutation on $[n]$ such that i is not followed by $i+1$, and \bar{i} is not followed by $\overline{i+1}$, for $1 \leq i \leq n-1$.

For example, $3\bar{2}\bar{4}1\bar{5}$ is a relative derangement in B_5 , while $41\bar{5}\bar{2}\bar{3}$ is not. Let Q_n^B be the number of relative derangements of type B . Our main result is the following type B analogue of the above relation (1.1).

Theorem 1.3. For $n \geq 2$, we have

$$Q_n^B = D_n^B + D_{n-1}^B. \quad (1.4)$$

The first few values of Q_n^B and D_n^B are given below:

n	1	2	3	4	5	6	7	8	...
Q_n^B	2	6	34	262	2562	30278	419234	6651846	...
D_n^B	1	5	29	233	2329	27949	391285	6260561	...

In accordance with the relation (1.4), we adopt the convention that $D_0^B = 1$.

One way to prove the above result for Q_n^B and D_n^B is to derive the following formula for Q_n^B by using the principle of inclusion-exclusion:

$$Q_n^B = n! \cdot 2^n + \sum_{k=1}^{n-1} (-1)^k \cdot \binom{n-1}{k} \cdot (n-k)! \cdot 2^{n-k}. \quad (1.5)$$

However, the details of the algebraic proof will be omitted. Instead, we will provide a combinatorial proof by introducing the structure of signed skew derangements.

2 Signed Skew Derangements

In this section, we first introduce the notion of signed skew derangements and establish a correspondence between signed relative derangements and signed skew derangements. Then we give a characterization of signed permutations that correspond to signed skew derangements. Then we show how to transform a signed skew derangement into a signed derangement. This leads to a combinatorial interpretation of the relation (1.4).

Recall that a skew derangement f on $[n]$ is a bijection from $[n]$ onto $\{0, 1, \dots, n-1\}$ with $f(i) \neq i$ for any $i \in [n]$, see [3]. For signed permutations, we will define signed skew derangements, or skew derangements of type B . Let us begin with the definition of a signed set. A signed set is a set with some elements bearing bars, then a signed set on $[n]$ can be considered the underlying set of a signed permutation. In other words, a signed set on $[n]$ is just the set $[n]$ with some elements bearing bars. For example, $X = \{1, \bar{2}, 3, 4, \bar{5}\}$ is a signed set on $\{1, 2, 3, 4, 5\}$.

Given a signed set X on $[n]$, we denote by $X - 1$ the signed set obtained from X by subtracting 1 from each element in X , where we define the subtraction for barred elements by the rule

$$\bar{i} - 1 = \overline{i - 1}. \quad (2.1)$$

Conversely, the addition to a barred element is given by

$$\bar{i} + 1 = \overline{i + 1}. \quad (2.2)$$

Definition 2.1. *Let X be a signed set on $[n]$. A signed skew derangement on $[n]$ is a bijection f from X to $Y = X - 1$ such that $f(x) \neq x$ for any $x \in X$, where x may be a barred element.*

For example, let $n = 2$, $X = \{\bar{1}, \bar{2}\}$ and $Y = \{\bar{0}, \bar{1}\}$. Then there are two bijections from X to Y : $f_1(\bar{1}) = \bar{0}$, $f_1(\bar{2}) = \bar{1}$ and $f_2(\bar{1}) = \bar{1}$, $f_2(\bar{2}) = \bar{0}$. From the above definition, we can see only the bijection f_1 is a signed skew derangement. The following theorem establishes a bijection between signed relative derangements and signed skew derangements.

Theorem 2.2. *There is a one-to-one correspondence between the set of signed relative derangements on $[n]$ and the set of signed skew derangements on $[n]$.*

Proof. First, given a signed relative derangement $\pi = \pi_1\pi_2\cdots\pi_n$ on $[n]$, we proceed to construct a signed skew derangement f on $[n]$. Let u be the maximum element in the signed permutation $\pi_1\pi_2\cdots\pi_n$ with respect to the order (1.2). Note that in the case of signed permutations, the maximum element is not necessarily the element n . Suppose that $\pi_k = u$. Let us consider the segment $\pi_1\pi_2\cdots\pi_k$. Define

$$f(\pi_1) = \pi_2 - 1, \quad f(\pi_2) = \pi_3 - 1, \quad \dots, \quad f(\pi_{k-1}) = \pi_k - 1, \quad f(\pi_k) = \pi_1 - 1,$$

subject to the above subtraction rule (2.1) if an element π_t is a barred element.

By the definition of signed relative derangement, we claim that f satisfies the condition of a signed skew derangement with respect to the elements $\pi_1, \pi_2, \dots, \pi_k$, namely,

$$f(\pi_1) \neq \pi_1, \quad f(\pi_2) \neq \pi_2, \quad \dots, \quad f(\pi_k) \neq \pi_k.$$

For any $r = 1, 2, \dots, k - 1$, since π is a signed relative derangement, in view of the addition operation (2.2) we see that $\pi_{r+1} \neq \pi_r + 1$ no matter whether π_r is a barred element or not. So we have

$$f(\pi_r) = \pi_{r+1} - 1 \neq \pi_r$$

for $r = 1, 2, \dots, k - 1$. We now consider π_k . Since π_k is the maximum element of π , we find $\pi_1 - 1 \neq \pi_k$. This implies that $f(\pi_k) = \pi_1 - 1 \neq \pi_k$.

Now we can repeat the above procedure for the remaining sequence $\sigma = \pi_{k+1}\pi_{k+2}\cdots\pi_n$. The next step is still to choose the maximum element π_t in σ , then assign the images of f for the elements $\pi_{k+1}, \pi_{k+2}, \dots, \pi_t$. If there are still elements left, we may iterate this procedure until f is completely determined.

It remains to construct the inverse map. Given a signed skew derangement f on $[n]$, we aim to find the corresponding signed relative derangement.

Suppose f is a signed skew derangement from a signed set X to $X - 1$. The first step is to determine π_1 . Assume that u is the maximum element in X with respect to the order (1.2). Then we set $\pi_1 = f(u) + 1$, subject to the above addition rule (2.2) if $f(u)$ is a barred element. Suppose π_r is already located. If $\pi_r \neq u$, then we set $\pi_{r+1} = f(\pi_r) + 1$, using the above rule (2.2) if $f(\pi_r)$ is a barred element, and repeat this process until we reach a step when $\pi_k = u$ for some k .

At this point, we have obtained the segment $\pi_1\pi_2\cdots\pi_k$. Since $f(\pi_r) \neq \pi_r$, we see that $\pi_{r+1} \neq \pi_r + 1$, for $r = 1, \dots, k$. If $k < n$, then we may choose the maximum element

among the remaining elements in X after removing the elements $\pi_1, \pi_2, \dots, \pi_k$, and iterate the above procedure until we obtain the desired signed relative derangement. Thus, we have shown that our construction is a bijection. ■

For example, the signed relative derangement $\bar{7} 8 6 \bar{1} \bar{5} \bar{3} 4 2$ corresponds to the following signed skew derangement:

$$\begin{aligned} f(\bar{7}) &= 8 - 1 = 7, & f(8) &= \bar{7} - 1 = \bar{6}, & f(6) &= 6 - 1 = 5, & f(\bar{1}) &= \bar{5} - 1 = \bar{4}, \\ f(\bar{5}) &= \bar{3} - 1 = \bar{2}, & f(\bar{3}) &= 4 - 1 = 3, & f(4) &= \bar{1} - 1 = \bar{0}, & f(2) &= 2 - 1 = 1. \end{aligned}$$

We now turn our attention to a combinatorial interpretation of the fact that the number of signed skew derangements on $[n]$ equals $D_n^B + D_{n-1}^B$. Our strategy based on the above theorem goes as follows. Given a signed skew derangement f on $[n]$, if we require $f(n) \neq 0$ or $f(\bar{n}) \neq 0$, then f can be viewed as a signed derangement g on $[n]$ because we can replace 0 with n or $\bar{0}$ with \bar{n} ; otherwise, f can be viewed as a signed derangement g on $[n-1]$. In these two situations, we should define $g(i) = f(i)$ or $g(i) = f(\bar{i})$.

However, in order to see the above strategy is valid we need to give a rigorous reasoning. As the first step, we give a characterization of signed permutations on $\{0, 1, \dots, n-1\}$ that correspond to signed skew derangements on $[n]$. Let us consider bijections from a signed set X on $[n]$ to $X-1$. Assume that the elements of X are arranged by the increasing order of their underlying elements, say, $X = \{\sigma_1, \sigma_2, \dots, \sigma_n\}$. It is easy to observe the fact that a bijection f from X to $X-1$ is determined by the signed permutation $\pi = \pi_1 \pi_2 \cdots \pi_n$, where $\pi_i = f(\sigma_i)$. In fact, this is a bijection, because for any signed permutation π on $\{0, 1, \dots, n-1\}$, the elements $\{\pi_1, \pi_2, \dots, \pi_n\}$ determines the signed set $X-1$, which in turn determines X . Hence the map f from X to $X-1$ is easily constructed. The signed permutation π is called the *representation* of f .

For the above signed skew derangement f , we have

$$X = \{\sigma_1, \sigma_2, \dots, \sigma_8\} = \{\bar{1}, 2, \bar{3}, 4, \bar{5}, 6, \bar{7}, 8\}$$

and $\pi = \bar{4} 1 3 \bar{0} \bar{2} 5 7 \bar{6}$.

The following lemma gives a characterization of signed permutations which are representations of signed skew derangements. A bar associated with an element is intuitively considered as a sign. Moreover, for a signed permutation $\pi = \pi_1 \pi_2 \cdots \pi_n$, an element π_i is called a *fixed point* if $\pi_i = i$, whereas it is called a *signed fixed point* if $\pi_i = i$ or \bar{i} . As will be seen, signed fixed points play an important role in establishing the correspondence between signed skew derangements and signed derangements.

Lemma 2.3. *Let π be a signed permutation on $\{0, 1, \dots, n-1\}$, and let X and f be the signed set and the bijection from X to $X-1$ determined by π . Then f has a fixed point if π has a signed fixed point π_i , and $i-1$ and i have the same sign in π .*

The above lemma can be restated as follows. A signed permutation π is a representation of a signed skew derangement if and only if $\pi_i = i$ implies that $\overline{i-1}$ appears in π , and $\pi_i = \bar{i}$ implies that $i-1$ appears in π .

Proof. Let π be a signed permutation on $\{0, 1, 2, \dots, n-1\}$. Let f be a bijection from X to $X-1$ such that π is the representation of f . Then $X-1$ is determined by the entries of π . Hence X is uniquely determined by π . Let $\sigma_1, \sigma_2, \dots, \sigma_n$ be the elements of X arranged in the increasing order of the underlying elements of X . If f has a fixed point, say, $f(x) = x$, for some $x = \sigma_i$. Then we have $\sigma_i = i$ or \bar{i} , and $f(\sigma_i) = \sigma_i = \pi_i$. Since f is a bijection from X to $X-1$, σ_i is a barred element if and only if $i-1$ is a barred element. Thus, we conclude that π_i and $i-1$ have the same sign. This completes the proof. ■

The above characterization indicates that signed skew derangements can be viewed as an intermediate structure between signed relative derangements and signed derangements. Using this characterization of representations of signed skew derangements on $[n]$, we first consider a class of such signed permutations that are in one-to-one correspondence with signed derangements on $[n-1]$.

Lemma 2.4. *There is a bijection between the set of representations of signed skew derangements on $[n]$ that are of the form $\pi = \pi_1\pi_2 \cdots \pi_{n-1}0$ and the set of signed derangements on $[n-1]$.*

For example, there are five signed derangements on $\{1, 2\}$: $\bar{1}\bar{2}$, 21 , $2\bar{1}$, $\bar{2}1$, $\bar{2}\bar{1}$. In the meantime, there are five representations signed skew derangements on $\{1, 2, 3\}$ that are of the form $\pi_1\pi_20$: $\bar{1}\bar{2}0$, 210 , $2\bar{1}0$, $\bar{2}10$, $\bar{2}\bar{1}0$. As in this example, special attention should be paid to the signed derangement $\bar{1}\bar{2}$ with signed fixed points, and to the representation $\bar{1}\bar{2}0$ which also have signed fixed points. In general, we can establish a correspondence as given in the following proof.

Proof. Let $\pi = \pi_1\pi_2 \cdots \pi_{n-1}0$ be a representation of a signed skew derangement on $[n]$. We aim to construct a signed derangement on $[n-1]$ from π . If $\pi_1\pi_2 \cdots \pi_{n-1}$ has no signed fixed point, then it is automatically the desired signed derangement.

We now consider that case when there are some signed fixed points, namely, there exist some i such that $\pi_i = i$ or \bar{i} . Taking the signed fixed point π_i with minimum index i , we observe that whether π_i has a bar or not is determined solely by the appearance of $i-1$ in the sense that it is a barred element or an unbarred element. Iterating this argument, we may deduce that the signed fixed points are uniquely determined by the remaining elements in π . Hence we may always put \bar{i} as the signed fixed points in order to obtain a signed derangement.

Conversely, given a signed derangement $\tau = \tau_1\tau_2 \cdots \tau_{n-1}$, we may identify the signed fixed points τ_i . By the same argument as in the previous paragraph, we can determine the signed fixed points according to the characterization of representations of signed skew derangements so that the resulting signed permutation on $\{0, 1, \dots, n-1\}$ corresponds

to a signed skew derangement on $[n]$. This completes the proof. ■

For example, consider the signed skew derangement f on $\{1, 2, \dots, 8\}$ which has the following representation

$$f(1) f(2) f(\bar{3}) f(\bar{4}) f(5) f(\bar{6}) f(\bar{7}) f(8) = \bar{6} \bar{2} 1 4 \bar{3} 7 \bar{5} 0.$$

It corresponds to the signed derangement $\bar{6} \bar{2} 1 4 \bar{3} 7 \bar{5}$ on $\{1, 2, \dots, 7\}$.

To complete the combinatorial proof of Theorem 1.3, it suffices to consider the second case for the representations of signed skew derangements. The following lemma deals with this case.

Lemma 2.5. *There is a one-to-one correspondence between representations $\pi = \pi_1 \pi_2 \cdots \pi_n$ of signed skew derangements on $[n]$ with $\pi_n \neq 0$ and signed derangements on $[n]$.*

For example, there are five representation $\pi = \pi_1 \pi_2$ of signed skew derangements on $\{1, 2\}$ with $\pi_2 \neq 0$: $01, 0\bar{1}, \bar{0}1, \bar{0}\bar{1}, 1\bar{0}$.

Proof. First, we show that from a representation $\pi = \pi_1 \pi_2 \cdots \pi_n$ of a signed skew derangement with $\pi_n \neq 0$ we can construct a signed derangement $\tau = \tau_1 \tau_2 \cdots \tau_n$. If there is no signed fixed point in π , then we can replace 0 or $\bar{0}$ by n or \bar{n} in π depending whether 0 or $\bar{0}$ appears. Since $\pi_n \neq 0$, we have $\tau_n \neq n$ and so the resulting signed permutation is a signed derangement on $[n]$.

Otherwise, there are some signed fixed points π_i ($1 \leq i \leq n-1$), namely, $\pi_i = i$ or \bar{i} . Using the same argument as in the proof of Lemma 2.4, we see that the signed fixed points are completely determined by the remaining elements in the signed permutation. So we may set all the signed fixed points to barred elements in π . Finally, we may replace 0 by n or $\bar{0}$ by \bar{n} to get a signed derangement τ on $[n]$.

It is clear that the above procedure is reversible. This completes the proof. ■

For example, consider the signed skew derangement f on $\{1, 2, \dots, 8\}$ which has the following representation

$$f(\bar{1}) f(2) f(\bar{3}) f(4) f(\bar{5}) f(6) f(\bar{7}) f(8) = \bar{4} 1 3 \bar{0} \bar{2} 5 7 \bar{6}.$$

The corresponding signed derangement is $\bar{4} 1 \bar{3} \bar{8} \bar{2} 5 7 \bar{6}$.

It is easy to see that Theorem 1.3 follows from the above two lemmas. To conclude this paper, we remark that our bijection between signed relative derangements and signed skew derangements can be restricted to ordinary permutations. Hence the classical relation (1.1) is a consequence of Theorem 1.3.

Acknowledgments. The authors are grateful to the referee for helpful suggestions. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, the Ministry of Science and Technology, and the National Science Foundation of China.

References

- [1] T. Andreescu and Z. Feng, *A Path to Combinatorics for Undergraduates*, Birkhäuser, Boston, 2004.
- [2] R.A. Brualdi, *Introductory Combinatorics*, 2nd Ed., North-Holland, Amsterdam, 1992.
- [3] W.Y.C. Chen, The skew, relative, and classical derangements, *Discrete Math.*, 160 (1996), 235–239.
- [4] R. J. Clarke, G.-N. Han, J. Zeng, A combinatorial interpretation of the Seidel generation of q -derangement numbers, *Ann. Combin.*, 4 (1997), 313-327.
- [5] C.-O. Chow, On derangement polynomials of type B , *Séminaire Lotharingien de Combinatorics*, 55 (2006), Artical B55b.
- [6] D. Hanson, K. Seyffarth and J.H. Weston, Matchings, derangements, rencontres, *Math. Magazine*, 56 (1983), 224-229.
- [7] R.P. Stanley, *Enumerative Combinatorics*, vol. 1, Cambridge University Press, Cambridge, 1997.
- [8] E.T.H. Wang, Elementary Problem 2947, *Amer. Math. Monthly*, 89 (1982), 334 (with solutions 92 (1985), 216).