

On the size of minimal unsatisfiable formulas *

Choongbum Lee †

Submitted: Oct 29, 2008; Accepted: Jan 21, 2009; Published: Jan 30, 2009
Mathematics Subject Classification: 05D99(Primary); 05C15, 68R10(Secondary)

Abstract

An unsatisfiable formula is called minimal if it becomes satisfiable whenever any of its clauses are removed. We construct minimal unsatisfiable k -SAT formulas with $\Omega(n^k)$ clauses for $k \geq 3$, thereby negatively answering a question of Rosenfeld. This should be compared to the result of Lovász [Studia Scientiarum Mathematicarum Hungarica 11, 1974, p113-114] which asserts that a critically 3-chromatic k -uniform hypergraph can have at most $\binom{n}{k-1}$ edges.

1 Introduction

Given n boolean variables x_1, \dots, x_n , a *literal* is a variable x_i or its negation \bar{x}_i ($1 \leq i \leq n$). A *clause* is a disjunction of literals and by k -*clause* we denote a clause of size k . A *CNF (Conjunctive Normal Form) formula* is a conjunction of clauses and a k -*SAT formula* is a CNF formula with only k -clauses. Throughout this article *formula* will mean a CNF formula and it will be given as a pair $F = (V, C)$ with variables $V = \{x_1, \dots, x_n\}$ and clauses C as collection of disjunction of literals $V \cup \bar{V}$. A formula is called *satisfiable* if there exists an assignment of values to variables so that the formula becomes true. A formula is called *minimal unsatisfiable* if it is not satisfiable but removing any clause makes it satisfiable.

Satisfiability of a formula is closely related to the 2-colorability of a hypergraph in the following sense. A formula is satisfiable if there is an assignment of values to variables in a way that no clauses have only false literals inside it. Similarly a hypergraph is 2-colorable if there is a way to color the vertices into two colors so that none of the edges become monochromatic. A hypergraph $H = (V, E)$ is called critically 3-chromatic if it is not 2 colorable but the deletion of any edge makes it 2 colorable. In this analogy, minimal unsatisfiable formulas correspond to critically 3-chromatic hypergraphs. Therefore it is

*This research forms part of the Ph.D thesis written by the author under the supervision of Prof. Benny Sudakov.

†Department of Mathematics, UCLA, Los Angeles, CA, 90095. E-mail: choongbum.lee@gmail.com. Research supported in part by Samsung Scholarship.

natural to ask if similar results hold for both problems. In particular, we are interested if the same restriction on the number of clauses (edges, respectively) holds or not.

In the case of lower bounds, roughly the same estimate holds for both formulas and hypergraphs. Seymour [5] used linear algebra method to deduce that a critically 3-chromatic hypergraph $H = (V, E)$ must satisfy $|E| \geq |V|$ if there is no isolated vertex. The corresponding bound for CNF formulas appeared in Aharoni and Linial [1] where they quote an unpublished work of M. Tarsi to prove that minimal unsatisfiable formula $F = (V, C)$ must satisfy $|C| \geq |V| + 1$ if every variable is contained in some clause.

For uniform hypergraphs there are also known upper bound results. Lovász [3] proved that any critically 3-chromatic k -uniform hypergraph has at most $\binom{n}{k-1}$ edges. This result is asymptotically tight, as was shown by Toft [6] who constructed critically 3-chromatic k -uniform hypergraphs with $\Omega(n^{k-1})$ edges.

Motivated by these results, Rosenfeld [4] asked if the analogy also holds for minimal unsatisfiable k -SAT formulas.

Question. *Should minimal unsatisfiable k -SAT formulas have at most $O(n^{k-1})$ clauses?*

It is not difficult to show that this conjecture is true for $k = 2$ and we will give the simple proof of this in section 2. However for k -SAT formulas with $k \geq 3$ we show that surprisingly the answer for the question of Rosenfeld is negative. In section 3 we will construct minimal unsatisfiable k -SAT formulas with $\Omega(n^k)$ clauses.

2 2-SAT formulas

First we give explicit minimal unsatisfiable 2-SAT formulas. Consider the 2-SAT formula $F^{(2)} = (V^{(2)}, C^{(2)})$ where $V^{(2)} = \{y_1, y_2, \dots, y_{2l}\}$ and $C^{(2)} = \{y_i \vee y_{i+1}, \bar{y}_i \vee \bar{y}_{i+1} : i = 1, 2, \dots, 2l - 1\} \cup \{y_1 \vee \bar{y}_{2l}\} \cup \{\bar{y}_1 \vee y_{2l}\}$. $F^{(2)}$ is unsatisfiable because if $y_i = y_{i+1}$ for some i then either $y_i \vee y_{i+1}$ or $\bar{y}_i \vee \bar{y}_{i+1}$ is false and otherwise if $y_i = \bar{y}_{i+1}$ for all $1 \leq i \leq 2l - 1$ then $y_1 = \bar{y}_{2l}$ and this time either $y_1 \vee \bar{y}_{2l}$ or $\bar{y}_1 \vee y_{2l}$ will become false. To prove that $F^{(2)}$ is minimal unsatisfiable, we only check that deleting $y_1 \vee y_2$ or $y_1 \vee \bar{y}_{2l}$ makes the new formula satisfiable as other clauses can be checked similarly. In each case, the assignment of $(y_1 = y_2 = \text{false}, y_3 = \dots = y_{2l-1} = \text{true}, y_4 = \dots = y_{2l} = \text{false})$ and $(y_1 = y_3 = \dots = y_{2l-1} = \text{false}, y_2 = y_4 = \dots = y_{2l} = \text{true})$ will make the remaining clauses true.

Next we prove the linear upper bound of number of clauses in minimal unsatisfiable 2-SAT formulas.

Proposition 1. *Minimal unsatisfiable 2-SAT formulas have at most $4n - 2$ clauses.*

Proof. Given a minimal unsatisfiable 2-SAT formula $F = (V, C)$, let's consider the *implication graph* D of this 2-SAT formula which is the directed graph D over the vertices $V \cup \bar{V}$ with two directed edges corresponding to each clause $z_1 \vee z_2 \in C$ given as $\bar{z}_1 \rightarrow z_2$ and $\bar{z}_2 \rightarrow z_1$. Aspvall, Plass and Tarjan [2] proved that 2-SAT is unsatisfiable if and only if its implication graph has a strongly connected component which contains both x_i and

\bar{x}_i for some index i . Therefore the unsatisfiability of F implies the existence of directed path from x_i to \bar{x}_i and from \bar{x}_i and x_i in D for some index i . Now observe that the minimality of F forces every clause $z_1 \vee z_2 \in C$ to have at least one of its corresponding edge $\bar{z}_1 \rightarrow z_2$ or $\bar{z}_2 \rightarrow z_1$ in these directed paths. As otherwise deleting the clause will not change the unsatisfiability of F (because it still contains both directed paths). Since D has $2n$ vertices, there can be at most $4n - 2$ edges in the two directed paths. Therefore $|C| \leq 4n - 2$. \square

3 k -SAT formulas

In this section we construct minimal unsatisfiable k -SAT formulas on n variables with $\Omega(n^k)$ clauses. For simplicity we describe in details the construction of 3-SAT formulas only. This construction can be easily generalized for all k . Informally, start with a minimal unsatisfiable “almost” 3-SAT formula with $\Omega(n^3)$ clauses where “almost” means that only a small number of clauses is not of size 3. Then transform this formula into a “genuine” 3-SAT formula by replacing the clauses of size greater than 3 by 3-clauses while keeping the minimal unsatisfiable property. During the process the number of variables will not increase too much and therefore we will end up with a 3-SAT formula that we have promised. Now we should make it into a formal argument.

The following lemma will allow us to change the size of a clause in the formula. This lemma is a modified version of Theorem 1 and 4 in [6] which were originally used by Toft to construct k -uniform hypergraphs with $\Omega(n^{k-1})$ edges.

Let $F_X = (V_X, C_X), F_Y = (V_Y, C_Y)$ be formulas with disjoint sets of variables (that is, $V_X \cap V_Y = \emptyset$) and $c_0 = z_1 \vee z_2 \vee \dots \vee z_k \in C_X$ be a k -clause of F_X where $k \leq |C_Y|$. For an arbitrary surjective map h from C_Y to $\{z_1, z_2, \dots, z_k\}$, let the formula $F_Z = (V_Z, C_Z)$ be as following.

- $V_Z = V_X \cup V_Y \quad C_Z = (C_X \setminus \{c_0\}) \cup \{c_y \vee h(c_y) : c_y \in C_Y\}$

Lemma 2. *If F_X and F_Y are minimal unsatisfiable formulas, then F_Z constructed as above is also a minimal unsatisfiable formula.*

Proof. Let’s first show that F_Z is unsatisfiable. For arbitrary values of V_X there must exist a clause $c_x \in C_X$ which is false. If $c_x \neq c_0$ then we are done as $c_x \in C_Z$ so assume that $c_x = c_0$. Since every literal $x \in c_0$ is false, a clause of the form $c_y \vee h(c_y)$ is true if and only if c_y is true. But F_Y is unsatisfiable so there must exist a clause c_y which is false and therefore F_Z is unsatisfiable.

Next we prove that removing any clause $c_z \in C_Z$ makes F_Z satisfiable. First assume that $c_z \in C_X \setminus \{c_0\}$. Then give values to V_X so that every clause in C_X except c_z is satisfied. Since $c_z \neq c_0$, there must exist a literal $x \in c_0$ which is true. Pick a clause $c' \in h^{-1}(x) \subset C_Y$ ($h^{-1}(x)$ is non-empty because h is surjective) and give V_Y the values which make every clause except c' in C_Y true. Observe that every clause in $C_X \setminus \{c_0\}$ except c_z is true by values of V_X and every clause in $\{c_y \vee h(c_y) : c_y \in C_Y\}$ is true either by values of V_Y or the literal x . Now assume that $c_z = c' \vee x \in \{c_y \vee h(c_y) : c_y \in C_Y\}$ and

give V_X the values which make every clause except c_0 true and give V_Y the values which makes every clause except c' true. This assignment of values will make every clause but $c_z \in C_Z$ true and thus we are done. \square

Next step is to construct an “almost” 3-SAT formula with many clauses. Let $V_0 = \{x_1, x_2, \dots, x_{6m}\}$ and look at the formula $F_0 = (V_0, C_0)$ with clauses given as,

$$\bullet C_0 = \{x_{i_1} \vee x_{i_2} \vee x_{i_3} : 1 \leq i_1 \leq 2m, 2m + 1 \leq i_2 \leq 4m, 4m + 1 \leq i_3 \leq 6m\} \\ \cup \{\bar{x}_1 \vee \bar{x}_2 \vee \dots, \bar{x}_{2m}\} \cup \{\bar{x}_{2m+1} \vee \dots \vee \bar{x}_{4m}\} \cup \{\bar{x}_{4m+1} \vee \dots \vee \bar{x}_{6m}\}$$

Informally, partition the variables V into three equal parts V_1, V_2, V_3 and consider every clauses $x_1 \vee x_2 \vee x_3$ with $x_i \in V_i$ and add three more clauses $\bar{V}_1, \bar{V}_2, \bar{V}_3$. Note that this formula contains $(2m)^3 + 3$ clauses.

Claim 3. F_0 is a minimal unsatisfiable formula.

Proof. Let's first prove that F_0 is unsatisfiable. Assume that the three clauses $\bar{x}_1 \vee \bar{x}_2 \vee \dots \vee \bar{x}_{2m}$, $\bar{x}_{2m+1} \vee \dots \vee \bar{x}_{4m}$, $\bar{x}_{4m+1} \vee \dots \vee \bar{x}_{6m}$ are all true. Then there must exist $1 \leq i_1 \leq 2m, 2m + 1 \leq i_2 \leq 4m, 4m + 1 \leq i_3 \leq 6m$ such that $x_{i_1} = x_{i_2} = x_{i_3} = \text{false}$. But this will make the clause $x_{i_1} \vee x_{i_2} \vee x_{i_3}$ false. Therefore F_0 is unsatisfiable.

Now assume that we remove a clause c . If $c = x_{i_1} \vee x_{i_2} \vee x_{i_3}$ for some i_1, i_2, i_3 then assigning $x_{i_1} = x_{i_2} = x_{i_3} = \text{false}$ and everything else true will make the remaining part satisfiable. On the other hand if $c = \bar{x}_1 \vee \bar{x}_2 \vee \dots \vee \bar{x}_{2m}$ then assigning $x_1 = x_2 = \dots = x_{2m} = \text{true}$ and everything else false will make the remaining part satisfiable. Similar assignment will work for clauses $\bar{x}_{2m+1} \vee \dots \vee \bar{x}_{4m}$ and $\bar{x}_{4m+1} \vee \dots \vee \bar{x}_{6m}$. \square

Construction

Note that the formula F_0 is “almost” a 3-SAT formula in the sense that there are only three clauses whose size is not 3. Use Lemma 2 with $F_X = F_0$, $c_0 = \bar{x}_1 \vee \bar{x}_2 \vee \dots \vee \bar{x}_{2m} \in C_0$ and $F_Y = F^{(2)}$ where $F^{(2)}$ is a minimal unsatisfiable 2-SAT formula with m variables and $2m$ clauses as constructed in section 2. The obtained formula F_1 is a minimal unsatisfiable formula over $6m + m = 7m$ variables and has only two clauses whose size are not 3. (All new clauses are 3-clauses.)

Repeat the same process with the remaining two $2m$ -clauses to obtain a minimal unsatisfiable formula F_2 whose every clause has size 3 i.e. F_2 is a 3-SAT formula over $n = 9m$ variables. Note that it still contains the original 3-clauses $\{x_{i_1} \vee x_{i_2} \vee x_{i_3} : 1 \leq i_1 \leq 2m, 2m + 1 \leq i_2 \leq 4m, 4m + 1 \leq i_3 \leq 6m\}$. There are $8m^3 = (\frac{2}{9}n)^3$ such clauses and therefore this 3-SAT formula F_2 contains $\Omega(n^3)$ clauses.

For $k \geq 4$, minimal unsatisfiable k -SAT formulas with $\Omega(n^k)$ clauses can be constructed similarly. Use $F_0^{(k)} = (V_0^{(k)}, C_0^{(k)})$ where,

$$\bullet V_0^{(k)} = \{x_1, x_2, \dots, x_{mk}\} \quad (m = \lfloor \frac{n}{k} \rfloor)$$

$$\bullet C_0^{(k)} = \{x_{i_1} \vee x_{i_2} \vee \dots \vee x_{i_k} : (t-1)m + 1 \leq i_t \leq tm, 1 \leq t \leq k\} \\ \cup \left(\bigcup_{s=1}^k \{\bar{x}_{(s-1)m+1} \vee \bar{x}_{(s-1)m+2} \vee \dots \vee \bar{x}_{sm}\} \right)$$

By the same process as above one can verify that $F_0^{(k)}$ is minimal unsatisfiable. Then replace the m -clauses by k -clauses using Lemma 2 and minimal unsatisfiable $(k-1)$ -SAT formulas. The final formula will be a minimal unsatisfiable k -SAT formula with $\Omega(n^k)$ clauses. Details are omitted.

Concluding remarks

- Toft [6] also constructed k -color critical r -uniform hypergraphs ($k \geq 4, r \geq 2$) with $\Omega(n^r)$ edges. Since 3-color critical r -uniform hypergraphs can have at most $O(n^{r-1})$ edges (Lovasz [3]), we can observe an interesting jump from 3-color critical uniform hypergraphs to k -color critical uniform hypergraphs ($k \geq 4$). A similar phenomena, namely that minimal unsatisfiable 2-SAT formulas have $O(n)$ clauses but there are minimal unsatisfiable k -SAT formulas with $\Omega(n^k)$ clauses ($k \geq 3$), occurs also in the case of formulas. It would be interesting to see a more direct connection.
- It was pointed out by the referee that based on the case $k = 3, 4, 5$ of the above construction one can obtain a richer family of minimal unsatisfiable k -SAT formulas ($k \geq 6$) with $\Omega(n^k)$ clauses as follows. Let $F_1 = (V_1, C_1)$ and $F_2 = (V_2, C_2)$ be a k_1 -SAT and a k_2 -SAT minimal unsatisfiable formula respectively, on distinct sets of variables. Then the formula $F^* = (V_1 \cup V_2, \{c_1 \vee c_2 : c_1 \in C_1, c_2 \in C_2\})$ is a $(k_1 + k_2)$ -SAT minimal unsatisfiable formula on $|V_1| + |V_2|$ variables and with $|C_1| \times |C_2|$ clauses. We will omit the simple proof of this fact. It is worth to note that the density of clauses in a k -SAT obtained from the original and this construction is both $O(1/k^k)$.

Acknowledgement. I gratefully thank Benny Sudakov for his advice and guidance and the anonymous referee for the careful reading and valuable suggestions. I am also thankful to Po-Shen Loh and Boris Bukh for the fruitful discussions.

References

- [1] R. Aharoni, N. Linial, *Minimal non-two-colorable hypergraphs and minimal unsatisfiable formulas*, Journal of Combinatorial Theory, Series A 43, 1986, p196-204
- [2] B. Aspvall, M. F. Plass, R. E. Tarjan, *A linear-time algorithm for testing the truth of certain quantified boolean formulas*, Information Processing Letters 8(3), 1979, p121-123
- [3] L. Lovász, *chromatic number of hypergraphs and linear algebra*, Studia Scientiarum Mathematicarum Hungarica 11, 1974, p113-114
- [4] M. Rosenfeld, *private communication*
- [5] P. D. Seymour, *On the two-colouring of hypergraphs*, Quart. J. Math. Oxford 25, 1974, p303-312
- [6] B. Toft, *On Colour-critical hypergraphs*, Colloquia Mathematica Societatis Janos Bolyai 10, 1973, p1445-1457