

Triviality of the 2D stochastic Allen-Cahn equation*

Martin Hairer[†] Marc D. Ryser[‡] Hendrik Weber[§]

Abstract

We consider the stochastic Allen-Cahn equation driven by mollified space-time white noise. We show that, as the mollifier is removed, the solutions converge weakly to 0, independently of the initial condition. If the intensity of the noise simultaneously converges to 0 at a sufficiently fast rate, then the solutions converge to those of the deterministic equation. At the critical rate, the limiting solution is still deterministic, but it exhibits an additional damping term.

Keywords: SPDEs; Allen-Cahn equation; white noise; stochastic quantisation.

AMS MSC 2010: 60H15; 81T08.

Submitted to EJP on January 14, 2012, final version accepted on May 29, 2012.

Supersedes arXiv:1201.3089v1.

1 Introduction

We consider the following evolution equation on the two-dimensional torus \mathbb{T}^2 :

$$du = (\Delta u + u - u^3) dt + \sigma dW, \quad u(0) = u^0. \quad (\Phi)$$

Here u^0 is a suitably regular initial condition, σ a positive constant, and W an $L^2(\mathbb{T}^2)$ -valued cylindrical Wiener process defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. In other words, at least at a formal level, $\frac{dW}{dt}$ is space-time white noise.

This equation and variants thereof have a long history. The deterministic part of the equation is the L^2 gradient flow of the Ginzburg-Landau free energy functional

$$\int_{\mathbb{T}^2} \left(\frac{1}{2} |\nabla u(x)|^2 + V(u(x)) \right) dx,$$

with the potential energy V given by the standard double-well function $V(u) = \frac{1}{4}(u^2 - 1)^2$, see [14]. This provides a phenomenological model for the evolution of an order parameter describing phase coexistence in a system without preservation of mass. At large scales,

*Supported by the Institute of Mathematical Statistics (IMS) and the Bernoulli Society.

[†]University of Warwick, UK. E-mail: m.hairer@warwick.ac.uk

[‡]Duke University, USA. E-mail: ryser@math.duke.edu

[§]University of Warwick, UK. E-mail: hendrik.weber@warwick.ac.uk

the dynamic of phase boundaries is known to converge to the mean curvature flow [1, 9, 12].

The noise term σdW accounts for thermal fluctuations at positive temperature. On a formal level the choice of space-time white noise is natural, because it satisfies the right fluctuation-dissipation relation. At least for finite-dimensional gradient flows it is natural to take the bilinear form that determines the mechanism of energy dissipation as covariance of the noise, as this guarantees the invariance of the right Gibbs measure under the dynamics. Naively extending this observation to the current infinite dimensional context yields (Φ) .

White noise driven equations such as (Φ) are known to be ill-posed in space-dimension $d \geq 2$ [17, 8]. Actually, the linearised version of (Φ) (simply remove the term u^3) admits only distribution-valued solutions for $d \geq 2$. For any $\kappa > 0$ these solutions take values in the Sobolev space $H^{\frac{2-d}{2}-\kappa}$, but they do not take values in $H^{\frac{2-d}{2}}$. In general, it is impossible to apply nonlinear functions to elements of these spaces and the standard approach to construct solutions of (Φ) [8, 11] fails.

In the present article, we introduce a cutoff at spatial lengths of order ε and we study the limit as $\varepsilon \rightarrow 0$ for finite noise strength for (Φ) . More precisely, we set

$$W_\varepsilon(t) = \sum_{|k| \leq 1/\varepsilon} e_k \beta_k(t), \quad \varepsilon > 0,$$

where $\{e_k\}_{k \in \mathbb{Z}^2}$ is the Fourier basis on \mathbb{T}^2 , and $\{\beta_k\}_{k \in \mathbb{Z}^2}$ are complex Brownian motions that are i.i.d. except for the reality condition $\bar{\beta}_k = \beta_{-k}$. We thus consider

$$du_\varepsilon = (\Delta u_\varepsilon + u_\varepsilon - u_\varepsilon^3) dt + \sigma(\varepsilon) dW_\varepsilon, \quad u_\varepsilon(0) = u^0, \tag{\Phi_\varepsilon}$$

and study the weak limit of u_ε as $\varepsilon \rightarrow 0$.

The main result of this article can loosely be formulated as follows (a precise statement will be given in Theorems 2.1 and 2.2 below):

Theorem 1.1. *Let σ be bounded and such that $\lim_{\varepsilon \rightarrow 0} \sigma^2(\varepsilon) \log 1/\varepsilon = \lambda^2 \in [0, +\infty]$. If $\lambda^2 = +\infty$, then u_ε converges weakly to 0, in probability. Otherwise, it converges weakly in probability to the solution w_λ of*

$$\partial_t w_\lambda = \Delta w_\lambda - \left(\frac{3}{8\pi} \lambda^2 - 1\right) w_\lambda - w_\lambda^3, \quad w_\lambda(0) = u^0. \tag{\Psi_\lambda}$$

Remark 1.2. *The result for constant σ was conjectured in [16], based on numerical simulations.*

Remark 1.3. *The borderline case $\lambda \neq 0$ is particularly interesting as it provides an example of stochastic damping: in the limit as $\varepsilon \rightarrow 0$, the stochastic forcing is converted into an additional deterministic damping term, $-\frac{3}{8\pi} \lambda^2 w_\lambda$, to the Allen-Cahn equation. In particular, if $\lambda^2 > \frac{8\pi}{3}$, the zero-solution becomes globally attracting.*

Remark 1.4. *Recently, there has been a lot of interest in (Φ) in the regime where the noise is small [13, 3, 5]. There, the authors studied (Φ) in arbitrary space dimension on the level of large deviation theory. As in (Φ_ε) they consider a modified version of (Φ) where the noise term dW is replaced by a noise term dW_ε with a finite spatial correlation length ε . For this modified equation, solutions can be constructed in a standard way and a large deviation principle à la Freidlin-Wentzell can be obtained. One can then show that the rate functionals converge as $\varepsilon \rightarrow 0$. The large deviation principle however is not uniform in ε ; this procedure corresponds to taking the amplitude of the noise much smaller than ε . The results obtained in this article quantify how small the noise should be as a function of ε in order for the solutions of (Φ) to be close to the deterministic equation.*

Remark 1.5. *We believe that the weak convergence to 0 as $\varepsilon \rightarrow 0$ actually holds for a much larger class of potentials. Actually, one would expect it to be true whenever $\lim_{|u| \rightarrow \infty} V''(u) = +\infty$. The proof given in this article does however depend crucially on the fact that $V(u) \sim u^4$ for large values of u .*

The main tools used in our proofs are provided by the theory of stochastic quantisation. Actually, in the context of Euclidean Quantum Field Theory the question of existence of the formal invariant measure of (Φ) has been treated in the seventies (see e.g. [10]). Then, it had been observed that this measure, the so called Φ_2^4 field, can be defined, but only if a logarithmically diverging lower order term is subtracted. The corresponding stochastic dynamical system (i.e. the renormalised version of (Φ)) has also been constructed [15, 2, 6]. Note that although this renormalised equation,

$$du = (\Delta u + u - :u^3:) dt + \sigma dW ,$$

formally resembles (Φ) it does not have a natural interpretation as a phase field model.

Our main argument is a modification of the construction provided in [6]. We present here a brief heuristic argument for the case $\sigma \equiv 1$. First, let $C_\varepsilon > 1$ and add and subtract the term $C_\varepsilon u_\varepsilon$ to (Φ_ε) to get

$$du_\varepsilon = (\Delta u_\varepsilon - (C_\varepsilon - 1)u_\varepsilon - u_\varepsilon(u_\varepsilon^2 - C_\varepsilon)) dt + \sigma dW_\varepsilon . \tag{1.1}$$

The key idea is to choose C_ε in such a way that, for small values of ε , the term $u_\varepsilon(u_\varepsilon^2 - C_\varepsilon)$ is equal to the Wick product $:u_\varepsilon^3:$ with respect to the Gaussian structure given by the invariant measure of the linearised system (which itself depends on C_ε). Since, given the results in [6], one would expect $:u_\varepsilon^3:$ to at least remain bounded as $\varepsilon \rightarrow 0$, it is not surprising that the additional strong damping term $-C_\varepsilon u_\varepsilon$ causes the solution to vanish in the limit.

2 Notations and Main Result

In order to formulate our results, we first introduce the class of Besov spaces that we will work with. As in [6] we choose to work in Besov spaces, because they satisfy the right multiplicative inequalities (see Lemma A.2). Denote by (\cdot, \cdot) the L^2 inner product, and by $\{e_k(x) = \frac{1}{2\pi} e^{ikx}\}_{k \in \mathbb{Z}^2}$ the corresponding orthonormal Fourier basis. Throughout the article, we work with periodic Besov spaces $\mathcal{B}_{p,r}^s(\mathbb{T}^2)$, where $p, r \geq 1$ and $s \in \mathbb{R}$. These spaces are defined as the closure of $C^\infty(\mathbb{T}^2)$ under the norm

$$\|u\|_{\mathcal{B}_{p,r}^s(\mathbb{T}^2)} := \left(\sum_{q=0}^{\infty} 2^{qrs} \|\Delta_q u\|_{L^p(\mathbb{T}^2)}^r \right)^{1/r} , \tag{2.1}$$

where the Δ_q are the Littlewood-Paley projection operators given by $\Delta_0 u = (e_0, u) e_0$ and

$$\Delta_q u = \sum_{2^{q-1} \leq |k| < 2^q} (e_k, u) e_k, \quad q \geq 1.$$

Regarding the exponents appearing in these Besov spaces, we will restrict ourselves throughout this article to exponents p, r and s such that

$$p \geq 4, \quad r \geq 1, \quad -\frac{2}{7p} < s < 0 . \tag{2.2}$$

We now reformulate Theorem 1.1 more precisely. The case $\lambda^2 = +\infty$ is given by the following:

Theorem 2.1. Assume $u^0 \in \mathcal{B}_{p,r}^s$ such that (2.2) holds. Then for all $\varepsilon > 0$ and $T > 0$, there exists a unique mild solution u_ε . If $\sigma(\varepsilon)$ is bounded uniformly in ε and satisfies $\lim_{\varepsilon \rightarrow 0} \sigma^2(\varepsilon) \log(1/\varepsilon) = +\infty$ then, for all $\delta \in (0, T)$, $\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon\|_{\mathcal{C}([\delta, T]; \mathcal{B}_{p,r}^s)} = 0$ in probability.

On the other hand, when $\sigma^2(\varepsilon) \log(1/\varepsilon)$ converges to a finite limit, we have

Theorem 2.2. Assume $u^0 \in \mathcal{B}_{p,r}^s$ such that (2.2) holds. If $\lim_{\varepsilon \rightarrow 0} \sigma^2(\varepsilon) \log(1/\varepsilon) = \lambda^2 \in \mathbb{R}$, then $\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - w_\lambda\|_{\mathcal{C}([0, T]; \mathcal{B}_{p,r}^s)} = 0$ in probability, where w_λ is the unique solution to (Ψ_λ) .

Remark 2.3. If σ decays sufficiently fast, for example $\sigma(\varepsilon) \sim \varepsilon^\tau$ for some $\tau > 0$, then the conclusion of Theorem 2.2 actually holds in the space of space-time continuous functions.

To conclude this section, we introduce some concepts borrowed from the theory of stochastic quantization. Since we are not concerned with the dynamics of quantised fields, we only introduce the notions necessary for the proof techniques used below, and refer to [7] for a general introduction to the topic. Consider the linear version of equation (1.1), namely

$$dz_\varepsilon = (\Delta z_\varepsilon - (C_\varepsilon - 1) z_\varepsilon) dt + \sigma(\varepsilon) dW_\varepsilon . \tag{2.3}$$

For $C_\varepsilon > 1$, this equation has a unique invariant measure on $L^2(\mathbb{T}^2)$, which we denote by μ_ε . It is μ_ε that will play the role of the “free field” in the present article. Under μ_ε , the k th Fourier component of z_ε is a centred complex Gaussian random variable with variance $\frac{\sigma^2(\varepsilon)}{2} (C_\varepsilon - 1 + |k|^2)^{-1}$. Furthermore, distinct Fourier components are independent, except for the reality condition $z_\varepsilon(-k) = \overline{z_\varepsilon(k)}$.

As a consequence of translation invariance, one has the identity

$$\begin{aligned} D_\varepsilon^2 &:= \int_{L^2} |\phi_\varepsilon(x)|^2 \mu_\varepsilon(d\phi_\varepsilon) = \left(\frac{1}{2\pi}\right)^2 \int_{L^2} \|\phi_\varepsilon\|_{L^2}^2 \mu_\varepsilon(d\phi_\varepsilon) \\ &= \frac{1}{8\pi^2} \sum_{|k| \leq 1/\varepsilon} \frac{\sigma^2(\varepsilon)}{C_\varepsilon - 1 + |k|^2} . \end{aligned} \tag{2.4}$$

We then define the Wick powers of any field u_ε with respect to the Gaussian structure given by μ_ε by

$$:u_\varepsilon^n: = D_\varepsilon^n H_n(u_\varepsilon / D_\varepsilon) ,$$

where H_n denotes the n th Hermite polynomial. In this article, we will only ever use the Wick powers for $n \leq 3$, for which one has the identities

$$:u_\varepsilon^1: = u_\varepsilon , \quad :u_\varepsilon^2: = u_\varepsilon^2 - D_\varepsilon^2 , \quad :u_\varepsilon^3: = u_\varepsilon^3 - 3 D_\varepsilon^2 u_\varepsilon . \tag{2.5}$$

From now on, whenever we use the notation $:u_\varepsilon^n:$, (2.5) is what we refer to. For any two expressions A and B depending on ε , we will throughout this article use the notation $A \lesssim B$ to mean that there exists a constant C independent of ε (and possibly of other relevant parameters clear from the respective contexts) such that $A \leq C B$.

3 Trivial limit for strong noise

In this section, we provide the proof of Theorem 2.1. First, in Subsection 3.1, we explain the “correct” choice of the renormalization constant C_ε in (1.1). In Subsection 3.2, we then obtain bounds on the linearised equation, as well as its Wick powers. Finally, in Subsection 3.3, we obtain a bound on the remainder and we combine these results in order to conclude.

3.1 Fixing the renormalization constant

For $C_\varepsilon > 1$, we rewrite (Φ_ε) as

$$du_\varepsilon = (A_\varepsilon u_\varepsilon - u_\varepsilon(u_\varepsilon^2 - C_\varepsilon)) dt + \sigma(\varepsilon)dW_\varepsilon, \tag{3.1}$$

where the linear operator A_ε is given by $A_\varepsilon = \Delta - (C_\varepsilon - 1)$. Motivated by the heuristic arguments provided in Section 1, the goal of this section is to determine C_ε in such a way that the nonlinear term $u_\varepsilon(u_\varepsilon^2 - C_\varepsilon)$ is equal to the Wick product $:u_\varepsilon^3:$. It then follows from (2.4) and (2.5) that C_ε is implicitly determined by the equation

$$C_\varepsilon = 3D_\varepsilon^2 = \frac{3}{8\pi^2} \sum_{|k| \leq 1/\varepsilon} \frac{\sigma^2(\varepsilon)}{C_\varepsilon - 1 + |k|^2}. \tag{3.2}$$

To describe the behavior of the solution to (3.2), we shall use the notation $A_\varepsilon \sim B_\varepsilon$ to mean $\lim_{\varepsilon \rightarrow 0} A_\varepsilon/B_\varepsilon = 1$.

Lemma 3.1. *For any values of the parameters, equation (3.2) has a unique solution $C_\varepsilon > 1$. If σ is uniformly bounded and such that $\lim_{\varepsilon \rightarrow 0} \sigma^2(\varepsilon) \log(1/\varepsilon) = \infty$, then one has*

$$C_\varepsilon \sim \frac{3}{4\pi} \sigma^2(\varepsilon) \log \frac{1}{\varepsilon}. \tag{3.3}$$

In particular, $\lim_{\varepsilon \rightarrow 0} C_\varepsilon = +\infty$.

Before we proceed to the proof of this result, we state the following very useful result:

Lemma 3.2. *Let $a, R \geq 1$. Then there exists a constant C such that the bound*

$$\left| \sum_{|k| \leq R} \frac{1}{a + |k|^2} - \pi \log \left(1 + \frac{R^2}{a} \right) \right| \leq \frac{C}{\sqrt{a}} \left(1 \wedge \frac{R}{\sqrt{a}} \right), \tag{3.4}$$

holds. Here, the sum goes over elements $k \in \mathbb{Z}^2$.

Proof. The second expression on the left is nothing but $\int_{|k| \leq R} \frac{dk}{a + |k|^2}$, so we want to bound the difference between the sum and the integral. Using the monotonicity and positivity of the function $x \mapsto \frac{1}{a + x^2}$ and restricting ourselves to one quadrant, we see that one has the bounds

$$\sum_{\substack{|k| \leq R \\ k_i > 0}} \frac{1}{a + |k|^2} \leq \frac{1}{4} \int_{|k| \leq R} \frac{dk}{a + |k|^2} \leq \sum_{\substack{|k| \leq R \\ k_i \geq 0}} \frac{1}{a + |k|^2}.$$

As a consequence, the required error is bounded by

$$4 \sum_{k=0}^{\lfloor R \rfloor} \frac{1}{a + k^2} \leq \frac{4}{a} + 4 \int_0^R \frac{dx}{a + x^2}.$$

The required bound follows at once, using the fact that a and R are bounded away from 0 by assumption. □

Proof of Lemma 3.1. Since the right hand side in (3.2) decreases from ∞ down to 0 as the left hand side grows from 1 to ∞ , it follows immediately that (3.2) always has a unique solution $C_\varepsilon > 1$.

Since, by Lemma 3.2, one has $\sum_{|k| \leq \frac{1}{\varepsilon}} \frac{1}{1 + |k|^2} \sim 2\pi \log \frac{1}{\varepsilon}$ and since by assumption $\sigma^2(\varepsilon) \log \frac{1}{\varepsilon} \rightarrow \infty$, there exists ε_0 such that

$$\frac{3}{8\pi^2} \sum_{|k| \leq 1/\varepsilon} \frac{\sigma^2(\varepsilon)}{1 + |k|^2} \geq 2, \tag{3.5}$$

for all $\varepsilon < \varepsilon_0$. As a consequence, we have $C_\varepsilon \geq 2$ for such values of ε , and we will use this bound from now on. On the other hand, if we know that $C_\varepsilon \geq 2$, then C_ε is bounded from above by the left hand side of (3.5), so that

$$C_\varepsilon \leq K\sigma^2(\varepsilon) \log \frac{1}{\varepsilon}, \tag{3.6}$$

for some constant K and for ε small enough.

It now follows from Lemma 3.2 that

$$C_\varepsilon = \frac{3\sigma^2(\varepsilon)}{8\pi} \log \left(1 + \frac{1}{\varepsilon^2(C_\varepsilon - 1)} \right) + R_\varepsilon, \tag{3.7}$$

for some remainder R_ε which is uniformly bounded as $\varepsilon \rightarrow 0$. Since, by (3.6), the first term on the right hand side goes to ∞ , this shows that R_ε is negligible in (3.7), so that

$$C_\varepsilon \sim \frac{3\sigma^2(\varepsilon)}{8\pi} \log \left(\frac{1}{\varepsilon^2 C_\varepsilon} \right) = \frac{3\sigma^2(\varepsilon)}{8\pi} \left(\log \frac{1}{\varepsilon^2} - \log C_\varepsilon \right).$$

Since C_ε is negligible with respect to $\frac{1}{\varepsilon^2}$ by (3.6), the claim follows. □

3.2 Bounds on the linearised equation

We split the solution to (3.1) into two parts by introducing the stochastic convolution

$$z_\varepsilon(t) := \sigma(\varepsilon) \int_{-\infty}^t e^{(t-s)A_\varepsilon} dW_\varepsilon(s), \tag{3.8}$$

and performing the change of variables $v_\varepsilon(t) := u_\varepsilon(t) - z_\varepsilon(t)$. With these notations, v_ε solves

$$\begin{aligned} \partial_t v_\varepsilon &= A_\varepsilon v_\varepsilon - (v_\varepsilon^3 + 3v_\varepsilon^2 z_\varepsilon + 3v_\varepsilon z_\varepsilon^2 + :z_\varepsilon^3:) & (\Phi_\varepsilon^{aux}) \\ v_\varepsilon(0) &= u^0 - z_\varepsilon(0). \end{aligned}$$

We thus split the original problem into two parts: first, we show that the stochastic convolution converges to 0, then we show that the remainder v_ε also converges to 0.

By construction, the stochastic convolution (3.8) is a stationary process and its invariant measure is given by μ_ε . We first establish a general estimate for its renormalized powers $:z_\varepsilon^n:$, which will be useful for bounding v_ε later on. Throughout this section, we assume that $\lim_{\varepsilon \rightarrow 0} \sigma^2(\varepsilon) \log(1/\varepsilon) = \infty$ and that C_ε is given by (3.2). We then have:

Lemma 3.3. *Let $r, k, p \geq 1, s < 0$. Then, for all $n \in \mathbb{N}$, we have*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \|:z_\varepsilon^n:\|_{\mathcal{B}_{p,r}^s}^k = 0. \tag{3.9}$$

Proof. Following the calculations of the proof of [6, Lemma 3.2], we see that

$$\mathbb{E} \|:z_\varepsilon^n:\|_{\mathcal{B}_{p,r}^s}^k \lesssim \|\gamma_\varepsilon\|_{H^{\beta_n}}^{\frac{kn}{2}}, \tag{3.10}$$

where $\beta_n = 1 + \frac{rks}{2^n p}$ and

$$\gamma_\varepsilon(x) = \sum_{|k| \leq 1/\varepsilon} \frac{\sigma^2(\varepsilon)}{C_\varepsilon - 1 + |k|^2} e_k(x).$$

Since

$$\|\gamma_\varepsilon\|_{H^{\beta_n}}^2 = \sum_{|k| \leq 1/\varepsilon} \frac{\sigma^4(\varepsilon)(1 + |k|^2)^{\beta_n}}{(C_\varepsilon - 1 + |k|^2)^2},$$

and $\beta_n < 1$, the claim follows from the boundedness of σ and the fact that $C_\varepsilon \rightarrow \infty$. □

Corollary 3.4. *Let $n, p, r \geq 1$ and $s < 0$. Then $z_\varepsilon^n : \in L^p([0, T]; \mathcal{B}_{p,r}^s)$ \mathbb{P} -a.s., for all $\varepsilon > 0$. In particular,*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \|z_\varepsilon^n\|_{L^p(0,T;\mathcal{B}_{p,r}^s)} = 0. \tag{3.11}$$

Proof. This follows from the stationarity of z_ε , Fubini’s theorem and Lemma 3.3. \square

We establish now the main result of this subsection.

Proposition 3.5. *Consider the stochastic convolution z_ε defined in (3.8) and let $p, r \geq 1$, $s < 0$ and $T > 0$. Then*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \|z_\varepsilon\|_{\mathcal{C}([0,T];\mathcal{B}_{p,r}^s)} = 0. \tag{3.12}$$

Proof. We begin by decomposing the stochastic convolution into two parts,

$$z_\varepsilon(t) = e^{tA_\varepsilon} z_\varepsilon(0) + \sigma(\varepsilon) \int_0^t e^{(t-s)A_\varepsilon} dW_\varepsilon(s).$$

The bound on the first term follows from Lemma 3.3 and Lemma A.3, so it remains to focus on the second term, which we denote hereafter as $\bar{z}_\varepsilon(t)$. In order to bound it, we use the *factorization method*, see [8, p. 128], as well as [11, p. 47] for a more detailed presentation. Recalling that

$$\int_\sigma^t (t-s)^{\alpha-1} (s-\sigma)^{-\alpha} ds = \frac{\pi}{\sin \pi \alpha},$$

we fix $\alpha \in (0, \frac{1}{2})$ and rewrite \bar{z}_ε as

$$\bar{z}_\varepsilon(t) = \frac{\sin \pi \alpha}{\pi} \int_0^t e^{(t-s)A_\varepsilon} Y_\varepsilon(s) (t-s)^{\alpha-1} ds, \tag{3.13}$$

where

$$Y_\varepsilon(s) := \sigma(\varepsilon) \int_0^s (s-\sigma)^{-\alpha} e^{(s-\sigma)A_\varepsilon} dW_\varepsilon(\sigma).$$

Next, we introduce the mapping $\Gamma_\varepsilon : y \mapsto \Gamma_\varepsilon y$ defined by

$$\Gamma_\varepsilon y(t) := \frac{\sin \pi \alpha}{\pi} \int_0^t e^{(t-s)A_\varepsilon} y(s) (t-s)^{\alpha-1} ds,$$

and show that $\Gamma_\varepsilon : L^q([0, T]; \mathcal{B}_{p,r}^s) \rightarrow \mathcal{C}([0, T]; \mathcal{B}_{p,r}^s)$ is a bounded mapping for $q > 1/\alpha$. First, it is a consequence of the strong continuity of e^{tA_ε} that $\Gamma_\varepsilon y \in \mathcal{C}([0, T]; \mathcal{B}_{p,r}^s)$ for all $y \in \mathcal{C}([0, T]; \mathcal{B}_{p,r}^s)$ such that $y(0) = 0$ [11, p. 48]. Next, observe that $s \mapsto (t-s)^{\alpha-1}$ is in $L^{\bar{q}}([0, t])$ for all $\bar{q} \in [1, (1-\alpha)^{-1})$, and hence we can use Hölder’s inequality to deduce that for all $q > \frac{1}{\alpha}$,

$$\sup_{t \in [0, T]} \|\Gamma_\varepsilon y(t)\|_{\mathcal{B}_{p,r}^s} \lesssim \|y\|_{L^q([0, T]; \mathcal{B}_{p,r}^s)}. \tag{3.14}$$

A standard density argument allows us to conclude that $\Gamma_\varepsilon : L^q([0, T]; \mathcal{B}_{p,r}^s) \rightarrow \mathcal{C}([0, T]; \mathcal{B}_{p,r}^s)$ is indeed a bounded mapping for $q > 1/\alpha$.

To conclude the proof, we assume for the moment that there exist $K_\varepsilon > 0$ such that

$$\sup_{t \in [0, T]} \mathbb{E} \|Y_\varepsilon(t)\|_{\mathcal{B}_{p,r}^s} \leq K_\varepsilon, \quad \lim_{\varepsilon \rightarrow 0} K_\varepsilon = 0. \tag{3.15}$$

From (3.15), it then follows that

$$\mathbb{E} \|Y_\varepsilon\|_{L^q([0,T];\mathcal{B}_{p,r}^s)} \leq \left(T \sup_{t \in [0,T]} \mathbb{E} \|Y_\varepsilon\|_{\mathcal{B}_{p,r}^s}^q \right)^{1/q} \lesssim T^{1/q} K_\varepsilon, \tag{3.16}$$

where the first inequality is due to Jensen’s inequality and Fubini’s theorem, and the second inequality follows from (3.15) in conjunction with Fernique’s theorem. By (3.16), $Y_\varepsilon \in L^q([0,T];\mathcal{B}_{p,r}^s)$ \mathbb{P} -a.s. and hence $\bar{z}_\varepsilon = \Gamma_\varepsilon Y_\varepsilon \in \mathcal{C}([0,T];\mathcal{B}_{p,r}^s)$ \mathbb{P} -a.s. Furthermore, it follows from (3.14)–(3.16) that

$$\mathbb{E} \sup_{t \in [0,T]} \|\bar{z}_\varepsilon(t)\|_{\mathcal{B}_{p,r}^s} \lesssim \mathbb{E} \|Y_\varepsilon\|_{L^q([0,T];\mathcal{B}_{p,r}^s)} \lesssim K_\varepsilon,$$

so that $\|\bar{z}_\varepsilon\|_{\mathcal{C}([0,T];\mathcal{B}_{p,r}^s)} \rightarrow 0$ in probability, as required.

It remains to establish (3.15). By definition of the Besov norm (2.1) and Jensen’s inequality,

$$\mathbb{E} \|Y_\varepsilon(t)\|_{\mathcal{B}_{p,r}^s} \leq \left(\sum_{q=0}^\infty 2^{qrs} \mathbb{E} \|\Delta_q Y_\varepsilon(t)\|_{L^p}^r \right)^{1/r}. \tag{3.17}$$

As a consequence, (3.15) follows if we can show that

$$\mathbb{E} \|\Delta_q Y_\varepsilon(t)\|_{L^p}^p \leq K_\varepsilon 2^{qp\tau}, \tag{3.18}$$

for some $\tau < |s|$ and some $K_\varepsilon \rightarrow 0$.

Fix now $q \in \mathbb{N}$. Thanks to Fubini’s theorem, the Gaussianity of $\Delta_q Y_\varepsilon(t)$, and the independence of its different Fourier components,

$$\begin{aligned} \mathbb{E} \|\Delta_q Y_\varepsilon(t)\|_{L^p}^p &= \int_{\mathbb{T}^2} \mathbb{E} \left| \sum_{2^{q-1} \leq |k| < 2^q} (Y_\varepsilon(t), e_k) e_k(\xi) \right|^p d\xi \\ &\lesssim \int_{\mathbb{T}^2} \left(\mathbb{E} \left| \sum_{2^{q-1} \leq |k| < 2^q} (Y_\varepsilon(t), e_k) e_k(\xi) \right|^2 \right)^{p/2} d\xi \\ &\lesssim \int_{\mathbb{T}^2} \left(\sum_{2^{q-1} \leq |k| < 2^q} \mathbb{E} |(Y_\varepsilon(t), e_k)|^2 \right)^{p/2} d\xi. \end{aligned} \tag{3.19}$$

Itô’s isometry and the definition of A_ε yield

$$\begin{aligned} \mathbb{E} |(Y_\varepsilon(t), e_k)|^2 &\leq \sigma^2(\varepsilon) \left[2 \left(C_\varepsilon - 1 + |k|^2 \right) \right]^{2\alpha-1} \int_0^\infty e^{-\tau} \tau^{-2\alpha} d\tau \\ &\lesssim \sigma^2(\varepsilon) \left(C_\varepsilon - 1 + |k|^2 \right)^{2\alpha-1}, \end{aligned} \tag{3.20}$$

where the last inequality is due to $2\alpha < 1$. Inserting (3.20) back into (3.19) we obtain the bound

$$\begin{aligned} \mathbb{E} \|\Delta_q Y_\varepsilon(t)\|_{L^p}^p &\lesssim \sigma^p(\varepsilon) \left(\sum_{2^{q-1} \leq |k| < 2^q} \left(\frac{1}{C_\varepsilon - 1 + |k|^2} \right)^{1-2\alpha} \right)^{p/2} \\ &\lesssim \sigma^p(\varepsilon) \left(\frac{2^{2q\tau}}{(C_\varepsilon - 1)^\delta} \sum_{2^{q-1} \leq |k| < 2^q} \frac{1}{|k|^{2+2\tau-4\alpha-2\delta}} \right)^{p/2}, \end{aligned}$$

which is valid for all $\tau > 0$ and all $\delta \in (0, 1 - 2\alpha)$. Since we can make both α and δ arbitrarily small, we can in particular choose them in such a way that $2\alpha + \delta < \tau < |s|$, so that the exponent is strictly greater than 2. This implies that the corresponding inverse power of $|k|$ is summable over all k , so that (3.18) is satisfied. \square

3.3 Bounds on the remainder

First, we need a technical lemma for the mapping \mathcal{M}^ε , defined as

$$(\mathcal{M}^\varepsilon y)(t) := e^{tA_\varepsilon} (u^0 - z_\varepsilon(0)) + \int_0^t e^{(t-\tau)A_\varepsilon} \sum_{l=0}^3 a_l y^l(\tau) : z_\varepsilon^{3-l}(\tau) : d\tau, \tag{3.21}$$

where the a_l are some real-valued constants. In order to formulate the results of this section, we introduce the Banach space

$$\mathcal{E}_T := \mathcal{C}([0, T]; \mathcal{B}_{p,r}^s) \cap L^p([0, T]; \mathcal{B}_{p,r}^{\bar{s}}),$$

equipped with the usual maximum norm

$$\|x\|_{\mathcal{E}_T} := \max \left(\|x\|_{\mathcal{C}([0, T]; \mathcal{B}_{p,r}^s)}, \|x\|_{L^p([0, T]; \mathcal{B}_{p,r}^{\bar{s}})} \right). \tag{3.22}$$

Regarding the parameters appearing in \mathcal{E}_T , we shall usually assume that (p, r, s, \bar{s}) satisfy the bounds

$$p \geq 4, \quad r \geq 1, \quad \bar{s} = 2s + \frac{2}{p}, \quad -\frac{2}{7p} < s < 0. \tag{3.23}$$

Lemma 3.6. Fix $\varepsilon > 0, T > 0$, and assume (3.23). Then there exist positive constants δ and K_ε with $\lim_{\varepsilon \rightarrow 0} K_\varepsilon = 0$ such that

$$\begin{aligned} \|\mathcal{M}^\varepsilon y\|_{\mathcal{E}_T} &\leq (1 + K_\varepsilon T^\delta) \|u^0 - z_\varepsilon(0)\|_{\mathcal{B}_{p,r}^s} \\ &\quad + K_\varepsilon T^\delta \sum_{l=0}^3 \| : z_\varepsilon^{3-l} : \|_{L^p([0, T]; \mathcal{B}_{p,r}^s)} \|y\|_{\mathcal{E}_T}^l. \end{aligned} \tag{3.24}$$

Proof. The bound of the first term on the right-hand side of (3.21) is given in Proposition A.4. Next, we split the second term into two parts, $\Omega_\varepsilon^1 + \Omega_\varepsilon^2$, where

$$\begin{aligned} \Omega_\varepsilon^1(t, y) &= \int_0^t e^{(t-\tau)A_\varepsilon} \sum_{l=0}^2 a_l : z_\varepsilon^{3-l}(\tau) : y^l(\tau) d\tau, \\ \Omega_\varepsilon^2(t, y) &= \int_0^t e^{(t-\tau)A_\varepsilon} y^3(\tau) d\tau. \end{aligned}$$

We bound Ω_ε^1 first. Since $((l + 1) - 1)s + 1 - 2/p > 0$ for $l = 0, 1, 2$, we can employ Lemma A.1 to find that there exist $\delta > 0$ and K_ε as in the statement such that

$$\left\| \int_0^t e^{(t-\tau)A_\varepsilon} : z_\varepsilon^{3-l}(\tau) : y^l(\tau) d\tau \right\|_{\mathcal{E}_T} \leq K_\varepsilon T^\delta \| : z_\varepsilon^{3-l} : y^l \|_{L^{p/(l+1)}([0, T]; \mathcal{B}_{p,r}^{(2l+1)s})}.$$

Using Lemma A.2 and adding up the respective contributions yields the terms with $l = 0, 1, 2$ on the right-hand side of (3.24).

We now bound Ω_ε^2 . Since $y \in \mathcal{E}_T$ and $s < \bar{s}$, the embedding $\mathcal{B}_{p,r}^{\bar{s}} \hookrightarrow \mathcal{B}_{p,r}^s$ implies that $y \in L^p([0, T]; \mathcal{B}_{p,r}^s)$. From Lemma A.1 with $n = 3$ and Lemma A.2 with $l = 2$, it follows again that there exist δ and K_ε such that

$$\begin{aligned} \left\| \int_0^t e^{(t-\tau)A_\varepsilon} y^3(\tau) d\tau \right\|_{\mathcal{E}_T} &\leq K_\varepsilon T^\delta \|y y^2\|_{L^{p/3}([0, T]; \mathcal{B}_{p,r}^{5s})} \\ &\leq K_\varepsilon T^\delta \|y\|_{L^p([0, T]; \mathcal{B}_{p,r}^{\bar{s}})}^3, \end{aligned} \tag{3.25}$$

which is the term with $l = 3$ on the right-hand side of (3.24). □

Lemma 3.7. *Let $\varepsilon > 0$, assume (3.23) and consider (Φ_ε^{aux}) with $u^0 \in \mathcal{B}_{p,r}^s$. Then for all $T > 0$, there exists \mathbb{P} -a.s. a unique mild solution $v_\varepsilon \in \mathcal{E}_T$.*

Proof. The existence of unique local solutions to (Φ_ε^{aux}) follows from (3.24) and is shown in detail in [6, Prop. 4.4]. Furthermore, a fixed point argument in a weighted supremum norm shows that $v_\varepsilon(T^*) \in \mathcal{C}(\mathbb{T}^2)$. Since, for $\mathcal{C}(\mathbb{T}^2)$ -valued initial datum, (Φ_ε^{aux}) admits a unique global solution in $\mathcal{C}([0, T]; \mathcal{C}(\mathbb{T}^2)) \cap \mathcal{C}((0, \infty), \mathcal{C}^\infty(\mathbb{T}^2))$, see e.g. [11, Thm. 6.4; Prop. 6.23], the claim follows from the fact that this space is a subspace of \mathcal{E}_T . \square

Before we state the main result of this section, we introduce the Banach space

$$\mathcal{E}_T^\delta := \mathcal{C}([\delta, T]; \mathcal{B}_{p,r}^s) \cap L^p([0, T]; \mathcal{B}_{p,r}^{\bar{s}}), \quad \delta \in [0, T],$$

equipped with the norm $\|x\|_{\mathcal{E}_T^\delta} := \|x\|_{\mathcal{C}([\delta, T]; \mathcal{B}_{p,r}^s)} + \|x\|_{L^p([0, T]; \mathcal{B}_{p,r}^{\bar{s}})}$. With this notation, we have:

Proposition 3.8. *Assume (3.23) and consider the sequence of regularized problems (Φ_ε^{aux}) with fixed initial condition $u^0 \in \mathcal{B}_{p,r}^s$. For all $T > 0$, the unique global solution $v_\varepsilon \in \mathcal{E}_T$ from Lemma 3.7 converges to zero in sense that, for every $\delta \in (0, T)$, $\lim_{\varepsilon \rightarrow 0} \|v_\varepsilon\|_{\mathcal{E}_T^\delta} = 0$ in probability.*

Proof. We introduce the stopping time $\tau_{\varepsilon, \delta}$ as

$$\tau_{\varepsilon, \delta} := T \wedge \inf \left\{ t \geq \delta : \|v_\varepsilon\|_{\mathcal{E}_t^\delta} \geq 1 \right\}, \tag{3.26}$$

with the convention that $\tau_{\varepsilon, \delta} = T$ if the set is empty. Next, we establish the limit

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \|v_\varepsilon\|_{\mathcal{E}_{\tau_{\varepsilon, \delta}}^\delta} = 0. \tag{3.27}$$

Recalling that v_ε solves the fixed point equation $\mathcal{M}^\varepsilon v_\varepsilon = v_\varepsilon$, we can use Lemma 3.6, combined with

$$\sup_{t \in [\delta, T]} \|e^{tA_\varepsilon} (u^0 - z_\varepsilon(0))\|_{\mathcal{B}_{p,r}^s} \leq e^{-\delta C_\varepsilon} \| (u^0 - z_\varepsilon(0)) \|_{\mathcal{B}_{p,r}^s}, \tag{3.28}$$

to show that there exists $\gamma > 0$ and K_ε with $\lim_{\varepsilon \rightarrow 0} K_\varepsilon = 0$ such that

$$\begin{aligned} \|v_\varepsilon\|_{\mathcal{E}_{\tau_{\varepsilon, \delta}}^\delta} &\leq K_\varepsilon (1 + T^\gamma) \|u^0 - z_\varepsilon(0)\|_{\mathcal{B}_{p,r}^s} \\ &\quad + K_\varepsilon T^\gamma \sum_{l=0}^3 \|v_\varepsilon\|_{\mathcal{E}_{\tau_{\varepsilon, \delta}}^\delta}^l \|z_\varepsilon^{3-l}\|_{L^p([0, \tau_{\varepsilon, \delta}^\delta]; \mathcal{B}_{p,r}^s)}. \end{aligned}$$

Since $\|v_\varepsilon\|_{\mathcal{E}_{\tau_{\varepsilon, \delta}}^\delta} \leq 1$ by construction, the claim (3.27) then follows from Lemma 3.3 and Corollary 3.4. Since, by the definition of $\tau_{\varepsilon, \delta}$, this implies that $\lim_{\varepsilon \rightarrow 0} \mathbb{P}(\tau_{\varepsilon, \delta} < T) = 0$, the claim follows. \square

Proof of Theorem 2.1. Since $u_\varepsilon = z_\varepsilon + v_\varepsilon$, the claim follows from Propositions 3.5 and 3.8, in conjunction with the embedding $\mathcal{B}_{\bar{p}, r}^{\bar{s}} \hookrightarrow \mathcal{B}_{p, r}^s$, which holds if $\bar{s} \geq s$ and $\bar{p} \geq p$. \square

4 Deterministic limit for weak noise

In this section, we give the proof of Theorem 2.2. The technique of proof is almost identical to the previous section, but we define objects in a slightly different way. This time, we define an operator $A = \Delta - 1$, and we set

$$z_\varepsilon(t) := \sigma(\varepsilon) \int_{-\infty}^t e^{(t-s)A} dW_\varepsilon(s). \tag{4.1}$$

We furthermore define all of our Wick products with respect to the law μ_ε of z_ε , so that all throughout this section (2.5) holds, but with D_ε given by

$$D_\varepsilon^2 = \frac{1}{8\pi^2} \sum_{|k| \leq 1/\varepsilon} \frac{\sigma^2(\varepsilon)}{1 + |k|^2}.$$

Note that, by Lemma 3.2, one has

$$\lim_{\varepsilon \rightarrow 0} D_\varepsilon^2 = \frac{\lambda^2}{8\pi}.$$

As before, we rewrite the solution to (Φ_ε) as $u_\varepsilon = v_\varepsilon + z_\varepsilon$, where v_ε is solution to

$$\partial_t v_\varepsilon = Av_\varepsilon + (2 - 3D_\varepsilon^2)(v_\varepsilon + z_\varepsilon) + \sum_{l=0}^3 a_l v_\varepsilon^l :z_\varepsilon^{3-l}:, \tag{4.2}$$

with initial condition $v_\varepsilon(0) = u^0 - z_\varepsilon(0)$ and suitable constants a_l .

Note first that one has the following result:

Proposition 4.1. *Let z_ε be defined as in (4.1). Then, for every $T > 0$ and every $n > 0$, the limits*

$$\lim_{\varepsilon \rightarrow 0} \|z_\varepsilon\|_{\mathcal{C}([0,T]; \mathcal{B}_{p,r}^s)} = 0, \quad \lim_{\varepsilon \rightarrow 0} \|:z_\varepsilon^n:\|_{L^p([0,T]; \mathcal{B}_{p,r}^s)} = 0,$$

hold in probability.

Proof. It follows from [6, Lem. 3.2] that

$$\mathbb{E} \|:z_\varepsilon^n:\|_{\mathcal{B}_{p,r}^s} \lesssim \sigma^n(\varepsilon) \rightarrow 0,$$

as $\varepsilon \rightarrow 0$. The proof that z_ε also converges to 0 in $\mathcal{C}([0, T]; \mathcal{B}_{p,r}^s)$ is virtually identical to the proof of Proposition 3.5, so we omit it. \square

It remains to establish that $\lim_{\varepsilon \rightarrow 0} \|v_\varepsilon - w_\lambda\|_{\mathcal{C}([0,T]; \mathcal{B}_{p,r}^s)} = 0$ in probability, which is the content of the following result:

Proposition 4.2. *Assume (3.23) and let $u_0 \in \mathcal{B}_{p,r}^s$. Let $v_\varepsilon \in \mathcal{E}_T$ be the unique mild solution to (4.2), and $w_\lambda \in \mathcal{E}_T$ the unique solution to (Ψ_λ) . Then*

$$\lim_{\varepsilon \rightarrow 0} \|v_\varepsilon - w_\lambda\|_{\mathcal{E}_T} = 0$$

in probability.

Proof. Setting $\delta_\varepsilon = 3D_\varepsilon^2 - \frac{3\lambda^2}{8\pi}$ and $a_\lambda = 1 - \frac{3\lambda^2}{8\pi}$, we can rewrite the equations for v_ε and w_λ as

$$\begin{aligned} \partial_t v_\varepsilon &= \Delta v_\varepsilon + a_\lambda v_\varepsilon - v_\varepsilon^3 - \delta_\varepsilon v_\varepsilon + (2 - 3D_\varepsilon^2)z_\varepsilon + \sum_{l=0}^2 a_l v_\varepsilon^l :z_\varepsilon^{3-l}:, \\ \partial_t w_\lambda &= \Delta w_\lambda + a_\lambda w_\lambda - w_\lambda^3. \end{aligned}$$

Setting $\rho_\varepsilon = v_\varepsilon - w_\lambda$, we see that ρ_ε solves the following evolution equation:

$$\begin{aligned} \partial_t \rho_\varepsilon &= (\Delta + a_\lambda)\rho_\varepsilon - \rho_\varepsilon(v_\varepsilon^2 + v_\varepsilon w_\lambda + w_\lambda^2) \\ &\quad + (2 - 3D_\varepsilon^2)z_\varepsilon - \delta_\varepsilon v_\varepsilon + \sum_{l=0}^2 a_l v_\varepsilon^l :z_\varepsilon^{3-l}:. \end{aligned}$$

Setting $\hat{A} = \Delta + a_\lambda$, we have the mild formulation

$$\begin{aligned} \rho_\varepsilon(t) &= e^{\hat{A}t} \rho_\varepsilon(0) - \int_0^t e^{\hat{A}(t-s)} \rho_\varepsilon(s) (v_\varepsilon^2 + v_\varepsilon w_\lambda + w_\lambda^2)(s) ds \\ &\quad + (2 - 3D_\varepsilon^2) \int_0^t e^{\hat{A}(t-s)} z_\varepsilon(s) ds - \delta_\varepsilon \int_0^t e^{\hat{A}(t-s)} v_\varepsilon(s) ds \\ &\quad + \sum_{l=0}^2 a_l \int_0^t e^{\hat{A}(t-s)} v_\varepsilon^l(s) : z_\varepsilon^{3-l}(s) ds . \end{aligned}$$

It then follows from Lemmas A.1 and A.2 that

$$\begin{aligned} \|\rho_\varepsilon\|_{\mathcal{E}_T} &\lesssim \|\rho_\varepsilon(0)\|_{\mathcal{B}_{p,r}^s} + T^\delta \|\rho_\varepsilon\|_{\mathcal{E}_T} (\|v_\varepsilon\|_{L^p([0,T];\mathcal{B}_{p,r}^s)}^2 + \|w_\lambda\|_{L^p([0,T];\mathcal{B}_{p,r}^s)}^2) \\ &\quad + \delta_\varepsilon \|v_\varepsilon\|_{\mathcal{E}_T} + T^\delta \sum_{l=0}^2 (1 + \|v_\varepsilon\|_{\mathcal{E}_T}^l) \|z_\varepsilon^{3-l}\|_{L^p([0,T];\mathcal{B}_{p,r}^s)} . \end{aligned} \tag{4.3}$$

We now use the fact that there exists K such that the deterministic solution w_λ satisfies $\|w_\lambda\|_{\mathcal{E}_T} \leq K$. Setting $\tau_\varepsilon = \bar{T} \wedge \inf\{t : \|\rho_\varepsilon\|_{\mathcal{E}_t} \geq 1\}$ for some $\bar{T} \leq T$ such that $\bar{T}^\delta ((K+1)^2 + K^2) \leq \frac{1}{2}$, it follows from (4.3) that

$$\|\rho_\varepsilon\|_{\mathcal{E}_{\tau_\varepsilon,\delta}} \lesssim \|\rho_\varepsilon(0)\|_{\mathcal{B}_{p,r}^s} + \bar{T}^\delta \sum_{l=0}^2 (1 + K)^l (\delta_\varepsilon + \|z_\varepsilon^{3-l}\|_{L^p([0,T];\mathcal{B}_{p,r}^s)}) .$$

This bound can easily be iterated, and the claim then follows similarly to the proof of Proposition 3.8. \square

A Technical results

In this appendix, we collect a few technical results.

Lemma A.1. *Let $A = \Delta - \Lambda$ for $\Lambda \geq 1$ and let $f \in L^{p/n}([0, T]; \mathcal{B}_{p,r}^{(2n-1)s})$ with $p > n \geq 1$, $s < 0$ and $\bar{s} = 2/p + 2s$ such that*

$$(n-1)s + 1 - \frac{n}{p} > 0. \tag{A.1}$$

Then there exists $\delta > 0$ such that

$$\left\| \int_0^t e^{(t-\tau)A} f(\tau) d\tau \right\|_{\mathcal{E}_T} \leq K(\Lambda) T^\delta \|f\|_{L^{p/n}([0,T];\mathcal{B}_{p,r}^{(2n-1)s})} ,$$

with a constant $K(\Lambda)$ such that $\lim_{\Lambda \rightarrow \infty} K(\Lambda) = 0$.

Proof. Modulo straightforward modifications yielding $K(\Lambda) \rightarrow 0$, the proof is identical to the proof of [6, Lem. 3.6]. \square

Lemma A.2. *Let $n, p, r \geq 1$, $s < 0$, $\bar{s} = 2/p + 2s$ such that $|s| < \frac{2}{p(2n+1)}$ and $l < n$. Assume that $g_i \in L^p([0, T]; \mathcal{B}_{p,r}^{\bar{s}})$ for $i = 1, \dots, l$ and $h \in L^p([0, T]; \mathcal{B}_{p,r}^s)$. Then, there exists a constant $C > 0$ such that*

$$\|h g_1 \cdots g_l\|_{L^{p/(l+1)}([0,T];\mathcal{B}_{p,r}^{(2l+1)s})} \leq C \|h\|_{L^p([0,T];\mathcal{B}_{p,r}^s)} \prod_{j=1}^l \|g_j\|_{L^p([0,T];\mathcal{B}_{p,r}^{\bar{s}})} . \tag{A.2}$$

Proof. This is a straightforward modification of [6, Cor. 3.5]. \square

Lemma A.3. *Let $p, r \geq 1$ and $\bar{s} < s$. Then, there exists a constant $C > 0$ such that*

$$\|e^{t\Delta}x\|_{\mathcal{B}_{p,r}^{\bar{s}}} \leq Ct^{\frac{\bar{s}-s}{2}} \|x\|_{\mathcal{B}_{p,r}^s} \quad \forall x \in \mathcal{B}_{p,r}^{\bar{s}}.$$

Proof. The estimate follows from [4, Lem. 2.4] and the definition of the Besov norm (2.1). \square

Corollary A.4. *Let $s < 0$, $r, p \geq 1$, and $\bar{s} = 2s + \frac{2}{p}$. Define the operator $A = \Delta - \Lambda$ and recall the \mathcal{E}_T -norm as defined in (3.22). Then there exists $\delta > 0$ such that for all $\Lambda > 1$,*

$$\|e^{tA}x\|_{\mathcal{E}_T} \leq (1 + C(\Lambda)T^\delta) \|x\|_{\mathcal{B}_{p,r}^s}, \quad \forall x \in \mathcal{B}_{p,r}^s,$$

where $\lim_{\Lambda \rightarrow \infty} C(\Lambda) = 0$.

Proof. The bound on the $C([0, T]; \mathcal{B}_{p,r}^s)$ norm is trivial. Using Proposition A.3, we obtain for arbitrary $\gamma > 0$

$$\begin{aligned} \|e^{tA}x\|_{L^p([0,T]; \mathcal{B}_{p,r}^s)} &\leq \frac{K}{\Lambda^{\gamma/p}} \left(\int_0^T \frac{1}{t^\gamma} \|e^{t\Delta}x\|_{\mathcal{B}_{p,r}^s} dt \right)^{1/p} \\ &\leq \frac{K}{\Lambda^{\gamma/p}} \|x\|_{\mathcal{B}_{p,r}^s} \left(\int_0^T t^{p(s-\bar{s})/2-\gamma} dt \right)^{1/p} \\ &\leq \frac{K}{\Lambda^{\gamma/p}} \|x\|_{\mathcal{B}_{p,r}^s} T^{|s|/2-\gamma/p}. \end{aligned}$$

Choosing $\gamma < \frac{p}{2}|s|$, the claim follows. \square

References

- [1] S. Allen and J. Cahn. A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening. *Acta Metall.* **27**, no. 6, (1979), 1085–1095.
- [2] S. Albeverio and M. Röckner. Stochastic differential equations in infinite dimensions: solutions via Dirichlet forms. *Probab. Theory Related Fields* **89**, no. 3, (1991), 347–386. MR-1113223
- [3] F. Barret, A. Bovier, and S. Méléard. Uniform estimates for metastable transition times in a coupled bistable system. *Electron. J. Probab.* **15**, no. 12, (2010), 323–345. MR-2609590
- [4] H. Bahouri, J. Chemin, and R. Danchin. *Fourier analysis and nonlinear partial differential equations*, vol. 343 of *Grundlehren der mathematischen Wissenschaften Series*. Springer Verlag, 2010.
- [5] S. Cerrai, and M. Freidlin. Approximation of quasi-potentials and exit problems for multidimensional RDE’s with noise. *Trans. Amer. Math. Soc.* **363**, no. 7, (2011), 3853–3892. MR-2775830
- [6] G. Da Prato and A. Debussche. Strong solutions to the stochastic quantization equations. *Ann. Probab.* **31**, no. 4, (2003), 1900–1916. MR-2016604
- [7] G. Da Prato and L. Tubaro. Wick powers in stochastic PDEs: an introduction. *Technical Report UTM 711, University of Trento* (2007).
- [8] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, vol. 45 of *Encyclopedia of mathematics and its applications*. Cambridge University Press, 1992. MR-1207136
- [9] L.C. EVANS, H.M. Soner, and P.E. Souganidis. Phase transitions and generalized motion by mean curvature. *Commun. Pure Appl. Math.* **45**, no. 9, (1992), 1097–1123. MR-1177477
- [10] J. Glimm and A. Jaffe. *Quantum Physics: A Functional Integral Point of View*. Second ed., Springer-Verlag, New York, 1987. MR-0887102
- [11] M. Hairer. An introduction to stochastic PDEs. <http://www.hairer.org/Teaching.html>, 2009. Unpublished lecture notes.

- [12] T. Ilmanen. Convergence of the Allen-Cahn equation to Brakke’s motion by mean curvature. *J. Differential. Geom.* **38**, no. 2, (1993), 417–461. MR-1237490
- [13] R. Kohn, F. Otto, M. Reznikoff, and E. Vanden-Eijnden. Action minimization and sharp-interface limits for the stochastic Allen-Cahn equation. *Commun. Pure Appl. Math.* **60**, no. 3, (2007), 393–438. MR-2284215
- [14] L. Landau, and L. Ginzburg. On the theory of superconductivity. *J. Expt. Theor. Phys.* **20**, (1950), 1064–1082.
- [15] G. Parisi, and Y.S. Wu. Perturbation theory without gauge fixing. *Sci. Sinica* **24**, no. 4, (1981), 483–496. MR-0626795
- [16] M.D. Ryser, N. Nigam, and P.F. Tupper. On the well-posedness of the stochastic Allen-Cahn equation in two dimensions. *J. Comp. Phys.* **231**, no. 6, (2012), 2537–2550.
- [17] J. Walsh. An introduction to stochastic partial differential equations. *École d’Été de Probabilités de Saint Flour XIV-1984* 265–439. MR-0876085

Acknowledgments. MH acknowledges financial support by the EPSRC through grant EP/D071593/1 and the Royal Society through a Wolfson Research Merit Award. Both MH and HW were supported by the Leverhulme Trust through a Philip Leverhulme Prize. MDR is grateful to P.F. Tupper and N. Nigam for fruitful discussions, and acknowledges financial support from a Hydro-Québec Doctoral Fellowship. We would like to thank F. Otto for suggesting to also consider the case $\sigma(\varepsilon) \rightarrow 0$. All three authors are grateful for the relaxed atmosphere at the Newton institute, where this collaboration was initiated during the 2010 “Stochastic PDEs” programme.