

## THE HYPER ORDER AND FIXED POINTS OF SOLUTIONS OF LINEAR DIFFERENTIAL EQUATIONS

NAN LI AND LIANZHONG YANG\*

ABSTRACT. In this paper, we obtain a precise estimation of the hyper order of solutions for a class of higher order linear differential equation, and also investigate the exponents of convergence of the fixed points of solutions and their first derivatives for the second order case. These results generalize the results of Gundersen-Steinbart, Wittich and Chen-Shon.

**AMS Subject Classification:** 30D35.

**Keywords:** Differential equation, Hyper order, Fixed points.

### 1. INTRODUCTION

In this paper, we will use standard notations from the value distribution theory of meromorphic functions (see [15] [20]). We suppose that  $f(z)$  is a meromorphic function in whole complex plane  $\mathbf{C}$ . In addition, we denote the order of growth of  $f(z)$  by  $\sigma(f)$ , and also use the notation  $\sigma_2(f)$  to denote the hyper-order of  $f(z)$ , defined by

$$\sigma_2(f) = \limsup_{r \rightarrow \infty} \frac{\log \log T(r, f)}{\log r}.$$

To give the precise estimate of fixed points, we define the exponent of convergence of fixed points by  $\tau(f)$

$$\tau(f) = \limsup_{r \rightarrow \infty} \frac{\log N(r, \frac{1}{f-z})}{\log r},$$

and also the hyper-exponent of convergence of (distinct) fixed points by  $\tau_2(f)$ ( $\bar{\tau}_2(f)$ )

$$\tau_2(f) = \limsup_{r \rightarrow \infty} \frac{\log \log N(r, \frac{1}{f-z})}{\log r},$$

$$\bar{\tau}_2(f) = \limsup_{r \rightarrow \infty} \frac{\log \log \bar{N}(r, \frac{1}{f-z})}{\log r}.$$

Recently, many scholars devoted to investigating the growth of solutions of complex differential equations, see [1-3, 5-9, 11, 12, 16-19, 21].

---

\*Corresponding author.

Consider the second order homogeneous linear periodic differential equation

$$f'' + P(e^z)f' + Q(e^z)f = 0, \quad (1.1)$$

where  $P(z)$  and  $Q(z)$  are polynomials in  $z$  and not both constants. It is well known that every solution of  $f$  is an entire.

Suppose  $f \not\equiv 0$  is a solution of (1.1) and if  $f$  satisfies the condition

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{r} = 0, \quad (1.2)$$

then, we say that  $f$  is a nontrivial subnormal solution of (1.1).

Wittich [17] investigated the subnormal solution of (1.1), and obtained the form of all subnormal solutions in the following theorem.

**Theorem A.** If  $f(\not\equiv 0)$  is a subnormal solution of (1.1), then  $f$  must have the form

$$f(z) = e^{cz}(h_0 + h_1e^z + \cdots + h_me^{mz}) \quad (1.3)$$

where  $m \geq 0$  is an integer and  $c, h_0, \dots, h_m$  are constants with  $h_0 \neq 0$  and  $h_m \neq 0$ .

Gundersen and Steinbart [12] refined Theorem A and got the following theorem.

**Theorem B.** Under the assumption of Theorem A, the following statements hold.

(i) if  $\deg P > \deg Q$  and  $Q \not\equiv 0$ , then, any subnormal solution  $f \not\equiv 0$  of (1.1) must have the form

$$f(z) = \sum_{k=0}^m h_k e^{-kz},$$

where  $m \geq 1$  is an integer and  $h_0, h_1, \dots, h_m$  are constants with  $h_0 \neq 0$  and  $h_m \neq 0$ .

(ii) if  $\deg P \geq 1$  and  $Q = 0$ , then any subnormal solution of equation (1.1) must be a constant,

(iii) if  $\deg P < \deg Q$ , then the subnormal solution of equation (1.1) is  $f = 0$ .

Chen and Shon [6] investigate more general equation than (1.1), and get the following theorem.

Set

$$a_j(z) = a_{jd_j}z^{d_j} + a_{j(d_j-1)}z^{d_j-1} + \cdots + a_{j1}z + a_{j0} \quad (1.4)$$

$$b_k(z) = b_{kd_k}z^{d_k} + b_{k(d_k-1)}z^{d_k-1} + \cdots + b_{k1}z + b_{k0} \quad (1.5)$$

where  $d_j \geq 0$ ,  $m_k \geq 0$  ( $j = 1, \dots, n$ ,  $k = 1, \dots, s$ ) are integers.  $a_{jd_j}, \dots, a_{j0}$ ;  $b_{kd_k}, \dots, b_{k0}$  are constants.  $a_{jd_j} \neq 0$ ,  $b_{kd_k} \neq 0$ .

**Theorem C.** Let  $a_n(z), \dots, a_1(z)$ ,  $b_s(z), \dots, b_1(z)$  be polynomials and satisfy (1.4) and (1.5), and  $a_n(z)b_s(z) \neq 0$ . Suppose that

$$P(e^z) = a_n(z)e^{nz} + \cdots + a_1(z)e^z, \quad Q(e^z) = b_s(z)e^{sz} + \cdots + b_1(z)e^z.$$

If  $n \neq s$ , then every solution  $f (\neq 0)$  of equation

$$f'' + P(e^z)f' + Q(e^z)f = 0 \quad (1.6)$$

satisfies  $\sigma_2(f) = 1$ .

For the higher-order linear homogeneous differential equation

$$f^{(k)} + P_{k-1}(e^z)f^{(k-1)} + \cdots + P_0(e^z)f = 0, \quad (1.7)$$

where  $P_j(e^z)$  ( $j = 0, \dots, k-1$ ) are polynomials in  $z$ , many papers were devoted to investigate the solutions of (1.7) (see [3] [5] [7] [8] [16]).

In [7] Chen and Shon consider the existence of subnormal solution of (1.7) and obtain the following theorem.

**Theorem D.** Let  $P_j(z)$  ( $j = 0, \dots, k-1$ ) be polynomials in  $z$  such that all constant terms of  $P_j$  are equal to zero and  $\deg P_j = m_j$ , that is,

$$P_j(e^z) = a_{jm_j}e^{m_jz} + a_{j(m_j-1)}e^{(m_j-1)z} + \cdots + a_{j1}e^z,$$

where  $a_{jm_j}, a_{j(m_j-1)}, \dots, a_{j1}$  are constants and  $a_{jm_j} \neq 0$ ;  $m_j \geq 1$  are integers. Suppose that there exists  $m_s$  ( $s \in \{0, \dots, k-1\}$ ) satisfying

$$m_s > \max\{m_j : j = 0, \dots, s-1, s+1, \dots, k-1\} = m.$$

Then one has the following properties.

(i) If  $P_0 \neq 0$ , then (1.7) has no nontrivial subnormal solution and every solution of (1.7) is of hyper order  $\sigma_2(f) = 1$ .

(ii) If  $P_0 \equiv \cdots \equiv P_{d-1} \equiv 0$  and  $P_d \neq 0$  ( $d < s$ ), then any polynomials with degree  $\leq d-1$  are subnormal solutions of (1.7) and all other solutions  $f$  of (1.7) satisfy  $\sigma_2(f) = 1$ .

It is natural to ask the following question: whether the result of Theorem B can be generalized to the higher order case under the condition

of Theorem C. In this paper, we first investigate the problem and obtain the following result.

Set

$$a_{jm_i}(z) = a_{jm_id_{jm_i}}z^{d_{jm_i}} + a_{jm_i(d_{jm_i}-1)}z^{d_{jm_i}-1} + \cdots + a_{jm_i1}z + a_{jm_i0} \quad (1.8)$$

where  $d_{jm_i} \geq 0$  ( $j = 1, \dots, n$ ) are integers,  $a_{jm_id_{jm_i}}, \dots, a_{jm_i0}$  are constants,  $a_{jm_id_{jm_i}} \neq 0$ .

**Theorem 1.** Let  $a_{jm_i}(z)$  be polynomials and satisfy (1.8). Suppose that

$$P_j(e^z) = a_{jm_j}(z)e^{m_j z} + \cdots + a_{j1}(z)e^z \quad (1.9)$$

where  $a_{jm_i}(z) \not\equiv 0$ . If there exists an integer  $s$  ( $s \in \{0, \dots, k-1\}$ ) satisfying

$$m_s > \max\{m_j : j = 0, \dots, s-1, s+1, \dots, k-1\} = m, \quad (1.10)$$

then every nonconstant solution  $f$  of equation

$$f^{(k)} + P_{k-1}(e^z)f^{(k-1)} + \cdots + P_0(e^z)f = 0 \quad (1.11)$$

satisfies  $\sigma_2(f) = 1$  if one of the following condition holds.

- (1)  $s = 0$  or  $1$ .
- (2)  $s \geq 2$  and  $\deg a_{0j}(z) > \deg a_{ij}(z)$  ( $i \neq 0$ ).

For almost four decades, a lot of results have been obtained on the fixed points of general transcendental meromorphic function. However, there are few studies on fixed points of differential polynomials generated by solutions of differential equation. In 2000, Z.X.Chen [4] first pointed out the relation between the exponent of convergence of distinct fixed points and the rate of growth of solutions of second-order linear differential equations with entire coefficients. In this paper, we continue to investigate the relation between the hyper-exponent of convergence of distinct fixed points and the rate of growth of solutions for a higher order case.

**Theorem 2.** Under the assumption of Theorem 1, if  $zP_0(e^z) + P_1(e^z) \not\equiv 0$ , then we have every nonconstant solution  $f$  of equation (1.11) satisfies

$$\tau_2(f) = \bar{\tau}_2(f) = \sigma_2(f) = 1.$$

In particular, we investigate the exponents of convergence of the fixed points of solutions and their first derivatives for a second order equation (1.6). we will prove the following theorems:

**Theorem 3.** Let  $a_n(z), \dots, a_1(z), b_s(z), \dots, b_1(z)$  be polynomials and satisfy (1.4) and (1.5), and  $a_n(z)b_s(z) \neq 0$ . Suppose that

$$P(e^z) = a_n(z)e^{nz} + \dots + a_1(z)e^z, \quad Q(e^z) = b_s(z)e^{sz} + \dots + b_1(z)e^z.$$

If  $s \neq n$ , then every solution  $f (\neq 0)$  of equation (1.6) satisfy  $\lambda(f - z) = \lambda(f' - z) = \sigma(f) = \infty$  and  $\lambda_2(f - z) = \lambda_2(f' - z) = \sigma_2(f)$ .

## 2. SOME LEMMAS

**Lemma 1.** ([20]) Let  $f_j(z) (j = 1, \dots, n) (n \geq 2)$  be meromorphic functions,  $g_j(z) (j = 1, \dots, n)$  be entire functions, and satisfy

- (1)  $\sum_{j=1}^n e^{g_j(z)} \equiv 0$ ;
- (2) when  $1 \leq j < k \leq n$ , then  $g_i(z) - g_k(z)$  is not a constant;
- (3) when  $1 \leq j \leq n, 1 \leq h < k \leq n$ , then

$$T(r, f_j) = o\{T(r, e^{g_h - g_k})\} \quad (r \rightarrow \infty, r \notin E),$$

where  $E \subset (1, \infty)$  is of finite linear measure or logarithmic measure. Then,  $f_j(z) \equiv 0 (j = 1, \dots, n)$ .

**Lemma 2** Let  $P_j(e^z), m_j, m_s, m$  and  $a_{ij}(z)$  satisfy the hypotheses of Theorem 1. Then equation (1.11) has no nonconstant polynomial solution.

Proof. Suppose that  $f_0 = b_n z^n + \dots + b_1 z + b_0 (n \geq 1, b_n, \dots, b_0$  are constants,  $b_n \neq 0)$  is a nonconstant solution of (1.11).

If  $n \geq s$ , then  $f_0^{(s)} \neq 0$ . Substituting  $f_0$  into (1.11) and taking  $z = r$ , we conclude that

$$\begin{aligned} & |a_{sm_s d_{sm_s}}| r^{d_{sm_s}} e^{m_s r} |b_n n(n-1) \dots (n-s+1)| r^{n-s} (1 + o(1)) \\ & \leq | -P_s(e^z) f_0^{(s)}(z) | \\ & \leq |f_0^{(k)}(z)| + |P_{k-1}(e^z) f_0^{(k-1)}(z)| + \dots + |P_{s+1}(e^z) f_0^{(s+1)}(z)| \\ & \quad + |P_{s-1}(e^z) f_0^{(s-1)}(z)| + \dots + |P_0(e^z) f_0(z)| \\ & \leq M r^d e^{mr} (1 + o(1)). \end{aligned} \tag{2.1}$$

Since  $m_s > m$  we see that (2.1) is a contradiction.

Obviously, when  $s = 0$  or  $1$ , we can get that the equation (1.11) has no nonconstant polynomial solution from the above process.

If  $n < s$ , then

$$P_n(e^z) f_0^{(n)}(z) + \dots + P_0(e^z) f_0(z) = 0. \tag{2.2}$$

Set  $\max\{m_i : i = 0, \dots, n\} = h$ . If  $m_j < h$ , then we can rewrite

$$\begin{aligned} P_j(e^z) &= a_{jh}(z) e^{hz} + \dots + a_{j(m_j+1)}(z) e^{(m_j+1)z} \\ & \quad + a_{jm_j}(z) e^{m_j z} + \dots + a_{j1}(z) e^z \quad (j = 0, \dots, n), \end{aligned} \tag{2.3}$$

where  $a_{jh}(z) = \cdots = a_{j(m_j+1)}(z) = 0$ .

Thus we conclude by (2.2) and (2.3) that

$$\begin{aligned} & (a_{nh}(z)f_0^{(n)} + a_{(n-1)h}(z)f_0^{(n-1)} + \cdots + a_{0h}f_0)e^{hz} + \cdots \\ & \quad + (a_{nj}(z)f_0^{(n)} + a_{(n-1)j}(z)f_0^{(n-1)} + \cdots + a_{0j}f_0)e^{jz} + \cdots \\ & \quad + (a_{n1}(z)f_0^{(n)} + a_{(n-1)1}(z)f_0^{(n-1)} + \cdots + a_{01}f_0)e^z = 0. \end{aligned} \quad (2.4)$$

Set

$$Q_j(z) = a_{nj}(z)f_0^{(n)} + a_{(n-1)j}(z)f_0^{(n-1)} + \cdots + a_{0j}f_0 \quad (j = 1, \dots, h). \quad (2.5)$$

Since  $f_0$  and  $a_{ij}(z)$  are polynomials, we see that

$$m(r, Q_j) = o\{m(r, e^{(\alpha-\beta)z})\} \quad (1 \leq \beta < \alpha \leq h). \quad (2.6)$$

By Lemma 1 and (2.4)-(2.6), we conclude that

$$Q_1(z) \equiv Q_2(z) \equiv \cdots \equiv Q_h(z) \equiv 0. \quad (2.7)$$

Since  $\deg f_0 > \deg f_0' > \cdots > \deg f_0^{(n)}$  and  $\deg a_{0j}(z) > \deg a_{ij}(z)$  ( $i \neq 0$ ), by (2.5) and (2.7), we get a contradiction.

**Lemma 3.** [11] Let  $f(z)$  be an entire function and suppose that  $|f^{(k)}(z)|$  is unbounded on some ray  $\arg z = \theta$ . Then, there exists an infinite sequence of points  $z_n = r_n e^{i\theta}$  ( $n = 1, 2, \dots$ ), where  $r_n \rightarrow \infty$ , such that  $f^{(k)}(z_n) \rightarrow \infty$  and

$$\frac{|f^{(j)}(z)|}{|f^{(k)}(z)|} \leq |z_n|^{(k-j)}(1 + o(1)) \quad (j = 0, \dots, k-1). \quad (2.8)$$

**Lemma 4.** [10] Let  $f(z)$  be a transcendental meromorphic function with  $\sigma(f) = \sigma < \infty$ , Let  $\Gamma = \{(k_1, j_1), \dots, (k_m, j_m)\}$  be a finite set of distinct pairs of integers satisfying  $k_i > j_i \geq 0$  for  $i = 1, 2, \dots, m$ . Also let  $\varepsilon > 0$  be a given constant, then there exists a set  $E \subset [0, 2\pi)$  that has linear measure zero, such that if  $\psi \in [0, 2\pi) \setminus E$ , then there is constant  $R_0 = R_0(\psi) > 1$  such that for all  $z$  satisfying  $\arg z = \psi$  and  $|z| \geq R_0$  and for all  $(k, j) \in \Gamma$ , we have

$$\frac{|f^{(k)}(z)|}{|f^{(j)}(z)|} \leq |z|^{(k-j)(\sigma-1+\varepsilon)}. \quad (2.9)$$

**Remark 1** Obviously, in Lemma 4, if  $\psi \in [0, 2\pi) \setminus E$  is replaced by  $\psi \in [-\frac{\pi}{2}, \frac{3\pi}{2}) \setminus E$ , then (2.9) still holds.

**Lemma 5.** [5] Let  $f(z)$  be an entire function with  $\sigma(f) = \sigma < \infty$ . Suppose that there exists a set  $E \subset [0, 2\pi)$  that has linear measure

zero, such that for any ray  $\arg z = \theta_0 \in [0, 2\pi) \setminus E$ ,  $|f(re^{i\theta_0})| \leq Mr^k$  ( $M = M(\theta_0) > 0$  is a constant and  $k > 0$  is a constant independent of  $\theta_0$ ). Then  $f(z)$  is a polynomial with  $\deg f \leq k$ .

**Lemma 6.** [8] Let  $A_0, \dots, A_{k-1}$  be entire functions of finite order. If  $f(z)$  is a solution of equation

$$f^{(k)} + A_{k-1}f^{(k-1)} + \dots + A_0f = 0,$$

then  $\sigma_2(f) \leq \max\{\sigma(A_j) : j = 0, \dots, k-1\}$ .

**Lemma 7.** [9] Let  $g(z)$  be an entire function of infinite order with the hyper-order  $\sigma_2 = \sigma$ , and let  $\nu(r)$  be the central index of  $g$ . Then,

$$\limsup_{r \rightarrow \infty} \frac{\log \log \nu(r)}{\log r} = \sigma_2(g) = \sigma.$$

**Lemma 8.** [6] Let  $f(z)$  be an entire function of infinite order with  $\sigma_2 = \alpha$  ( $0 \leq \alpha < \infty$ ), and a set  $E \subset [1, \infty)$  have a finite logarithmic measure. Then, there exists  $\{z_k = r_k e^{i\theta_k}\}$ , such that  $|f(z_k)| = M(r_k, f)$ ,  $\theta_k \in [-\frac{\pi}{2}, \frac{3\pi}{2})$ ,  $\lim_{k \rightarrow \infty} \theta_k = \theta_0 \in [-\frac{\pi}{2}, \frac{3\pi}{2})$ ,  $r_k \notin E$ , and  $r_k \rightarrow \infty$ , such that (1) if  $\sigma_2(f) = \alpha$  ( $0 < \alpha < \infty$ ), then for any given  $\varepsilon_1$  ( $0 < \varepsilon_1 < \alpha$ ),

$$\exp\{r_k^{\alpha-\varepsilon_1}\} < \nu(r_k) < \exp\{r_k^{\alpha+\varepsilon_1}\}. \quad (2.10)$$

(2) if  $\sigma(f) = \infty$  and  $\sigma_2(f) = 0$ , then for any given  $\varepsilon_2$  ( $0 < \varepsilon_2 < \frac{1}{2}$ ), and any large  $M$  ( $> 0$ ), we have, as  $r_k$  sufficiently large,

$$r_k^M < \nu(r_k) < \exp\{r_k^{\varepsilon_2}\}. \quad (2.11)$$

**Lemma 9.** [10] Let  $f$  be a transcendental meromorphic function, and  $\alpha > 1$  be a given constant. Then there exists a set  $E \subset (1, \infty)$  with finite logarithmic measure and a constant  $B > 0$  that depends only on  $\alpha$  and  $i, j$  ( $i < j, j \in \mathbb{N}$ ), such that for all  $z$  satisfying  $|z| = r \notin E \cup [0, 1]$ ,

$$\left| \frac{f^{(j)}(z)}{f^{(i)}(z)} \right| \leq B \left( \frac{T(\alpha r, f)}{r} (\log^\alpha r) \log T(\alpha r, f) \right)^{j-i}. \quad (2.12)$$

**Remark 2** From the proof of Lemma 9, we can see that the exceptional set  $E$  satisfies that if  $a_n$  and  $b_m$  ( $n, m = 1, 2, \dots$ ) denote all zeros and poles of  $f$ , respectively,  $O(a_n)$  and  $O(b_m)$  denote sufficiently small neighborhoods of  $a_n$  and  $b_m$ , respectively, then

$$E = \{|z| : z \in (\cup_{n=1}^{+\infty} O(a_n)) \cup (\cup_{m=1}^{+\infty} O(b_m))\}.$$

Hence, if  $f(z)$  is a transcendental entire function, and  $z$  is a point that satisfies  $|f(z)|$  to be sufficiently large, then (2.12) holds. For details see [7] Remark 2.10.

**Lemma 10.**(See [14] and Satz 21.2 of [13]) Let  $g$  be a non-constant entire function, and let  $0 < \delta < 1$ . There exists a set  $E \subset [1, \infty)$  of finite logarithmic measure with the following property. For  $r \in [1, \infty) \setminus E$ , the central index  $\nu(r)$  of  $g$  satisfies

$$\nu(r) \leq (\log M(r, g))^{1+\delta}.$$

**Lemma 11.** [3] Let  $A_0, \dots, A_{k-1}, F \not\equiv 0$  be finite order meromorphic functions. If  $f$  is a meromorphic solution the equation

$$f^{(k)} + A_{k-1}f^{(k-1)} + \dots + A_0f = F, \quad (2.13)$$

with  $\rho(f) = +\infty$  and  $\rho_2(f) = \rho$ , then  $f$  satisfies  $\bar{\lambda}(f) = \lambda(f) = \rho(f) = +\infty$  and  $\bar{\lambda}_2(f) = \lambda_2(f) = \rho_2(f) = \rho$ .

### 3. PROOF OF THEOREM 1

Suppose that  $f \not\equiv 0$  is a solution of (1.11), then,  $f$  is an entire function. By Lemma 2, we see that  $f$  is transcendental.

**First step.**we prove that  $\sigma(f) = \infty$ .

Assume that  $f$  is transcendental with  $\sigma(f) < \infty$ . By Lemma 4, we know that for any given  $\varepsilon > 0$ , there exists a set  $E \subset [-\frac{\pi}{2}, \frac{3\pi}{2})$  having linear measure zero, such that if  $\psi \in [-\frac{\pi}{2}, \frac{3\pi}{2}) \setminus E$ , then there is a constant  $R_0 = R_0(\psi) > 1$  such that for all  $z$  satisfying  $\arg z = \psi$  and  $|z| = r > R_0$ , we have

$$\left| \frac{f^{(j)}(z)}{f^{(s)}(z)} \right| \leq r^{(\sigma-1+\varepsilon)(j-s)} \quad j = s+1, \dots, k. \quad (3.1)$$

**Case 1** Now we take a ray  $\arg z = \theta \in (-\frac{\pi}{2}, \frac{\pi}{2}) \setminus E$ . Then we have  $\cos \theta > 0$ . We assert that  $|f^{(s)}(re^{i\theta})|$  is bounded on the ray  $\arg z = \theta$ . If  $|f^{(s)}(re^{i\theta})|$  is unbounded on the ray  $\arg z = \theta$ , then by Lemma 3, there exists a sequence  $\{z_t = r_t e^{i\theta}\}$  such that as  $r_t \rightarrow \infty$ ,  $f^{(s)}(z_t) \rightarrow \infty$  and

$$\left| \frac{f^{(i)}(z_t)}{f^{(s)}(z_t)} \right| \leq r_t^{s-i}(1 + o(1)) \quad i = 0, \dots, s-1. \quad (3.2)$$

By (1.11), we get that

$$-P_s(e^{z_t}) = \frac{f^{(k)}(z_t)}{f^{(s)}(z_t)} + \sum_{j=0, j \neq s}^{k-1} [P_j(e^{z_t})] \frac{f^{(j)}(z_t)}{f^{(s)}(z_t)}. \quad (3.3)$$



$$\begin{aligned}
|P_s(e^{z_t})| &= |a_{sm_s}(z_t)e^{m_s z_t} + \cdots + a_1(z_t)e^{z_t}| \\
&\geq |a_{sm_s}(z_t)e^{m_s z_t}| - [|a_{s(m_s-1)}(z_t)e^{(m_s-1)z_t}| + \cdots + |a_{s1}(z_t)e^{z_t}|] \\
&= |a_{sm_s d_{sm_s}}| r^{d_{sm_s}} (1 + o(1)) e^{m_s r_t \cos \theta} \\
&\quad - [|a_{s(m_s-1) d_{s(m_s-1)}}| r^{d_{s(m_s-1)}} e^{(m_s-1)r_t \cos \theta} (1 + o(1)) + \cdots \\
&\quad + |a_{s1 d_{s1}}| r^{d_{s1}} e^{r_t \cos \theta} (1 + o(1))] \\
&\geq \frac{1}{2} |a_{sm_s d_{sm_s}}| r^{d_{sm_s}} e^{m_s r_t \cos \theta} (1 + o(1)) \tag{3.4}
\end{aligned}$$

and

$$|P_j(e^{z_t})| \leq 2 |a_{jm_j d_{jm_j}}| r_t^{d_{jm_j}} e^{m_j r_t \cos \theta} (1 + o(1)) \quad j \neq s. \tag{3.5}$$

By substituting (3.1) (3.2) (3.4) and (3.5) into (3.3), we obtain that

$$\frac{1}{2} |a_{sm_s d_{sm_s}}| r^{d_{sm_s}} e^{m_s r_t \cos \theta} (1 + o(1)) \leq 2 |a_{jm_j d_{jm_j}}| r_t^{d_{jm_j} + k\sigma} e^{m_j r_t \cos \theta} (1 + o(1)) \tag{3.6}$$

Since  $m_s > m$  and  $\cos \theta > 0$ , we know that when  $r_t \rightarrow \infty$ , (3.6) is a contradiction.

Hence when  $\arg z = \theta \in (-\frac{\pi}{2}, \frac{\pi}{2}) \setminus E$ , we have  $|f^{(s)}(re^{i\theta})| \leq M$ , so, on the ray  $\arg z = \theta \in (-\frac{\pi}{2}, \frac{\pi}{2}) \setminus E$ ,

$$|f(re^{i\theta})| \leq Mr^s. \tag{3.7}$$

**Case 2** Now we take a ray  $\arg z = \theta \in (\frac{\pi}{2}, \frac{3\pi}{2}) \setminus E$ . Then we have  $\cos \theta < 0$ . We assert that  $|f^{(k)}(re^{i\theta})|$  is bounded on the ray  $\arg z = \theta$ . If  $|f^{(k)}(re^{i\theta})|$  is unbounded on the ray  $\arg z = \theta$ , then by Lemma 3, there exists a sequence  $\{z_t = r_t e^{i\theta}\}$  such that as  $r_t \rightarrow \infty$ ,  $f^{(k)}(z_t) \rightarrow \infty$  and

$$\left| \frac{f^{(i)}(z_t)}{f^{(k)}(z_t)} \right| \leq r_t^{k-i} (1 + o(1)) \quad i = 0, \dots, k-1. \tag{3.8}$$

By (1.11), we get that

$$-1 = P_{k-1}(e^{z_t}) \frac{f^{(k-1)}(z_t)}{f^{(k)}(z_t)} + \cdots + P_0(e^{z_t}) \frac{f(z_t)}{f^{(k)}(z_t)}. \tag{3.9}$$

Since when  $r_t \rightarrow \infty$ ,

$$\begin{aligned}
|P_j(e^{z_t})| &= |a_{jm_j}(z_t)e^{m_j z_t} + a_{jm_{j-1}}(z_t)e^{(m_{j-1})z_t} + \cdots + a_{j1}(z_t)e^z| \\
&\leq |a_{jm_j d_{jm_j}}| r_t^{d_{jm_j}} e^{m_j r_t \cos \theta} (1 + o(1)) + \cdots \\
&\quad + |a_{j1 d_{j1}}| r^{d_{j1}} e^{r_t \cos \theta} (1 + o(1)). \tag{3.10}
\end{aligned}$$

By substituting (3.8) and (3.10) into (3.9), we obtain that

$$\begin{aligned}
 1 \leq & r_t [ |a_{k-1m_{k-1}d_{k-1m_{k-1}}}| r_t^{d_{k-1m_{k-1}}} e^{m_{k-1}r_t \cos \theta_t} (1 + o(1)) + \dots \\
 & + |a_{k-11d_{k-11}}| r_t^{d_{k-11}} e^{r_t \cos \theta_t} (1 + o(1)) ] + \dots \\
 & + r_t^k [ |a_{0m_0d_{0m_0}}| r_t^{d_{0m_0}} e^{m_0r_t \cos \theta_t} (1 + o(1)) \\
 & + \dots + |a_{01d_{01}}| r_t^{d_{01}} e^{r_t \cos \theta_t} (1 + o(1)) ]. \tag{3.11}
 \end{aligned}$$

Since  $\cos \theta_t < 0$ , when  $r_t \rightarrow \infty$ , by (3.11), we get  $1 \leq 0$ . This is a contradiction. Hence  $|f^{(k)}(re^{i\theta})| \leq M$  on the ray  $\arg z = \theta \in (\frac{\pi}{2}, \frac{3\pi}{2}) \setminus E$ . So, on the ray  $\arg z = \theta \in (\frac{\pi}{2}, \frac{3\pi}{2}) \setminus E$ , we have

$$|f(re^{i\theta})| \leq Mr^k. \tag{3.12}$$

Since the linear measure of  $E \cup \{-\frac{\pi}{2}, \frac{\pi}{2}\}$  is zero, by Lemma 5, (3.7) and (3.12), we know that  $f(z)$  is a polynomial. This contradicts our assumption that  $f(z)$  is transcendental. Therefore  $\sigma(f) = \infty$ .

**Second step.** We prove that  $\sigma_2(f) = 1$ .

By Lemma 6 and  $\sigma(P_i(e^z)) = 1$  ( $j = 0, \dots, k - 1$ ), we see that

$$\sigma_2(f) \leq \max\{\sigma(P_i(e^z))\} = 1. \tag{3.13}$$

Now we suppose that there exists a solution  $f_0$  satisfies  $\sigma_2(f_0) = \alpha < 1$ . Then we have

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, f_0)}{r} = 0. \tag{3.14}$$

By Lemma 9, we see that there exists a subset  $E_1 \subset (1, \infty)$  having finite logarithmic measure such that for all  $z$  satisfying  $|z| = r \notin E_1 \cup [0, 1]$ ,

$$\left| \frac{f_0^{(j)}(z)}{f_0(z)} \right| \leq M_0 [T(2r, f_0)]^{k+1}, \quad j = 1, \dots, k, \tag{3.15}$$

where  $M(> 0)$  is some constant.

From the Wiman-Valiron theory, there is a set  $E_2 \subset (1, \infty)$  having logarithmic measure  $lmE_2 < \infty$ , such that we can choose a  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_2$  and  $|f_0(z)| = M(r, f_0)$ , then we get

$$\frac{f_0^{(j)}(z)}{f_0(z)} = \left( \frac{\nu(r)}{z} \right)^j (1 + o(1)), \quad j = 0, \dots, k - 1. \tag{3.16}$$

where  $\nu(r)$  is the central index of  $f_0(z)$ .

By Lemma 8, we see that there exists a sequence  $\{z_t = r_t e^{i\theta_t}\}$  such that  $|f_0(z_t)| = M(r_t, f_0)$ ,  $\theta_t \in [-\frac{\pi}{2}, \frac{3\pi}{2})$ ,  $\lim \theta_t = \theta_0 \in [-\frac{\pi}{2}, \frac{3\pi}{2})$ ,  $r_t \notin$

$[0, 1] \cup E_1 \cup E_2$ ,  $r_t \rightarrow \infty$ , and if  $\alpha > 0$ , then for any given  $\varepsilon_1$  ( $0 < \varepsilon_1 < \min\{\alpha, 1 - \alpha\}$ ) and for sufficiently large  $r_k$ , we get by (2.10) that

$$\exp\{r_t^{\alpha-\varepsilon_1}\} < \nu(r_t) < \exp\{r_t^{\alpha+\varepsilon_1}\}; \quad (3.17)$$

if  $\alpha = 0$ , then by  $\sigma(f_0) = \infty$  and (2.11), we see that for any  $\varepsilon_2$  ( $0 < \varepsilon_2 < \frac{1}{2}$ ), and any large  $M$  ( $> 0$ ), we have, as  $r_t$  sufficiently large,

$$r_t^M < \nu(r_t) < \exp\{r_t^{\varepsilon_2}\}. \quad (3.18)$$

Since  $\theta_0$  may belong to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ , or  $(\frac{\pi}{2}, \frac{3\pi}{2})$ , or  $\{-\frac{\pi}{2}, \frac{\pi}{2}\}$ , we divide this proof into three cases.

**Case 1.** Suppose  $\theta_0 \in (-\frac{\pi}{2}, \frac{\pi}{2})$ . Then  $\cos\theta_0 > 0$ . We take  $\delta = \frac{1}{4}(\frac{\pi}{2} - |\theta_0|)$ . Thus  $[\theta_0 - \delta, \theta_0 + \delta] \subset (-\frac{\pi}{2}, \frac{\pi}{2})$ . By  $\theta_t \rightarrow \theta_0$ , we see that there is a constant  $N$  ( $> 0$ ), such that as  $t > N$ ,  $\theta_t \in [\theta_0 - \delta, \theta_0 + \delta]$ , and  $0 < \cos(|\theta_0| + \delta) \leq \cos\theta_t$ . By (3.14), we see that for any given  $\varepsilon_3$  ( $0 < \varepsilon_3 < \frac{1}{4(k+1)} \cos(|\theta_0| + \delta)$ ),

$$[T(2r_t, f_0)]^{k+1} \leq e^{\varepsilon_3(k+1)2r_t} \leq e^{\frac{1}{2} \cos(|\theta_0| + \delta)r_t} \leq e^{\frac{1}{2} \cos\theta_t r_t} \quad (3.19)$$

holds for  $n > N$ .

By (3.15) (3.16) and (3.19), we see that

$$\left(\frac{\nu(r_t)}{r_t}\right)^{k-s}(1 + o(1)) = \left|\frac{f_0^{(k-s)}(z_t)}{f_0(z_t)}\right| \leq M_0 [T(2r_t, f_0)]^{k+1} \leq M_0 e^{\frac{1}{2} \cos\theta_t r_t}. \quad (3.20)$$

By (1.11), we get

$$-\frac{f_0^{(s)}(z_t)}{f_0(z_t)} P_s(e^{z_t}) = \frac{f_0^{(k)}(z_t)}{f_0(z_t)} + \sum_{j=0, j \neq s}^{k-1} P_j(e^{z_t}) \frac{f_0^{(j)}(z_t)}{f_0(z_t)}. \quad (3.21)$$

Because  $\cos\theta_t > 0$  and (1.9), we get that

$$|P_s(e^{z_t})| = |a_{sm_s d_{sm_s}}| r^{d_{sm_s}} e^{m_s r_t \cos\theta_t} (1 + o(1)) \quad (3.22)$$

and

$$|P_j(e^{z_t})| \leq M_1 r_t^{d_{jm_j}} e^{m_j r_t \cos\theta_t} (1 + o(1)) \quad (j = 0, \dots, s-1, s+1, \dots, k-1). \quad (3.23)$$

Substituting (3.16) (3.22) and (3.23) into (3.21), we get for sufficiently large  $r_t$ ,

$$\begin{aligned} \left(\frac{\nu(r_t)}{r_t}\right)^s |a_{sm_s d_{sm_s}}| r^{d_{sm_s}} e^{m_s r_t \cos\theta_t} (1 + o(1)) &\leq \left(\frac{\nu(r_t)}{r_t}\right)^k (1 + o(1)) \\ &+ M_1 e^{m_j r_t \cos\theta_t} \sum_{j=0, j \neq s}^{k-1} r_t^{d_{jm_j}} \left(\frac{\nu(r_t)}{r_t}\right)^j (1 + o(1)) \end{aligned} \quad (3.24)$$

By (3.17) or (3.18),

$$\nu(r_t) > r_t^M > r_t. \quad (3.25)$$

By (3.20) (3.24) and (3.25), we get

$$\begin{aligned} |a_{sm_s d_{sm_s}}| r_t^{d_{sm_s}} e^{(m_s-m)r_t \cos \theta_t} (1 + o(1)) &\leq kM_1 \left(\frac{\nu(r_t)}{r_t}\right)^{k-s} r_t^d (1 + o(1)) \\ &\leq M_2 r_t^d e^{\frac{1}{2}r_t \cos \theta_t}. \end{aligned} \quad (3.26)$$

Since  $m_s - m \geq 1 > \frac{1}{2}$  and  $\cos \theta_t > 0$ , we see that (3.26) is a contradiction.

**Case 2.** Suppose  $\theta_0 \in (\frac{\pi}{2}, \frac{3\pi}{2})$ . By  $\cos \theta_0 < 0$  and  $\theta_t \rightarrow \theta_0$ , we see that for sufficiently large  $t$ , we have  $\cos \theta_t < 0$ . By (1.11) (3.16) and  $\cos \theta_t < 0$ , we get for sufficiently large  $r_t$ ,

$$e^{-m_s z_t} \frac{f^{(k)}(z_t)}{f(z_t)} = e^{-m_s z_t} P_{k-1}(e^{z_t}) \frac{f^{(k-1)}(z_t)}{f(z_t)} + \dots + e^{-m_s z_t} P_0(e^z). \quad (3.27)$$

From (1.9) and  $\cos \theta_t < 0$ , we get

$$\begin{aligned} |e^{-m_s z_t} P_j(e^{z_t})| &= |a_{jm_j}(z_t) e^{-(m_s-m_j)z_t} + \dots + a_{j1}(z_t) e^{-(m_s-1)z_t}| \\ &\leq B r_t^{d_{j1}} e^{-(m_s-1)r_t \cos \theta_t} (1 + o(1)). \end{aligned} \quad (3.28)$$

Substituting (3.16) (3.28) into (3.27), from (3.25) we have

$$e^{-m_s r_t \cos \theta_t} \nu(r_t) \leq M_3 r_t^d e^{-(m_s-1)r_t \cos \theta_t} (1 + o(1)). \quad (3.29)$$

If  $\alpha > 0$ , from (3.17) we have

$$\exp\{r_t^{\alpha-\varepsilon_3}\} e^{-m_s r_t \cos \theta_t} \leq M_3 r_t^d e^{-(m_s-1)r_t \cos \theta_t} (1 + o(1)). \quad (3.30)$$

Since  $\cos \theta_t < 0$  and  $\alpha < 1$ , we see (3.30) is a contradiction.

If  $\alpha = 0$ , from (3.18) we have

$$r_t^M e^{-m_s r_t \cos \theta_t} \leq M_3 r_t^d e^{-(m_s-1)r_t \cos \theta_t} (1 + o(1)). \quad (3.31)$$

Since  $\cos \theta_t < 0$ , we see (3.31) is also a contradiction.

**Case 3.** Suppose that  $\theta_0 = \frac{\pi}{2}$  or  $\theta_0 = -\frac{\pi}{2}$ . Since the proof for  $\theta_0 = -\frac{\pi}{2}$  is the same as the proof for  $\theta_0 = \frac{\pi}{2}$ , we only prove the case that  $\theta_0 = \frac{\pi}{2}$ . Since  $\theta_t \rightarrow \theta_0$ , for any given  $\varepsilon_4$  ( $0 < \varepsilon_4 < \frac{1}{10}$ ), we see that there is an integer  $K$  ( $> 0$ ), as  $t > K$ ,  $\theta_t \in [\frac{\pi}{2} - \varepsilon_4, \frac{\pi}{2} + \varepsilon_4]$ , and

$$z_t = r_t e^{i\theta_t} \in \bar{\Omega} = \{z : \frac{\pi}{2} - \varepsilon_4 \leq \arg \leq \frac{\pi}{2} + \varepsilon_4\}. \quad (3.32)$$

By Lemma 9, we see that there exist a subset  $E_3 \subset (1, \infty)$  having logarithmic measure  $lmE_3 < \infty$ , and a constant  $B > 0$  such that for

all  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_3$ , we have

$$\left| \frac{f_0^{(i)}(z)}{f_0^{(s)}(z)} \right| \leq B[T(2r, f_0^{(s)})]^{k-s+1} \quad (i = s+1, \dots, k). \quad (3.33)$$

Now we consider the property of  $f_0(re^{i\theta})$  on a ray  $\arg z = \theta \in \overline{\Omega} \setminus \{\frac{\pi}{2}\}$ . If  $\theta \in [\frac{\pi}{2} - \varepsilon_3, \frac{\pi}{2})$ , then  $\cos \theta > 0$ .

Since  $\sigma_2(f_0) < 1$ , we get that  $f_0$  satisfy (3.14). From  $T(r, f_0^{(s)}) \leq (s+1)T(r, f_0)$ , we get that  $f_0^{(s)}$  also satisfies (3.14). So for any given  $\varepsilon_5$  ( $0 < \varepsilon_5 < \frac{1}{4(k-s+1)} \cos \theta$ ), we have

$$[T(2r, f_0^{(s)})]^{k-s+1} \leq e^{\varepsilon_5(k-s+1)2r} \leq e^{\frac{1}{2}r \cos \theta}. \quad (3.34)$$

We assert that  $|f_0^{(s)}(re^{i\theta})|$  is bounded on the ray  $\arg z = \theta \in [\frac{\pi}{2} - \varepsilon_3, \frac{\pi}{2})$ . If  $|f_0^{(s)}(re^{i\theta})|$  is unbounded on the ray  $\arg z = \theta$ , then by Lemma 3, there exists a sequence  $\{y_j = R_j e^{i\theta}\}$  such that as  $R_j \rightarrow \infty$ ,  $f_0^{(s)}(y_j) \rightarrow \infty$  and

$$\left| \frac{f_0^{(i)}(y_j)}{f_0^{(s)}(y_j)} \right| \leq R_j^{s-i}(1+o(1)) \quad i = 0, \dots, s-1. \quad (3.35)$$

By Remark 2 and  $f_0^{(s)}(y_j) \rightarrow \infty$ , we know that  $|y_j| = R_j \notin E_3$ . By (3.33) and (3.34), we have for sufficiently large  $j$ ,

$$\left| \frac{f_0^{(j)}(y_j)}{f_0^{(s)}(y_j)} \right| \leq [T(2R_j, f_0^{(s)})]^{k-s+1} \leq B e^{\frac{1}{2}R_j \cos \theta} \quad j = s+1, \dots, k. \quad (3.36)$$

Substituting (3.35) and (3.36) into (1.11), we get

$$\begin{aligned} |a_{sm_s d_{sm_s}}| R_j^{d_{sm_s}} e^{m_s R_j \cos \theta} (1+o(1)) &= | -P_s(e^{y_j}) | \\ &\leq s M_4 R_j^{d_2} e^{m R_j \cos \theta} (1+o(1)) \\ &+ (k-s) M_5 e^{\frac{1}{2}R_j \cos \theta} R_j^{d_1} e^{m R_j \cos \theta} (1+o(1)). \end{aligned} \quad (3.37)$$

Since  $m_s > m + \frac{1}{2}$  and  $\cos \theta > 0$ , we get (3.37) is a contradiction.

Hence  $|f_0^{(s)}(re^{i\theta})|$  is bounded on the ray  $\arg z = \theta \in [\frac{\pi}{2} - \varepsilon_4, \frac{\pi}{2})$ . Set  $|f_0^{(s)}(re^{i\theta})| \leq M_6$ , then on the ray  $\arg z = \theta \in [\frac{\pi}{2} - \varepsilon_4, \frac{\pi}{2})$ ,

$$|f_0(re^{i\theta})| \leq M_7 r^s. \quad (3.38)$$

On the other hand, since  $r_t \notin [0, 1] \cup E_1 \cup E_2 \cup E_3$ , by Lemma 10, and (3.17) or (3.18), we see that for sufficiently large  $r$

$$\log M(r_t, f_0) \geq (\nu(r_t))^{\frac{1}{2}} \geq r_t^{\frac{M}{2}},$$

where  $M(> 1)$  is some constant. Since  $\{z_t\}$  satisfies  $|f_0(z_t)| = M(r_t, f_0)$ ,

$$|f_0(z_t)| \geq \exp\left\{r_t^{\frac{M}{2}}\right\}. \quad (3.39)$$

By (3.38) and (3.39), we see that for sufficiently large  $t$ ,  $\theta_t \notin [\frac{\pi}{2} - \varepsilon_3, \frac{\pi}{2}]$ , i.e.,

$$\theta_t \in [\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon_3]. \quad (3.40)$$

Thus there are two subcases: Subcase (i) there are infinitely many  $\theta_t$  in  $(\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon_3]$ ; Subcase (ii) there are only finitely many  $\theta_t$  in  $(\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon_3]$ . Now consider Subcase (i), all  $\theta_t \in (\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon_3]$  form a subsequence  $\theta_{t_j}$  of  $\theta_t$  and a corresponding subsequence  $z_{t_j} = r_{t_j} e^{i\theta_{t_j}}$  of  $z_t$ . For a subsequence  $\{z_{t_j}\} \subset \{z : \frac{\pi}{2} < \arg z = \theta \leq \frac{\pi}{2} + \varepsilon_3\}$ , using a similar method to that in the proof of Case 2, we can get a contradiction.

Consider Subcase (ii), we see that for sufficiently large  $t$ ,

$$\theta_t = \frac{\pi}{2}.$$

Thus, for sufficiently large  $t$ ,  $\cos \theta_t = 0$  and

$$\begin{aligned} |P_j(e^{z_t})| &= |a_{jm_j}(z_t)e^{m_j z_t} + \cdots + a_{j1}(z_t)e^{z_t}| \\ &\leq |a_{jm_j}(z_t)| + \cdots + |a_{j1}(z_t)| \leq M_8 r^d, \end{aligned} \quad (3.41)$$

where  $j = 0, \dots, k-1$  and  $M_8$  is a constant.

By (1.11) (3.16) (3.17) (or (3.18) ) and (3.41), we get that

$$\left| -\left(\frac{\nu(r_t)}{z_t}\right)^k (1 + o(1)) \right| = \left| -\frac{f_0^{(k)}(z_t)}{f_0(z_t)} \right| \leq k M_9 r_t^{d_3} \left(\frac{\nu(r_t)}{r_t}\right)^{k-1} (1 + o(1)),$$

i.e.,

$$\nu(r_t)(1 + o(1)) \leq k M_9 r_t^{d_3+1} (1 + o(1)).$$

By (3.17) (or (3.18) ), this is also a contradiction.

So we have  $\sigma_2(f) = \alpha = 1$ .

#### 4. PROOF OF THEOREM 2

From Theorem 1, we get  $\sigma_2(f) = 1$ .

Let  $g = f - z$ , then  $f = g + z$ . Substituting it into (1.11), we have

$$g^{(k)} + P_{k-1}(e^z)g^{(k-1)} + \cdots + P_0(e^z)g = -zP_0(e^z) - P_1(e^z). \quad (4.1)$$

Since  $zP_0(e^z) - P_1(e^z) \not\equiv 0$ , from Lemma 11 and  $\sigma_2(g) = 1$  we conclude  $\bar{\lambda}_2(g) = \lambda_2(g) = \sigma_2(g) = 1$ . So we have  $\bar{\tau}_2(f) = \tau_2(f) = \sigma_2(f) = 1$ .

#### 5. PROOF OF THEOREM 3

From Theorem 1, we get  $\sigma(f) = \infty$ .

(i) Let  $g = f - z$ , then  $f = g + z$ . Substituting it into (1.6), we have

$$g'' + P(e^z)g' + Q(e^z)g = -P(e^z) - Q(e^z)z. \quad (5.1)$$

Since  $n \neq s$ , we get that  $-P(e^z) - Q(e^z)z \not\equiv 0$ . From Lemma 11, we get  $\lambda(g) = \sigma(g) = \sigma(f) = \infty$  and  $\lambda_2(g) = \sigma_2(g) = \sigma_2(f) = \sigma$ . i.e.,

$\lambda(f - z) = \infty$  and  $\lambda_2(f - z) = \sigma$ .

(ii) Differentiating both sides of (1.6), we get that

$$f''' + P(e^z)f'' + (P'(e^z) + Q(e^z))f' + Q'(e^z)f = 0. \quad (5.2)$$

By (1.6), we get that

$$f = -\frac{f'' + P(e^z)f'}{Q(e^z)}. \quad (5.3)$$

Substituting (5.3) into (5.2), we get

$$f''' + [P(e^z) - \frac{Q'(e^z)}{Q(e^z)}]f'' + [P'(e^z) + Q(e^z) - \frac{Q'(e^z)}{Q(e^z)}P(e^z)]f' = 0. \quad (5.4)$$

Let  $g = f' - z$ , then  $f' = g + z$ ,  $f'' = g' + 1$ ,  $f''' = g''$ . Substituting these into (5.4), we get that

$$\begin{aligned} g'' + [P(e^z) - \frac{Q'(e^z)}{Q(e^z)}]g' + [P'(e^z) + Q(e^z) - \frac{Q'(e^z)}{Q(e^z)}P(e^z)]g \\ = \frac{Q'(e^z)}{Q(e^z)} - P(e^z) - [P'(e^z) + Q(e^z) - \frac{Q'(e^z)}{Q(e^z)}P(e^z)]z = h(z) \end{aligned} \quad (5.5)$$

Next we prove that  $h(z) \not\equiv 0$ .

If  $h(z) \equiv 0$ , then  $\frac{Q'(e^z)}{Q(e^z)} - P(e^z) \equiv [P'(e^z) + Q(e^z) - \frac{Q'(e^z)}{Q(e^z)}P(e^z)]z$ .

Since  $Q(z) \not\equiv 0$ , we have

$$Q'(e^z) - P(e^z)Q(e^z) \equiv [P'(e^z)Q(e^z) + Q^2(e^z) - Q'(e^z)P(e^z)]z. \quad (5.6)$$

If  $n < s$ , taking  $z = r$ , we have

$$e^{2sr}(1 + o(1)) \leq e^{(n+s)r}(1 + o(1)).$$

This is a contradiction.

So we have  $h(z) \not\equiv 0$ .

From Lemma 11, we get  $\lambda(g) = \sigma(g) = \sigma(f' - z) = \sigma(f) = \infty$  and  $\lambda_2(g) = \sigma_2(g) = \sigma_2(f' - z) = \sigma_2(f) = \sigma$ . i.e.,  $\lambda(f' - z) = \infty$  and  $\lambda_2(f' - z) = \sigma$ .

If  $n > s$ , taking  $z = r$ , we have

$$P(e^r) = a_n(r)e^{nr} + \dots + a_1(r)e^r, \quad Q(e^r) = b_s(r)e^{sr} + \dots + b_1(r)e^r.$$

We get

$$P'(e^r) = (a'_n(r) + na_n(r))e^{nr} + (a'_{n-1}(r) + (n-1)a_{n-1}(r))e^{(n-1)r} + \dots$$

and

$$Q'(e^r) = (b'_s(r) + sb_s(r))e^{sr} + (b'_{s-1}(r) + (s-1)b_{s-1}(r))e^{(s-1)r} + \dots$$

So we have

$$|P(e^r)Q(e^r) + P'(e^r)Q(e^r)r - P(e^r)Q'(e^r)r| = |(n-s)ra_n(r)b_s(r) + [a_n(r)b_s(r) + (a'_n(r)b_s(r) - b'_s(r)a_n(r))r]|e^{(n+s)r}(1+o(1)) \quad (5.7)$$

Since  $a_n(r)$ ,  $b_s(r)$  are polynomials and  $n > s$ , we get

$$\deg((n-s)ra_n(r)b_s(r)) > \deg[a_n(r)b_s(r) + (a'_n(r)b_s(r) - b'_s(r)a_n(r))r].$$

So we have

$$\begin{aligned} & |(n-s)ra_n(r)b_s(r) + [a_n(r)b_s(r) + (a'_n(r)b_s(r) - b'_s(r)a_n(r))r]| \\ & = Mr^{d_1}(1+o(1)) \neq 0. \end{aligned}$$

From (5.6), we have

$$\begin{aligned} Mr^{d_1}e^{(n+s)r}(1+o(1)) & = |P(e^r)Q(e^r) + P'(e^r)Q(e^r)r - P(e^r)Q'(e^r)r| \\ & = |Q'(e^r) - Q^2(e^r)r| \leq Br^{d_2}e^{2sr}(1+o(1)). \quad (5.8) \end{aligned}$$

Since  $n > s$ , we get a contradiction.

So we also have  $h(z) \neq 0$ .

From Lemma 11, we get  $\lambda(g) = \sigma(g) = \sigma(f' - z) = \sigma(f) = \infty$  and  $\lambda_2(g) = \sigma_2(g) = \sigma_2(f' - z) = \sigma_2(f) = \sigma$ . i.e.,  $\lambda(f' - z) = \infty$  and  $\lambda_2(f' - z) = \sigma$ .

#### ACKNOWLEDGEMENTS

The authors would like to thank the referee for his/her valuable suggestions. This work was supported by the NSF of Shandong Province, No.ZR2010AM030, P. R. China and the NNSF of China (No. 11171013 & No.11041005).

#### REFERENCES

- [1] B. Belaïdi, *Growth and oscillation theory of solutions of some linear differential equations*, Mat. Vesnik., **60** (2008), 233–246.
- [2] W. J. Chen and J. F. Xu *Growth of meromorphic solutions of higher-order linear differential equations*, Electronic Journal of Qualitative Theory of Differential Equations., **1** (2009), 1–13.
- [3] Z. X. Chen, *Zeros of meromorphic solutions of higher order linear differential equations*, Analysis., **14** (1994), 425–438.
- [4] Z. X. Chen, *The fixed points and hyper-order of solutions of second order complex differential equations(in Chinese)*, Acta Math. Sci. Ser. A Chin. Ed., **20** (2000), 425–432.
- [5] Z. X. Chen, *On the growth of solutions of a class of higher order differential equations*, Chin Ann of Math., **24B(4)** (2003), 501–508.
- [6] Z. X. Chen and K. H. Shon, *The hyper order of solutions of second order differential equations and subnormal solutions of periodic equations*, Taiwanese J. Math. **14** (2)(2010), 611–628.



- [7] Z. X. Chen and K. H. Shon, *On subnormal solutions of periodic differential equations*, Abstract and Applied Analysis., vol. 2010, Article ID 170762, 16 pages, 2010.
- [8] Z. X. Chen and K. H. Shon, *Numbers of subnormal solutions for higher order periodic differential equations*, Acta Mathematica Sinica, English Series., **27** (9) (2011), 1753–1768.
- [9] Z. X. Chen and C. C. Yang, *Some further results on the zeros and growths of entire solutions of second order linear differential equations*, Kodai Math. J., **22** (1999), 273–285.
- [10] G. Gundersen, *Estimates for the logarithmic derivative of a meromorphic function, plus similar estimates*, J. London Math. Soc., **37** (1988), 88–104.
- [11] G. Gundersen, *Finite order solutions of second order linear differential equations*, Trans. Amer. Math. Soc., **305** (1988), 415–429.
- [12] G. Gundersen and M. Steinbart, *Subnormal solutions of second order linear differential equation with periodic coefficients*, Results in Math., **25** (1994), 270–289.
- [13] G. Jank and L. Volkmann, *Einführung in die Theorie der ganzen und meromorphen Funktionen mit Anwendungen auf Differentialgleichungen*, Birkhäuser, Basel-Boston, 1985
- [14] J. K. Langley, *Integer points of entire functions*, Bull. London Math. Soc., **38** (2006) 239–249.
- [15] I. Laine, *Nevanlinna theory and complex differential equations*, W. de Gruyter, Berlin, (1993)
- [16] K. Liu and L. Z. Yang, *On the complex oscillation of higher order linear differential equations*, Bull. Korean Math. Soc., **46** (2009), 607–615.
- [17] H. Wittich, *Subnormale Lösungen der Differentialgleichung  $\omega'' + p(e^z)\omega' + q(e^z)\omega = 0$* , Nagoya Math. J., **30** (1967), 29–37.
- [18] J. F. Xu, *On the growth of meromorphic solutions of complex algebraic differential equations*, Journal of Shandong University(Natural Science), **45** (2010), 91–93.
- [19] L. Z. Yang, *Solution of a differential equation and its applications*, Kodai Math. J., **22** (1999), 458–464.
- [20] H. X. Yi and C. C. Yang, *Uniqueness theory of meromorphic functions*. Science Press, Beijing, (1995)
- [21] G.W. Zhang and A. Chen, *Fixed points of the derivative and  $k$ -th power solutions of complex linear differential equations in the unit disc*, Electronic Journal of Qualitative Theory of Differential Equations, **48** (2009), 1–9.

(Received September 7, 2012)

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG PROVINCE,  
250100, P.R.CHINA

*E-mail address:* nanli32787310@gmail.com

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG PROVINCE,  
250100, P.R.CHINA

*E-mail address:* lzyang@sdu.edu.cn