



NONCOMMUTATIVE DEFORMATION OF INSTANTONS AND VORTEXES

AKIFUMI SAKO

Communicated by Akira Yoshioka

Abstract. We study noncommutative (NC) instantons and vortexes. At first, we construct instanton solutions which are deformations of instanton solutions on commutative Euclidean four-space. We show that the instanton numbers of these NC instanton solutions coincide with the commutative solutions. Next, we also deform vortex solutions similarly and we show that their vortex numbers are unchanged under the NC deformation.

1. Introduction

Instanton connections in the four dimensional Yang-Mills theory are defined by

$$F^+ = \frac{1}{2}(1 + *)F = 0 \tag{1}$$

where F is a curvature two-form and $*$ is the Hodge star operator.

The NC instanton solutions were constructed with the ADHM method in [1, 15]. The ADHM construction which generate the instanton $U(N)$ gauge field require a pair of the two complex vector spaces $V = \mathbb{C}^k, W = \mathbb{C}^N$. Here k is an integer. Introduce $B_1, B_2 \in \text{Hom}(V, V), I \in \text{Hom}(W, V)$ and $J \in \text{Hom}(V, W)$ called ADHM data such that

$$\mu_{\mathbf{R}} := [B_1, B_1^\dagger] + [B_2, B_2^\dagger] + II^\dagger - J^\dagger J = \zeta \text{Id}_k \tag{2}$$

$$\mu_{\mathbf{C}} := [B_1, B_2] + IJ = 0. \tag{3}$$

Here ζ is a NC parameter and its detail will appear in the following. Using these ADHM data we can construct NC instanton and call it NC ADHM instanton in the following. NC ADHM instantons and deformed instantons from the commutative ADHM construction is unknown.

It was also shown that instanton numbers of NC ADHM instantons do not depend on the NC parameter [4, 5, 9, 16, 17]. These results may imply that the instanton