




Gâteaux Differentiability

B. Zlatanov

vol. 8, iss. 4, art. 113, 2007

[Title Page](#)

[Contents](#)



Page 1 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

# UPPER ESTIMATES FOR GÂTEAUX DIFFERENTIABILITY OF BUMP FUNCTIONS IN ORLICZ-LORENTZ SPACES

**B. ZLATANOV**

Department of Mathematics and Informatics  
Plovdiv University,  
24 “Tzar Assen” str.  
Plovdiv, 4000, Bulgaria  
EMail: [bzlatanov@gmail.com](mailto:bzlatanov@gmail.com)

*Received:* 29 January, 2007

*Accepted:* 17 November, 2007

*Communicated by:* C.P. Niculescu

*2000 AMS Sub. Class.:* 46B25, 40E30.

*Key words:* Orlicz function, Orlicz–Lorentz space, Smooth bump functions.

*Abstract:* Upper estimates for the order of Gâteaux smoothness of bump functions in Orlicz–Lorentz spaces  $d(w, M, \Gamma)$ ,  $\Gamma$  uncountable, are obtained. The best possible order of Gâteaux differentiability in the class of all equivalent norms in  $d(w, M, \Gamma)$  is found.

*Acknowledgements:* Research is partially supported by National Fund for Scientific Research of the Bulgarian Ministry of Education and Science, Contract MM-1401/04.

# Contents

1	Introduction	3
2	Preliminaries	5
3	Main Result	8
4	Auxiliary Lemmas	9
5	Gâteaux Differentiability of Bumps in $d(w, M, \Gamma)$ and $d(w, p, \Gamma)$	16



Gâteaux Differentiability

B. Zlatanov

vol. 8, iss. 4, art. 113, 2007

Title Page

Contents



Page 2 of 18

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

## 1. Introduction

The existence of higher order Fréchet smooth norms and bump functions and its impact on the geometrical properties of a Banach space have been subject to many investigations beginning with the classical result for  $L_p$ -spaces in [1] and [6]. An extensive study and bibliography may be found in [2]. As any negative result on the existence of Gâteaux smooth bump functions immediately applies to the problem of existence of Fréchet smooth bump functions and norms, the question arises of estimating the best possible order of Gâteaux smoothness of bump functions in a given Banach space. A variational technique (the Ekeland variational principle) was applied in [2] to show that in  $\ell_1(\Gamma)$ ,  $\Gamma$  uncountable, there is no continuous Gâteaux differentiable bump function. Following the same idea and using Stegall's variational principle, an extension of this result to Banach spaces with uncountable unconditional basis was given in [4] and to Banach spaces with uncountable symmetric basis in [9]. As an application in [4] it was shown that in  $\ell_p(\Gamma)$ ,  $\Gamma$  uncountable, there is no continuous  $p$ -times Gâteaux differentiable bump function when  $p$  is odd and there is no continuous  $([p] + 1)$ -times Gâteaux differentiable bump function in the case  $p \notin \mathbb{N}$ . This is essentially different from the case  $\ell_p(\mathbb{N})$ ,  $p$ -odd, where equivalent  $p$ -times Gâteaux differentiable and even uniformly Gâteaux differentiable norms are constructed (see [10] and [8] respectively). As examples of the main result in [9], Orlicz  $\ell_M(\Gamma)$  and Lorentz  $d(w, p, \Gamma)$ ,  $\Gamma$  uncountable are considered and estimates for the order of Gâteaux smoothness of bump functions are obtained. Recently a deep result on embedding of  $\ell_p$  spaces in Orlicz–Lorentz sequence spaces  $d_0(p, M)$  have been found in [5]. It is shown there that  $\ell_p \hookrightarrow d_0(w, M)$  iff  $\ell_p \hookrightarrow h_M$  iff  $p \in [\alpha_M, \beta_M]$ . From this result naturally arises the question of finding upper estimates for the order of Gâteaux smoothness of bump functions in Orlicz–Lorentz spaces.

It is worthwhile to mention that results about differentiability of bump functions



Gâteaux Differentiability

B. Zlatanov

vol. 8, iss. 4, art. 113, 2007

Title Page

Contents



Page 3 of 18


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-575b



Title Page

Contents



Page 4 of 18

Go Back

Full Screen

Close

in  $\ell_p(\Gamma)$  cannot be used directly for  $\ell_M(\Gamma)$  and  $d(w, p, \Gamma)$ . Indeed, in [3] it is proved that  $\ell_p(\mathbf{A})$  is isomorphic to a subspace of  $d(w, p, \Gamma)$  iff  $\mathbf{A}$  is countable. On the other hand  $\ell_M(\Gamma)$  for  $M \equiv t^p(1 + |\log t|)^q$  at zero,  $p \geq 1$ ,  $q \neq 0$ , contains an isomorphic copy of  $\ell_p(\mathbf{A})$  iff  $\mathbf{A}$  is countable. The problem of embedding  $\ell_p(\mathbf{A})$  or  $\ell_M(\mathbf{A})$  into  $d(w, M, \Gamma)$ ,  $\Gamma$  uncountable is open.


In this note we give one new application of the main result of [9] in Orlicz–Lorentz spaces  $d(w, M, \Gamma)$ ,  $\Gamma$  uncountable for finding upper estimates for the order of Gâteaux smoothness of bump functions.

Let  $U$  be an open set in a Banach space  $X$  and let  $f : U \rightarrow \mathbb{R}$  be continuous. Following [4] we shall say that  $f$  is  $G_{\omega, k}^0$ -smooth,  $k \in \mathbb{N}$  in  $U$  for some  $\omega : (0, 1] \rightarrow \mathbb{R}^+$ ,  $\lim_{t \rightarrow 0} t^{-k} \omega(t) = 0$  if for any  $x \in U$ ,  $y \in X$  the representation holds

$$f(x + ty) = f(x) + \sum_{i=1}^k \frac{t^i}{i!} f^{(i)}(x)(y^i) + R_f^k(x, y, t),$$

where  $f^{(i)}$ ,  $i = 1, 2, \dots, k$  are  $i$ -linear bounded symmetric forms on  $X$  and  $\lim_{t \rightarrow 0} \frac{|R_f^k(x, y, t)|}{\omega(|t|)} = 0$ .

If  $U = X$  we use the notation  $G_{\omega, [p]}^0$  instead of  $G_{\omega, [p]}^0(X)$  and  $G_k$ ,  $k \in \mathbb{N}$  for the set of all continuous  $k$ -times Gâteaux differentiable functions on  $X$ , for which  $\lim_{t \rightarrow 0} |R_f^k(x, y, t)|/|t|^k = 0$ . We say that the norm  $\|\cdot\|$  in  $X$  is  $k$ -times Gâteaux differentiable if it is from the class  $G_k(X \setminus \{0\})$ .

[Title Page](#)[Contents](#)

Page 5 of 18

[Go Back](#)[Full Screen](#)[Close](#)journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-575b

## 2. Preliminaries

We use the standard Banach space terminology from [7]. Let us recall that an Orlicz function  $M$  is an even, continuous, non-decreasing convex function such that  $M(0) = 0$  and  $\lim_{t \rightarrow \infty} M(t) = \infty$ . We say that  $M$  is a non-degenerate Orlicz function if  $M(t) > 0$  for every  $t > 0$ .

A weight sequence  $w = \{w_n\}_{n=1}^{\infty}$  is a positive decreasing sequence such that  $w_1 = 1$  and  $\lim_{n \rightarrow \infty} W(n) = \infty$ , where  $W(n) = \sum_{j=1}^n w_j$ , for any  $n \in \mathbb{N}$ .

The Orlicz–Lorentz space  $d(w, M, \Gamma)$  is the space of all real functions  $x = x(\alpha)$  defined on the set  $\Gamma$ , for which

$$I(\lambda x) = \sup \left\{ \sum_{i=1}^{\infty} w_i M(\lambda x(\alpha_i)) \right\} < \infty$$

for some  $\lambda > 0$ , where the supremum is taken over all sequences  $\{\alpha_i\}_{i=1}^{\infty}$  of different elements on  $\Gamma$ . There exists a sequence  $\{\alpha_i^*\}_{i=1}^{\infty}$ , such that  $|x(\alpha_1^*)| \geq |x(\alpha_2^*)| \geq \dots \geq |x(\alpha_i^*)| \geq \dots$ ,  $\lim_{i \rightarrow \infty} x(\alpha_i^*) = 0$ ,  $|x(\alpha^*)| = 0$  if  $\alpha \neq \alpha_i^*$  for  $i \in \mathbb{N}$  and  $I(\lambda x) = \sum_{i=1}^{\infty} w_i M(\lambda x(\alpha_i^*))$ . The space  $d(w, M, \Gamma)$ , equipped with the Luxemburg norm:


$$\|x\| = \inf \left\{ \lambda > 0 : I\left(\frac{x}{\lambda}\right) \leq 1 \right\}$$

is a Banach space.

By  $\text{supp } x$  we denote the set  $\{\alpha \in \Gamma : x(\alpha) \neq 0\}$ .

The symbol  $e_\gamma$ ,  $\gamma \in \Gamma$  will stand for the unit vectors.

If  $M(u) = u^p$ ,  $1 \leq p < \infty$  then  $d(w, M, \Gamma)$  is the Lorentz space  $d(w, p, \Gamma)$ . If  $w_i = 1$  for every  $i \in \mathbb{N}$  then  $d(w, M, \Gamma)$  is the Orlicz space  $\ell_M(\Gamma)$ . In this case we use the notation  $I(x) = \widetilde{M}(x)$



[Title Page](#)

[Contents](#)



Page 6 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

To every Orlicz function  $M$  the following numbers are associated:

$$\alpha_M = \sup \left\{ p > 0 : \sup_{0 < u, v \leq 1} \frac{M(uv)}{u^p M(v)} < \infty \right\},$$

$$\beta_M = \inf \left\{ q > 0 : \inf_{0 < u, v \leq 1} \frac{M(uv)}{u^q M(v)} > 0 \right\}.$$

We consider only spaces generated by an Orlicz function  $M$  satisfying the  $\Delta_2$ -condition at zero, i.e.,  $\beta_M < \infty$ , which implies of course that

$$(2.1) \quad M(uv) \geq u^q M(v), \quad u, v \in [0, 1]$$

for some  $q > \beta_M$  (see [7]).

Finally we mention that the unit vectors  $\{e_\gamma\}_{\gamma \in \Gamma}$  form a symmetric basis of  $d(w, M, \Gamma)$  with symmetric constant 1, which is boundedly complete [5], [7].

For a function  $g : (0, 1] \rightarrow \mathbb{R}^+$  denote:

$$d_M(g) = \sup \left\{ \frac{M(uv)}{g(u)M(v)} : u, v \in (0, 1] \right\}.$$

Let us recall a well known definition. Let  $X$  have symmetric basis  $\{e_\gamma\}_{\gamma \in \Gamma}$  with a symmetric constant 1 and let  $z \in X$ ,  $z \neq 0$ ,  $z = \sum_{i=1}^{\infty} u_i e_{\gamma_i}$ ,  $\gamma_i \neq \gamma_j$  for  $i \neq j$ . A sequence  $\{z_k\}_{k=1}^{\infty}$ ,  $z_k = \sum_{i=1}^{\infty} u_i e_{\alpha_{i,k}}$ ,  $\alpha_{i,k} \neq \alpha_{j,l}$  for  $(i, k) \neq (j, l)$ ,  $\alpha_{i,k} \in \Gamma$  is called a block basis generated by the vector  $z$ .

We will apply a general result for upper estimates for the order of Gâteaux smoothness of bump functions in a Banach space with a symmetric, boundedly complete basis with a symmetric constant 1, obtained in [9].



**Theorem 2.1.** [9] Let  $X$  be a Banach space, let  $\{e_\gamma\}_{\gamma \in \Gamma}$ ,  $\#\Gamma > \aleph_0$  be a symmetric, boundedly complete basis in  $X$  with a symmetric constant 1 and let:

$$\lim_{n \rightarrow \infty} \left\| \sum_{j=1}^n z_j \right\| n^{-\frac{1}{k}} = 0$$

for every  $z \in X$ .

Let  $\omega : [0, 1] \rightarrow \mathbb{R}^+$  be such that for every  $x \in X$  there exist  $y \in X$ ,  $\text{supp } y \cap \text{supp } x = \emptyset$  and a sequence  $t_n \searrow 0$ , which satisfy the inequality

$$\|x + t_n y\| - \|x\| \geq \omega(t_n), \quad n \in \mathbb{N}.$$

Then in  $X$  there is no continuous:

- (i)  $G_{\omega, k}^0$ -smooth bump when  $\omega(t) = o(t^k)$ ;
- (ii)  $G_{\omega, k+1}^0$ -smooth bump when  $\omega(t) = o(t^{k+1})$ ,  $k$ -even;
- (iii)  $k$ -times Gâteaux differentiable bump if  $\omega(t) = t^k$ ;
- (iv)  $(k+1)$ -times Gâteaux differentiable bump if  $\omega(t) = t^{k+1}$ ,  $k$ -even.

Title Page

Contents



Page 7 of 18

Go Back

Full Screen

Close

### 3. Main Result

**Theorem 3.1.** *Let  $M$  be an Orlicz function. If  $f$  is a continuous  $k$ -times Gâteaux differentiable bump function in  $d(w, M, \Gamma)$ , then*

$$k \leq E_M = \begin{cases} [\alpha_M], & d_M(t^{\alpha_M}) < \infty \\ \alpha_M - 1, & \alpha_M \in \mathbb{N}, d_M(t^{\alpha_M}) = \infty. \end{cases}$$



**Gâteaux Differentiability**

B. Zlatanov

vol. 8, iss. 4, art. 113, 2007

Title Page

Contents



Page 8 of 18


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Title Page

Contents

◀ ▶

◀ ▶

Page 9 of 18

Go Back

Full Screen

Close

## 4. Auxiliary Lemmas

To apply Theorem 2.1 for  $d(w, M, \Gamma)$  we need the following lemmas.

**Lemma 4.1.** *Let  $p \geq 1$  and let  $M$  be an Orlicz function satisfying the conditions  $\lim_{t \rightarrow 0} \frac{M(t)}{t^p} = 0$ ,  $d_M(t^p) = c < \infty$ . Then every block basis  $\{z_j\}_{j=1}^\infty$  of the unit vector basis  $\{e_\gamma\}_{\gamma \in \Gamma}$  in  $d(w, M, \Gamma)$ , generated by one vector, satisfies*

$$\lim_{n \rightarrow \infty} \left\| \sum_{j=1}^n z_j \right\| n^{-\frac{1}{p}} = 0.$$

*Proof.* Let  $z = \sum_{i=1}^\infty u_i e_{\gamma_i} \in d(w, M, \Gamma)$ . Let  $\{e_{j,i}\}_{i=1}^\infty$ ,  $j \in \mathbb{N}$  be disjoint subsets of  $\{e_\gamma\}_{\gamma \in \Gamma}$ . Then we define  $z_j = \sum_{i=1}^\infty u_i e_{j,i}$ . Let  $\mu(t) = \frac{M(t)}{t^p}$ . It follows that  $\lim_{t \rightarrow 0} \mu(t) = 0$  and  $\mu(t_1) \leq c\mu(t_2)$  for every  $0 < t_1 < t_2 \leq 1$ . Let  $\lambda_n(z) = \sum_{j=1}^n z_j$ . Then

$$I(\lambda_n(z)) = \sum_{i=1}^\infty \sum_{j=n(i-1)+1}^{ni} w_j M(u_i^*).$$

For every  $\varepsilon > 0$  there exists  $m \in \mathbb{N}$  such that

$$\sum_{i=m+1}^\infty \sum_{j=n(i-1)+1}^{ni} w_j M(u_i^*) < \frac{\varepsilon}{2c}.$$

By the definition of the function  $\mu$  it follows that

$$\begin{aligned} \sum_{i=1}^\infty \sum_{j=n(i-1)+1}^{ni} w_j |u_i^*|^p \mu \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right) &= \|\lambda_n(z)\|^p \sum_{i=1}^\infty \sum_{j=n(i-1)+1}^{ni} w_j M \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right) \\ &= \|\lambda_n(z)\|^p. \end{aligned}$$



[Title Page](#)

[Contents](#)



Page 10 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

Using the inequality

$$1 = I \left( \frac{\lambda_n(z)}{\|\lambda_n(z)\|} \right) = \sum_{i=1}^{\infty} \sum_{j=n(i-1)+1}^{ni} w_j M \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right) \geq \sum_{j=1}^n w_j M \left( \frac{u_1^*}{\|\lambda_n(z)\|} \right)$$

we get that  $\lim_{n \rightarrow \infty} \|\lambda_n(z)\|^{-1} = 0$ .

For every  $m \in \mathbb{N}$  we have

$$\sum_{i=1}^m \sum_{j=n(i-1)+1}^{ni} \frac{w_j |u_i^*|^p}{n} \mu \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right) \leq \frac{w_1 + w_2 \dots w_n}{n} \sum_{i=1}^m |u_i^*|^p \mu \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right).$$

Because  $\lim_{j \rightarrow \infty} w_j = 0$  it follows that for every  $\varepsilon > 0$  and every  $m \in \mathbb{N}$  there exists  $N \in \mathbb{N}$  such that for any  $n \geq N$  holds

$$\sum_{i=1}^m \sum_{j=n(i-1)+1}^{ni} \frac{w_j |u_i^*|^p}{n} \mu \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right) \leq \frac{\varepsilon}{2}.$$

On the other hand for all  $n \in \mathbb{N}$  such that  $\|\lambda_n(z)\|^{-1} \leq 1$  we can write the chain of inequalities

$$\begin{aligned} \sum_{i=m+1}^{\infty} \sum_{j=n(i-1)+1}^{ni} \frac{w_j |u_i^*|^p}{n} \mu \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right) &\leq c \sum_{i=m+1}^{\infty} \sum_{j=n(i-1)+1}^{ni} \frac{w_j |u_i^*|^p}{n} \mu(u_i^*) \\ &\leq c \sum_{i=m+1}^{\infty} \sum_{j=n(i-1)+1}^{ni} \frac{w_j}{n} M(u_i^*) \\ &\leq \frac{\varepsilon}{2n}. \end{aligned}$$



Title Page

Contents



Page 11 of 18

Go Back

Full Screen

Close

Therefore for every  $\varepsilon > 0$  and  $n \geq N$  we have

$$\frac{\|\lambda_n(z)\|^p}{n} = \sum_{i=1}^{\infty} \sum_{j=n(i-1)+1}^{ni} \frac{w_j |u_i^*|^p}{n} \mu \left( \frac{u_i^*}{\|\lambda_n(z)\|} \right) \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2n} < \varepsilon$$

and thus

$$\lim_{n \rightarrow \infty} \left\| \sum_{j=1}^n z_j \right\| n^{-\frac{1}{p}} = 0$$

□

**Lemma 4.2.** *Let  $d_M(\omega) = \infty$  then for any  $x \in d(w, M, \Gamma)$  there exist  $y \in d(w, M, \Gamma)$  with  $\text{supp } y \cap \text{supp } x = \emptyset$  and a sequence  $t_n \searrow 0$  such that*

$$(4.1) \quad \|x + t_n y\| \geq \|x\| + c\omega(t_n)$$

for some constant  $c > 0$  and any  $n \in \mathbb{N}$ .

*Proof.* We note first that

$$\liminf_{t \rightarrow 0} \frac{\omega(t)}{t} = 0.$$

If  $x = 0$ , choose sequence  $t_n \searrow 0$  such that  $\lim_{n \rightarrow \infty} \omega(t_n)/t_n = 0$ . Then (4.1) holds trivially for any  $y \neq 0$  with  $c = \|y\| > 0$ .

WLOG suppose that  $M(1) = 1$ .

Fix an arbitrary  $x = \sum_{n=1}^{\infty} x_n e_{\gamma_n} \in d(w, M, \Gamma)$  and  $\|x\| = 1$ . Just for simplicity of notation we will assume that  $|x_1| \geq |x_2| \geq \dots \geq |x_n| \geq \dots$ .

We will choose sequences  $t_n \searrow 0$  and  $v_n \searrow 0$  inductively:

1.  $t_1 = v_1 = u_1 = 1, k_0 = 0, k_1 = 1$ .



Title Page

Contents



Page 12 of 18

Go Back

Full Screen

Close

2. Find  $k_2 > k_1, k_2 \in \mathbb{N}$  such that

$$\frac{1}{2^1 \sum_{j=k_1+1}^{k_2} w_j} < M(v_1) \quad \text{and} \quad M(x_i) < \frac{M(t_1 v_1)}{2}$$

for  $i \geq k_2$ .

Find  $t_2 < t_1, v_2 < v_1$  such that

$$\frac{M(t_2 v_2)}{\omega(t_2) M(v_2)} > 2^2 \quad \text{and} \quad M(v_2) < \frac{1}{2^2 \sum_{j=k_1+1}^{k_2} w_j}.$$

3. Find  $k_3 > k_2, k_3 \in \mathbb{N}$  such that

$$\frac{1}{2^2 \sum_{j=k_2+1}^{k_3} w_j} < M(v_2) \quad \text{and} \quad M(x_i) < \frac{M(t_2 v_2)}{2}$$

for  $i \geq k_3$ .

Find  $t_3 < t_2, v_3 < v_2$  such that

$$\frac{M(t_3 v_3)}{\omega(t_3) M(v_3)} > 2^3 \quad \text{and} \quad M(v_3) < \frac{1}{2^3 \sum_{j=k_2+1}^{k_3} w_j}.$$

If we have chosen  $t_{n-1}, v_{n-1}$  and  $k_{n-1}$  then

4. Find  $k_n > k_{n-1}, k_n \in \mathbb{N}$  such that

$$\frac{1}{2^{n-1} \sum_{j=k_{n-1}+1}^{k_n} w_j} < M(v_{n-1}) \quad \text{and} \quad M(x_i) < \frac{M(t_{n-1} v_{n-1})}{2}$$

for  $i \geq k_n$ .



Title Page

Contents



Page 13 of 18

Go Back

Full Screen

Close

Find  $t_n < t_{n-1}, v_n < v_{n-1}$  such that

$$\frac{M(t_n v_n)}{\omega(t_n) M(v_n)} > 2^n \quad \text{and} \quad M(v_n) < \frac{1}{2^n \sum_{j=k_{n-1}+1}^{k_n} w_j}.$$

For a sequence  $\{A_n\}_{n=1}^{\infty}$  of finite disjoint subsets of  $\Gamma$ , such that  $A_n \cap \text{supp } x = \emptyset$ ,  $\#A_n = k_n - k_{n-1}$ , put

$$y_n = v_n \sum_{\gamma \in A_n} e_{\gamma} \quad \text{and} \quad y = \sum_{n=1}^{\infty} y_n.$$

Obviously

$$\begin{aligned} I(y) &= \sum_{n=1}^{\infty} \sum_{j=k_{n-1}+1}^{k_n} w_j M(v_n) \\ &= w_1 M(v_1) + \sum_{n=2}^{\infty} M(v_n) \sum_{j=k_{n-1}+1}^{k_n} w_j \\ &\leq 1 + \sum_{n=2}^{\infty} \frac{1}{2^n} < \infty, \end{aligned}$$

which ensures  $y \in d(w, M, \Gamma)$ . We have  $\text{supp } (x + t_n y) = \text{supp } x \cup (\bigcup_{n=1}^{\infty} A_n)$  for any  $t \neq 0$  and therefore

$$\begin{aligned} (4.2) \quad I(x + t_n y) - I(x) &\geq I(x + t_n y_n) - I(x) \\ &\geq \sum_{j=1}^{k_{n+1}} w_j M(x_j) + \sum_{j=k_{n+1}+1}^{k_{n+2}} w_j M(t_n v_n) \end{aligned}$$



[Title Page](#)

[Contents](#)



Page 14 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

$$\begin{aligned}
 & + \sum_{j=k_{n+2}+1}^{\infty} w_j M(x_{j+k_{n+1}-k_{n+2}}) - \sum_{j=1}^{\infty} w_j M(x_j) \\
 = & M(t_n v_n) \sum_{j=k_{n+1}+1}^{k_{n+2}} w_j - \sum_{j=k_{n+1}+1}^{k_{n+2}} w_j M(x_j) \\
 & + \sum_{j=k_{n+2}+1}^{\infty} w_j (M(x_{j+k_{n+1}-k_{n+2}}) - M(x_j)) \\
 \geq & \frac{1}{2} M(t_n v_n) \sum_{j=k_{n+1}+1}^{k_{n+2}} w_j \\
 \geq & \frac{1}{2} 2^n \omega(t_n) M(v_n) \sum_{j=k_{n+1}+1}^{k_{n+2}} w_j \\
 \geq & 2^{n-1} \omega(t_n) \frac{M(v_n)}{2^{n+1} M(v_{n+1})} \geq \frac{\omega(t_n)}{4}.
 \end{aligned}$$

Remove as many elements of the sequence  $\{t_n\}_{n=1}^{\infty}$  as necessary to obtain

$$0 < d_n = \|x + t_n y\| - 1 \leq 1$$

and keep the same notation for the remaining sequence. Now (2.1) implies

$$\begin{aligned}
 (4.3) \quad I(x + t_n y) - I(x) & = I\left(\|x + t_n y\| \frac{x + t_n y}{\|x + t_n y\|}\right) - 1 \\
 & \leq \|x + t_n y\|^q - 1 \\
 & = (1 + d_n)^q - 1 \leq q 2^{q-1} d_n,
 \end{aligned}$$

for some  $q > \beta_M$ .

Combining (4.2) and (4.3), we obtain

$$\|x + t_n y\| - 1 \geq c\omega(t_n),$$

where  $c = \frac{1}{q^{2q+1}}$ .

Now let  $x \neq 0$  be arbitrary. Find  $\bar{y}$  such that  $\text{supp } \bar{y} \cap \text{supp } x = \emptyset$  and  $\left\| \frac{x}{\|x\|} - t_n \bar{y} \right\| - 1 \geq c\omega(t_n)$ . Obviously for  $y = \|x\|\bar{y}$  we have

$$\|x + t_n y\| - \|x\| \geq c\|x\|\omega(t_n).$$

□



**Gâteaux Differentiability**

B. Zlatanov

vol. 8, iss. 4, art. 113, 2007

Title Page

Contents



Page 15 of 18

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 16 of 18

Go Back

Full Screen

Close

## 5. Gâteaux Differentiability of Bumps in $d(w, M, \Gamma)$ and $d(w, p, \Gamma)$

**Theorem 5.1.** *Let  $M$  be an Orlicz function and  $\omega : (0, 1] \rightarrow \mathbb{R}^+$ ,  $d_M(\omega) = \infty$ .*

- (i) *If  $\alpha_M \notin \mathbb{N}$  then there is no continuous  $G_{\omega, [\alpha_M]}^0$ -smooth bump function in  $d(w, M, \Gamma)$ ;*
- (ii) *If  $\alpha_M \in \mathbb{N}$  then there is no continuous  $G_{\omega, \alpha_M}^0$ -smooth bump function, provided  $d_M(t^{\alpha_M}) < \infty$  in  $d(w, M, \Gamma)$  and there is no continuous  $G_{\omega, \alpha_M - 1}^0$ -smooth bump function, provided  $d_M(t^{\alpha_M}) = \infty$  in  $d(w, M, \Gamma)$ .*

*Proof.* The proof in all cases is straightforward, applying Lemma 4.1 for appropriate  $p$ , Lemma 4.2 and Theorem 2.1. □

*Proof of Theorem 3.1.* The proof in the two cases is straightforward, applying Theorem 5.1. □

It is well known that in a Banach space  $X$  a norm of some order of smoothness generates a bump function with the same order of smoothness (see e.g. [2]), therefore the next corollary is a direct consequence of Theorem 3.1

**Corollary 5.2.** *Let  $M$  be an Orlicz function. If  $|\cdot|$  is an equivalent norm in  $d(w, M, \Gamma)$ , which is  $k$ -times Gâteaux differentiable then  $k \leq E_M$ .*

As a consequence of Theorem 5.1 and Theorem 3.1 we get for  $M(t) = t^p$ ,  $p \geq 1$  the results from [9].

**Corollary 5.3 ([9, Theorem 3]).** *Let  $p \geq 1$ ,  $w_n \searrow 0$ ,  $\sum_{n=1}^{\infty} w_n = \infty$  and  $\omega : (0, 1] \rightarrow \mathbb{R}^+$  be such that  $\omega(t) = o(t^p)$ . Then there is no continuous  $G_{\omega, [p]}^0$ -smooth bump function in  $d(w, p, \Gamma)$ .*



*Proof.* Indeed in this case  $\alpha_M = p$  and  $d_{t^p}(\omega) = \infty$ . If  $p \notin \mathbb{N}$  then by Theorem 5.1 i), it follows that there is no continuous  $G_{\omega, [p]}^0$ -smooth bump in  $d(w, M, \Gamma)$ . If  $p \in \mathbb{N}$  then  $d_{t^p}(t^p) = 1 < \infty$  and by Theorem 5.1 ii), there is no continuous  $G_{\omega, p}^0$ -smooth bump in  $d(w, M, \Gamma)$ .  $\square$

**Corollary 5.4 ([9, Corollary 2]).** *Let  $p \geq 1$ ,  $w_n \searrow 0$ ,  $\sum_{n=1}^{\infty} w_n = \infty$ . If  $f$  is a continuous  $k$ -times Gâteaux differentiable bump function in  $d(w, p, \Gamma)$ , then  $k \leq [p]$ .*

*Proof.* In this case it is obvious that  $d_{t^p}(t^{[p]}) < \infty$  and  $d_{t^p}(t^p) < \infty$ . Therefore by Theorem 3.1 it follows that  $k \leq [p]$ .  $\square$

Title Page

Contents



Page 17 of 18

Go Back

Full Screen

Close



## References

- [1] N. BONIC AND J. FRAMPTON, Smooth functions on Banach manifolds, *J. Math. Mech.*, **15** (1966), 877–898.
- [2] R. DEVILLE, G. GODEFROY AND V. ZIZLER, *Smoothness and Renormings in Banach Spaces*, Pitman Monographs and Surveys in Pure and Applied Mathematics, Vol 64, Logman Scientific and Technical, Harlow/New York, 1993.
- [3] F. HERNANDEZ AND S. TROYANSKI, On representation of uncountable symmetric basic sets and its applications, *Studia Math.*, **107** (1993), 287–304.
- [4] F. HERNANDEZ AND S. TROYANSKI, On Gâteaux differentiable bump functions, *Studia Math.*, **118** (1996), 135–143.
- [5] A. KAMINSKA AND Y. RAYNAUD, Isomorphic  $\ell_p$ -subspaces in Orlicz–Lorentz sequence spaces, *Proc. Amer. Math. Soc.*, **134** (2006), 2317–2327.
- [6] J. KURZWEIL, On approximation in real Banach space, *Studia Math.*, **14** (1954), 213–231.
- [7] J. LINDENSTRAUSS AND L. TZAFRIRI, *Classical Banach Spaces I, Sequence Spaces*, Springer–Verlag, Berlin, 1977.
- [8] R. MALEEV, Higher order uniformly Gateaux differentiable norms in Orlicz spaces, *Rocky Mountain J. Math.*, **28** (1995), 1117–1136.
- [9] R. MALEEV, G. NEDEV AND B. ZLATANOV, Gâteaux differentiability of bump functions in Banach spaces, *J. Math. Anal. Appl.*, **240** (1999), 311–323.
- [10] S. TROYANSKI, Gateaux differentiable norms in  $L_p$ , *Math. Ann.*, **287** (1990), 221–227.

Gâteaux Differentiability

B. Zlatanov

vol. 8, iss. 4, art. 113, 2007

Title Page

Contents



Page 18 of 18

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-575b