



Extension of a Theorem of Duffin and Schaeffer

Michael Coons

School of Mathematical and Physical Sciences
The University of Newcastle
Callaghan, NSW 2308
Australia

Michael.Coons@newcastle.edu.au

Abstract

Let $r_1, \dots, r_s : \mathbb{Z}_{n \geq 0} \rightarrow \mathbb{C}$ be linearly recurrent sequences whose associated eigenvalues have arguments in $\pi\mathbb{Q}$ and let $F(z) := \sum_{n \geq 0} f(n)z^n$, where $f(n) \in \{r_1(n), \dots, r_s(n)\}$ for each $n \geq 0$. We prove that if $F(z)$ is bounded in a sector of its disk of convergence, then it is a rational function. This extends a very recent result of Tang and Wang, who gave the analogous result when the sequence $f(n)$ takes on values of finitely many polynomials.

1 Introduction

In 1945, Duffin and Schaeffer [2] proved that *if a power series whose coefficients are taken from a finite set is bounded in a sector of the unit disk, then it is already a rational function.* In this short note, we prove the following extension.

Theorem 1. *Let $r_1, \dots, r_s : \mathbb{Z}_{n \geq 0} \rightarrow \mathbb{C}$ be linearly recurrent sequences whose associated eigenvalues have arguments in $\pi\mathbb{Q}$ and let $F(z) := \sum_{n \geq 0} f(n)z^n$, where $f(n) \in \{r_1(n), \dots, r_s(n)\}$ for each $n \geq 0$. If $F(z)$ is bounded in a sector of its disk of convergence, then it is a rational function.*

Our result, follows from a very recent result of Tang and Wang [3, Theorem 1.3], who used a method developed by Borwein, Erdélyi, and Littmann [1] to show that *if $p_1(z), \dots, p_s(z) \in \mathbb{C}[z]$ are polynomials and $F(z) := \sum_{n \geq 0} f(n)z^n$, where $f(n) \in \{p_1(n), \dots, p_s(n)\}$ for each $n \geq 0$, is bounded in a sector of the unit disk, then it is a rational function.*

2 Proof of Theorem 1

Proof of Theorem 1. Let $r_1, \dots, r_s : \mathbb{Z}_{n \geq 0} \rightarrow \mathbb{C}$ be linearly recurrent sequences whose associated eigenvalues have arguments in $\pi\mathbb{Q}$ and $f(n) \in \{r_1(n), \dots, r_s(n)\}$ for each $n \geq 0$. Then for each $i = 1, \dots, s$, there exist an integer ℓ , $\ell \times 1$ vectors \mathbf{u}_i and \mathbf{v}_i with complex entries, and an $\ell \times \ell$ matrix \mathbf{A} with complex entries such that $r_i(n) = \mathbf{u}_i^T \mathbf{A}_i^n \mathbf{v}_i$. Recall that $r_i(n)$ can be written in the eigenvalue expansion

$$r_i(n) = \sum_{j=0}^k \alpha_j n^{m_j-1} \lambda_j^n,$$

where $\lambda_1, \dots, \lambda_k$ are the eigenvalues of \mathbf{A}_i , which we have assumed have arguments in $\pi\mathbb{Q}$, m_j is the multiplicity of λ_j , and $\alpha_1, \dots, \alpha_k \in \mathbb{C}$.

Similarly, we can consider a “common” eigenvalue expansion of the linear recurrences r_1, \dots, r_s . In particular, let

$$\rho_1 < \rho_2 < \dots < \rho_D \tag{1}$$

be the distinct moduli of the collection of eigenvalues of $\mathbf{A}_1, \dots, \mathbf{A}_s$. Then there exist integers D and d and a finite collection of complex numbers $\{\alpha_{n,i,m,t}\}_{n \geq 0, 1 \leq i \leq s, 1 \leq m \leq D, 0 \leq t \leq d}$, many of which may be zero, such that

$$r_i(n) = \sum_{m=1}^D \sum_{t=0}^d \alpha_{n,i,m,t} n^t \rho_m^n \quad (i = 1, 2, \dots, s, \text{ and } n \geq 0). \tag{2}$$

We note that the collection $\{\alpha_{n,i,m,t}\}_{n \geq 0, 1 \leq i \leq s, 1 \leq m \leq D, 0 \leq t \leq d}$ is finite precisely because all of the eigenvalues associated to the linear recurrences have arguments that are rational multiples of π (this assumption in the statement of the theorem is crucial for this proof to work).

Now consider $F(z) := \sum_{n \geq 0} f(n)z^n$ satisfying the assumptions of the theorem. Suppose that $F(z)$ is bounded in a sector of its disk of convergence, that $F(z)$ is irrational, and that D is the minimal such number so that the previous two properties hold. We will prove the theorem by showing that such a minimal D cannot exist.

To this end, set $\Psi(z) := F(z/\rho_D)$, so that the coefficients $\psi(n)$ of $\Psi(z)$ satisfy $\psi(n) = f(n)\rho_D^{-n}$, $\Psi(z)$ is bounded in a sector of the unit disk, and $\Psi(z)$ is irrational. Using (2) we have that

$$\psi(n) = \alpha_{n,i_n,D,d} \cdot n^d + \alpha_{n,i_n,D,d-1} \cdot n^{d-1} + \dots + \alpha_{n,i_n,D,0} + r(n), \tag{3}$$

where for each n ,

$$r(n) \in \{r'_1(n), \dots, r'_s(n)\} \quad \text{with} \quad r'_i(n) = \sum_{m=1}^{D-1} \sum_{t=0}^d \alpha_{n,i,m,t} n^t (\rho_m/\rho_D)^n, \tag{4}$$

and where we have used the notation i_n to indicate the value of $i \in \{1, \dots, s\}$ for which $f(n) = r_i(n)$. By (1), (4), and the minimality of D , we have that the function $R(z) =$

$\sum_{n \geq 0} r(n)z^n$ is bounded throughout the unit disk (as it has radius of convergence at least $\rho_D/\rho_{D-1} > 1$) and is rational. Thus by (3), the function

$$\Psi(z) - R(z) = \sum_{n \geq 0} (\alpha_{n,i_n,D,d} \cdot n^d + \alpha_{n,i_n,D,d-1} \cdot n^{d-1} + \cdots + \alpha_{n,i_n,D,0}) z^n$$

is bounded in a sector of the unit disk and irrational. But by Tang and Wang's result, the function $\Psi(z) - R(z)$ is rational. This contradiction proves the theorem. \square

3 A question

The careful reader will notice that while Tang and Wang's result holds for polynomials in $\mathbb{C}[z]$, the linear recurrences in our theorem, while also being complex-valued, must have associated eigenvalues with arguments in $\pi\mathbb{Q}$. This is because if we let the eigenvalues be more general complex numbers, the arguments of the powers of those eigenvalues may not take only finitely many values. Our proof works as long as this finiteness condition holds.

Because of the necessity of this finiteness condition, Theorem 1 is essentially the best possible result that can be proved from the methods of Borwein, Erdélyi, and Littmann [1], and Tang and Wang [3]. An interesting generalization would be a statement as in Theorem 1 for general complex linear recurrences. To apply the same basic theory, one would need to answer the following question in the affirmative.

Question 2. Let $a : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{C}$ be given by

$$a(n) = \sum_{i=0}^M b_{i,n} \zeta_i^n,$$

where $b_{i,n}, \zeta_i \in B$, B is finite set of complex numbers, and ζ_i lies on the unit circle for each i . Suppose that for each $\varepsilon > 0$ there exist a positive integer q and complex numbers $\gamma_0, \dots, \gamma_{q-1}$ such that for all $n \geq 0$,

$$|\gamma_0 a(n) + \cdots + \gamma_{q-1} a(n+q-1) + a(n+q)| < \varepsilon.$$

Is it true that the sequence a is linearly recurrent?

4 Acknowledgment

We thank the referee and editor—their great care made this a better (and correct!) paper. This paper was written while visiting the Alfréd Rényi Institute of the Hungarian Academy of Sciences; we thank the Institute and its members for their kindness and support.

References

- [1] P. Borwein, T. Erdélyi, and F. Littmann, Polynomials with coefficients from a finite set, *Trans. Amer. Math. Soc.* **360** (2008), 5145–5154.
- [2] R. J. Duffin and A. C. Schaeffer, Power series with bounded coefficients, *Amer. J. Math.* **67** (1945), 141–154.
- [3] S. Tang and C. Wang, Rational-transcendental dichotomy of power series with a restriction on coefficients, *Monatsh. Math.* (2017), published online 24 May 2017.

2010 *Mathematics Subject Classification*: Primary 30B30 Secondary 11B37, 11J91.

Keywords: Power series, rational function, Duffin-Schaeffer theorem.

Received June 8 2017; revised versions received August 9 2017; September 5 2017; September 11 2017. Published in *Journal of Integer Sequences*, September 15 2017.

Return to [Journal of Integer Sequences home page](#).