

SEMIALGEBRAIC SARD THEOREM FOR GENERALIZED CRITICAL VALUES

K. KURDYKA, P. ORRO & S. SIMON

Abstract

We prove that a semialgebraic differentiable mapping has a generalized critical values set of measure zero. Moreover, if the mapping is C^2 we obtain, by a generalisation of Ehresmann's fibration theorem due to P. J. Rabier [20], a locally trivial fibration over the complement of this set. In the complex case, we prove that the set of generalized critical values of a polynomial mapping is a proper algebraic set.

1. Introduction

The usual Sard's theorem says that the set $K_0(f)$ of critical values of a C^p map $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ has zero Lebesgue measure when $p \geq \max(1, n - k + 1)$. The Ehresmann's fibration theorem asserts that a proper submersion is a locally trivial fibration. Thus $K_0(f)$ is a bifurcation set of a proper map and is a small set.

The fibration theorem has been generalized in different ways:

- For general non-proper functions, R. S. Palais introduced a condition, known as the (C) condition of Palais. Roughly speaking it means that the norm of the differential of f is separated from zero uniformly fibrewise. Palais proved that a function from a Hilbertian manifold M to \mathbb{R} satisfying condition (C) is still a fibration outside of $K_0(f)$. Later he generalized this result in the case M is a complete Finsler manifold (cf. [15]).
- For general non-proper mappings $f : M \rightarrow N$, where M and N are Finsler manifolds, P. J. Rabier [20] introduced the notion of strong

Received November 22, 1999.

submersion which generalizes condition (C) of Palais. The norm of the differential is replaced by $\nu(df)$ (as we explain in Section 2, the function ν of Rabier is simply the distance to the set of singular operators). Under the hypothesis of completeness of M and some technical assumption (which always holds for Hilbert manifolds) Rabier proves that every strong submersion is a fibration.

- For polynomials $f : \mathbb{C}^n \rightarrow \mathbb{C}$, it is well known that f is a fibration outside the bifurcation set $B(f)$, which is a finite set consisting of points in $K_0(f)$ and critical points at infinity. Many authors tried to make precise the bifurcation set: [16], [19], [20], [22] — introducing different conditions at infinity such as quasi-tame, Malgrange condition, M-tame,

In this paper we are interested in the case of semialgebraic mappings from \mathbb{R}^n to \mathbb{R}^k (or polynomial from \mathbb{C}^n to \mathbb{C}^k). We prove that the set of generalized critical values $K(f) = K_0(f) \cup K_\infty(f)$, where

$$K_\infty(f) := \{y \in \mathbb{K}^k : \exists x_l \in \mathbb{K}^n, |x_l| \rightarrow \infty \\ \text{s.t. } f(x_l) \rightarrow y \text{ and } (1 + |x_l|)\nu(df(x_l)) \rightarrow 0\},$$

is a zero-measure semialgebraic subset of \mathbb{K}^k (constructible if $\mathbb{K} = \mathbb{C}$).

Using Rabier's results [20], this gives a fibration theorem for f over the connected components of $\mathbb{K}^k - K(f)$. The main point for the fibration result is the completeness of \mathbb{K}^n . If we consider a map f defined on an open set of \mathbb{K}^n , a similar result is valid if f is without fibers adherent to ∂U . Another way of proving a fibration result for maps defined on open sets is to take a complete metric on U and define a new K_∞ -set relative to this metric.

The set $K_\infty(f)$ is called the set of asymptotic critical values (we use the notation of [20] but we take the definition given in Remark 6.10 [20], see our Section 2). The fact that $K_\infty(f)$ is finite, for semialgebraic functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$, plays a crucial role in a proof of the Gradient Conjecture of R. Thom (cf. [12], [13]). A different proof of the finiteness of $K_\infty(f)$ is given by D. d'Acunto ([1]) in a more general setting of o-minimal structures.

The proof of the main result, i.e., that for a semialgebraic mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$, the set $K_\infty(f)$ has zero measure (more precisely that $\dim K_\infty(f) < k$) is inspired by Y. Yomdin's quantitative Sard theorem for polynomials [23], see also S. K. Donaldson [4, Section 5]. We decompose the set on which $|x|\nu(df(x))$ is small in a parametric family of

L -regular cells (a L -regular cell is a subset of \mathbb{R}^n on which the geodesic distance is equivalent to the euclidian distance); and then, by giving an estimate for length of the image by a component of f of each cell (Lemma 3.6), we deduce the result. We give also an alternative proof of Lemma 3.6 based on a result of B. Teissier [21].

The theory of generalized critical values appears as an alternative to the stratification theory. Actually our main result (in the real case) can be also proved using the (w) condition of Kuo and Verdier. However our approach via L -regular sets is conceptually simpler, and moreover gives an explicit bound for the number of points in $K_\infty(f)$ in term of the degree of f for $k = 1$ (cf. Remark 3.2).

For a polynomial mapping $f : \mathbb{K}^n \rightarrow \mathbb{K}^k$ let $J_f \subset \mathbb{K}^k$ denote the set of points where f is not proper, we call it *the Jelonek set of f* . It was proved by Z. Jelonek that J_f is \mathbb{K} -uniruled (i.e., by each point passes a curve with a polynomial parametrization), if $n = k$ and f is dominant. Moreover in the complex case J_f is a hypersurface or an empty set. In Proposition 3.1 we prove that $K_\infty(f) = J_f$, if f is a C^1 semialgebraic mapping from \mathbb{R}^n to \mathbb{R}^n . We also prove (Theorem 3.4) that for polynomial dominant mappings from \mathbb{R}^n to \mathbb{R}^k , with compact regular fibers, the equality $K_\infty(f) = J_f$ holds as well. Moreover, by a result of Z. Jelonek [8], the set J_f is also \mathbb{R} -uniruled in this case. If the generic fibers are not compact, then of course $K_\infty(f) \neq J_f$. We conjecture that for a polynomial mapping $f : \mathbb{K}^n \rightarrow \mathbb{K}^k$ the set $K_\infty(f)$ is \mathbb{K} -uniruled.

In Section 2 we recall the definition of the ν function of Rabier and we give several equivalent definitions of ν . In particular we prove that this function is equal to the distance to the set of singular operators, at our knowledge this fact was only known in the finite dimensional case when the space of linear operators is endowed with the L^2 -norm. We decided to write down this part in the setting of Banach spaces since the theorem of Rabier has potentially a lot of applications in this case. Finally we recall another way for measuring the distance to the set of singular operators (valid for finite dimensional target), the so called Kuo distance which was introduced by T. C. Kuo in [14]. It turns out to be equivalent to ν and is more convenient in our construction. We discuss also two possible definitions of the set of generalized critical values and we recall some examples which show that these definitions are not equivalent.

The proofs of our main results are given in Section 3 in the real case, Theorem 3.1, and in Section 4, Theorem 4.1, in the complex case.

We are happy to thank P. J. Rabier and Z. Jelonek for valuable remarks on our paper.

2. Preliminaries and notations

2.1 Distance to the set of singular mappings

Let X, Y be Banach spaces (over $\mathbb{K} = \mathbb{R}$ or \mathbb{C}). We will denote by $\mathcal{L}(X, Y)$ the Banach space of linear continuous mappings from X to Y , and by Σ the singular set of $\mathcal{L}(X, Y)$ that is the subset of operators which are not onto. The distance function to Σ is defined as usual by

$$\text{dist}(A, \Sigma) = \inf_{B \in \Sigma} \|A - B\|.$$

If $Y = \mathbb{K}$, it is customary to write X' for $\mathcal{L}(X, \mathbb{K})$. For $A \in \mathcal{L}(X, Y)$, A^* stands for the algebraic adjoint operator in $\mathcal{L}(Y', X')$.

Definition 2.1 ([20]). Let $A \in \mathcal{L}(X, Y)$. Set

$$\nu(A) = \inf_{\|\varphi\|=1} \|A^*\varphi\|.$$

Notice that, since X, Y are Banach spaces, $\nu(A) > 0 \Leftrightarrow A$ is onto. The properties in the next proposition are well known and easy to establish, see for instance [20]

Proposition 2.1. *Let $A \in \mathcal{L}(X, Y)$.*

1. *If $Y = \mathbb{K}$ then $\nu(A) = \|A\|$.*
2. *If $A \in GL(X, Y)$ then $\nu(A) = \|A^{-1}\|^{-1}$.*
3. *ν is 1-Lipschitz.*

The two functions ν and $\text{dist}(\cdot, \Sigma)$ vanish on Σ . We prove that they are in fact the same. We begin by a lemma.

Lemma 2.1. *Let $A \in \mathcal{L}(X, Y)$ be a surjective map and $\epsilon > 0$. Then*

$$\text{dist}(A, \Sigma) \leq \nu(A) + \epsilon.$$

Proof. By definition of $\nu(A)$, there exists some $\varphi_\epsilon \in Y'$, $\|\varphi_\epsilon\| = 1$, such that $\|\varphi_\epsilon \circ A\| \leq \nu(A) + \epsilon$. Now, consider the topological sum $Y = \ker \varphi_\epsilon \oplus \langle y_\epsilon \rangle$, where y_ϵ is any vector satisfying $\varphi_\epsilon(y_\epsilon) = 1$. Then, denote

by p_ϵ the canonical projection onto $\langle y_\epsilon \rangle$ and define $B_\epsilon = A - p_\epsilon \circ A \in \Sigma$. Therefore,

$$\|A - B_\epsilon\| = \|p_\epsilon \circ A\| = \sup_{\|x\|=1} \|p_\epsilon \circ A(x)\| = \|y_\epsilon\| \cdot \|\varphi_\epsilon \circ A\|.$$

So that, $\frac{\text{dist}(A, \Sigma)}{\|y_\epsilon\|} \leq \nu(A) + \epsilon$.

Now, remember that if $\psi \in Y' - \{0\}$ then for all $y \in Y$, $\text{dist}(y, \ker \psi) = \frac{|\psi(y)|}{\|\psi\|}$. It follows that $\text{dist}(0, \ker \varphi_\epsilon + y_\epsilon) = 1$. Consequently, there exists a sequence u_n in $\ker \varphi_\epsilon$ such that $y_\epsilon^n := y_\epsilon + u_n$ satisfies $\varphi_\epsilon(y_\epsilon^n) = 1$ for all n and $\|y_\epsilon^n\| \searrow 1$. At last, apply the preceding construction in substituting y_ϵ by y_ϵ^n and get the result. q.e.d.

Proposition 2.2. *Let $A \in \mathcal{L}(X, Y)$. We have*

$$\nu(A) = \text{dist}(A, \Sigma).$$

Proof. The inequality $\nu(A) \leq \text{dist}(A, \Sigma)$ is a consequence of the fact that ν is 1-Lipschitz, whereas $\nu(A) \geq \text{dist}(A, \Sigma)$ follows from Lemma 2.1. q.e.d.

Remark 2.1. Note that $\nu(A)$ is the maximum radius R such that the open ball of center A and radius R does not intersect Σ .

Another characterization of ν is as follows:

Proposition 2.3. *Let $A \in \mathcal{L}(X, Y)$. Then*

$$\nu(A) = \sup\{r \geq 0 : B(0, r) \subset A(B(0, 1))\}.$$

Proof. Let R denote $\sup\{r \geq 0 : B(0, r) \subset A(B(0, 1))\}$.

We first prove that $\nu(A) \geq R$. Let $r < R$ and $\varphi \in Y'$ such that $\|\varphi\| = 1$. For all $\epsilon > 0$, there exists $y \in S_Y(0, 1)$ with $|\varphi(y)| \geq 1 - \epsilon$. Then, there exists $x \in B_X(0, 1)$ verifying $|\varphi(Ax)| \geq r|\varphi(y)| \geq r(1 - \epsilon)$. Hence $\|\varphi \circ A\| \geq r(1 - \epsilon)$, so $\nu(A) \geq r(1 - \epsilon)$, and finally, $\nu(A) \geq R$.

Now, we prove that $\nu(A) \leq R$. Let $\epsilon > 0$ and $y \in B_Y(0, R + \epsilon) \setminus A(B_X(0, 1))$. Denote $H = A^{-1}(\langle y \rangle)$. Then $A|_H$ defines $\lambda \in H'$ by $A(h) = \lambda(h)y$ for $h \in H$. Applying Hahn-Banach theorem, we extend λ to X and denote this extension by $\tilde{\lambda}$ which is such that $\|\tilde{\lambda}\| = \|\lambda\| < 1$. Now, $B_y = A - \tilde{\lambda}y \in \Sigma$ and $\|A - B_y\| \leq R + \epsilon$. Thus, $\nu(A) \leq R + \epsilon$.

q.e.d.

Assume that X, Y are complex normed vector spaces, then we have on them the induced structures of real normed vector spaces. Let $A : X \rightarrow Y$ be a continuous \mathbb{C} -linear map. We can consider $\nu(A)$ with respect to both structures. It follows immediatly from Proposition 2.3 that $\nu(A)$ is the same in the real and complex case. In particular, by Proposition 2.2 we have

$$(2.1) \quad \nu(A) = \text{dist}(A, \Sigma_{\mathbb{C}}) = \text{dist}(A, \Sigma_{\mathbb{R}}),$$

where $\Sigma_{\mathbb{C}}$ (resp. $\Sigma_{\mathbb{R}}$) is the set of nonsurjective \mathbb{C} -linear (resp. \mathbb{R} -linear) continuous maps from X to Y .

Till the end of Subsection 2.2 we will consider only the finite dimensional case.

Proposition 2.4. *If the norms on \mathbb{R}^n and \mathbb{R}^k are semialgebraic, then ν is semialgebraic on $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^k)$.*

Proof. Since Σ is a semialgebraic set in this case, the result is a consequence of Proposition 2.2 and the classical fact (cf. [2]) that the distance function to a semialgebraic set is semialgebraic (i.e., that its graph is semialgebraic). q.e.d.

Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , we will always consider \mathbb{K}^n equipped with an hermitian scalar product, which will be denoted by $(\cdot | \cdot)$.

The first and well known characterization of $\nu(A)$ is the following:

Proposition 2.5. *Let $A \in L(\mathbb{K}^n, \mathbb{K}^k)$. Then*

$$\nu(A) = \min\{|\mu| : \mu^2 \text{ is an eigenvalue of } AA^*\}.$$

We shall need an expression of ν in terms of gradients of components of a linear mapping. To this end we introduce the Kuo distance κ which is actually equivalent to ν , as we show in Proposition 2.6.

2.2 The Kuo distance

Definition 2.2 ([14]). Let $\eta_1, \dots, \eta_k \in \mathbb{K}^n$ for $k \in \mathbb{N}^*$. The Kuo distance between these vectors is defined by

$$\kappa(\eta_1, \dots, \eta_k) = \min_{1 \leq i \leq k} \text{dist}(\eta_i, \langle (\eta_j)_{j \neq i} \rangle),$$

where $\langle (\eta_j)_{j \neq i} \rangle$ is the vector space generated by the vectors $(\eta_j)_{j \neq i}$.

Proposition 2.6. *Let $A = (A_1, \dots, A_k) \in L(\mathbb{K}^n, \mathbb{K}^k)$. For $i \in \{1, \dots, k\}$, denote by η_i the gradient of A_i , then*

$$\nu(A) \leq \kappa(\eta_1, \dots, \eta_k) \leq \sqrt{k} \nu(A).$$

Proof. For $1 \leq i \leq k$, set $V_i = \{x \in \mathbb{K}^n : (x|\eta_j) = 0, j \neq i\}$. Then $V_i^\perp = \langle (\eta_j)_{j \neq i} \rangle$. So $\text{dist}(\eta_i, V_i^\perp) = |p_{V_i}(\eta_i)| = \sup\{|(x|\eta_i)| : x \in V_i, |x| \leq 1\}$ where $p_{V_i} : \mathbb{K}^n \rightarrow V_i$ is the projection. Denote $\xi_i = f(x_i)$ with $x_i \in V_i \cap S(1)$ ($S(r)$ is the sphere of radius r centered at 0) such that $|\xi_i| = |p_{V_i}(\eta_i)|$. Then $(\xi_i)_{1 \leq i \leq k}$ is an orthogonal family in \mathbb{K}^k . We conclude using the next Lemma 2.2. q.e.d.

Lemma 2.2. *Let (ξ_1, \dots, ξ_k) be an orthogonal basis of \mathbb{K}^k and $u \in S(1)$, then*

i) *there exists $i_0 \in \{1, \dots, k\}$ such that $\frac{1}{\sqrt{k}} \|\xi_{i_0}\| \leq |(u|\xi_{i_0})|$.*

ii) *let (e_1, \dots, e_k) be an orthonormal basis of \mathbb{K}^k , and denote by \mathcal{E} the ellipsoid defined in this basis by $\left\{ x : \sum_{i=1}^k \left| \frac{x_i}{a_i} \right|^2 = 1 \right\}$ where $0 < |a_1| \leq |a_2| \leq \dots \leq |a_k| \in \mathbb{R}_+^*$ for $1 \leq i \leq k$. Suppose that $\xi_i \in \mathcal{E}$ for $1 \leq i \leq k$. Then $\min_{1 \leq i \leq k} \|\xi_i\| \leq \sqrt{k} |a_1|$.*

Proof. Write $\|u\|^2 = 1 = \sum_{i=1}^k \left| \left(u \mid \frac{\xi_i}{\|\xi_i\|} \right) \right|^2$. Then, there necessarily exists $i_0 \in \{1, \dots, k\}$ such that $\left| \left(u \mid \frac{\xi_{i_0}}{\|\xi_{i_0}\|} \right) \right|^2 \geq \frac{1}{k}$, which gives i). Using i), take $i_0 \in \{1, \dots, k\}$ such that $\frac{1}{\sqrt{k}} \|\xi_{i_0}\| \leq |(e_1|\xi_{i_0})|$. Now, $|(e_1|\xi_{i_0})| \leq |a_1|$ because $\xi_{i_0} \in \mathcal{E}$. q.e.d.

It is convenient to use the Kuo distance in the following way: let $A = (A_1, \dots, A_k) \in L(\mathbb{K}^n, \mathbb{K}^k)$ and let η_i be the gradient of A_i . For $i \in \{1, \dots, k\}$ denote $V_i = \bigcap_{j \neq i} \ker A_j$. Observe that

$$\text{dist}(\eta_i, \langle (\eta_j)_{j \neq i} \rangle) = \|A_i|_{V_i}\|,$$

where $A_i|_{V_i}$ is the restriction of A_i to V_i . Recall that $\langle (\eta_j)_{j \neq i} \rangle$ is the vector space generated by the vectors $(\eta_j)_{j \neq i}$, hence it is the orthogonal

complement to V_i . So

$$(2.2) \quad \kappa(\eta_1, \dots, \eta_k) = \min_{1 \leq i \leq k} \|A_i|_{V_i}\|$$

and we denote this number by $\kappa(A)$.

From Proposition 2.6 we obtain immediately:

Corollary 2.1. *Let $A = (A_1, \dots, A_k) \in L(\mathbb{K}^n, \mathbb{K}^k)$, then*

$$\nu(A) \leq \kappa(A) \leq \sqrt{k} \nu(A).$$

2.3 Asymptotic critical values and fibration theorem

Let M, N be $C^{p \geq 2}$ manifolds and U an open subset of M . Suppose that M is complete and N is connected—although most definitions and results of this section are valid in the case of Finsler manifolds we will always suppose that M and N are Hilbert spaces. Suppose given a $C^{p \geq 2}$ -map $f : U \rightarrow N$, satisfying the condition that:

$$(2.3) \quad \text{There is no sequence } x_l \text{ in } U \text{ such that} \\ \lim_{l \rightarrow \infty} x_l \in \partial U \text{ and } \lim_{l \rightarrow \infty} f(x_l) \in N.$$

Then we will denote by $\tilde{K}(f)$ the following set:

$$\tilde{K}(f) = \{y \in N : \exists x_l \in U \ f(x_l) \rightarrow y \text{ and } \nu(df(x_l)) \rightarrow 0\}.$$

The set of critical values of f say

$$K_0(f) = \{y \in N : \exists x \in f^{-1}(y) \text{ s.t. } \nu(df(x)) = 0\}$$

is contained in $\tilde{K}(f)$. In fact $\tilde{K}(f)$ is the union of $K_0(f)$ and $\tilde{K}_\infty(f)$, an asymptotic critical set for f , which is defined by (NCS means no converging subsequences in \bar{U})

$$\tilde{K}_\infty(f) = \{y \in N : \exists x_l \in U \text{ with NCS } f(x_l) \rightarrow y \text{ and } \nu(df(x_l)) \rightarrow 0\}.$$

Clearly $\tilde{K}(f)$ is closed, and $\tilde{K}_\infty(f) = \emptyset$ if f is a proper map. Note that $\tilde{K}_\infty(f)$ is also closed if f is defined on an open subset U in a finite dimensional space.

In [20] one can find the following generalisation of Ehresmann's theorem:

Theorem 2.1 ([20]). *Let $V \subset N$ be a connected component of $N - \tilde{K}(f)$, then either $f^{-1}(V) = \emptyset$ or $f : f^{-1}(V) \rightarrow V$ is a locally trivial C^p -fibration.*

As a particular case this gives the classical Erhesmann's theorem for proper mappings.

In the case of finite dimensional manifolds, the classical Sard's theorem says that $K_0(f)$ is a subset of N of zero measure. In the infinite dimensional setting, if the spaces M and N are separable, one has a similar result for C^p -functions if $p \geq \max(k, 2)$ assuming k is an integer such that $\dim \ker f''(x) \leq k$ for all x ; also for C^r -Fredholm maps with $r > \max(\text{index}(f), 0)$ one can show that the set $K_0(f)$ has empty interior.

Without some extra hypothesis the set $\tilde{K}(f)$, as well as $\tilde{K}_\infty(f)$, may be quite large. Using the fact that $\tilde{K}(f)$ is closed it is not difficult to construct a C^∞ function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $\tilde{K}(f) = \mathbb{R}$. This may happen also for polynomials in more than 2 variables as the example below shows.

Example 2.1 ([17]). Consider the polynomial

$$f(x, y, z) = x + x^2y + x^4yz.$$

Then $K_0(f) = \emptyset$ and $\tilde{K}_\infty(f) = \mathbb{R}$. To see this take any point $a \in \mathbb{R}^*$ and the curve

$$\gamma_a(s) = \left(s, \frac{2a}{s^2}, -\frac{(1 + (4a)^{-1}s)}{2s^2} \right).$$

We have

$$f(\gamma_a(s)) = a + s \left(1 + \frac{1}{4a} \right) \rightarrow a$$

and

$$\nabla f(\gamma_a(s)) = \left(0, s^2 \left(\frac{1}{2} - \frac{s}{8a} \right), 2as^2 \right) \rightarrow 0$$

as $s \rightarrow 0$. Observe nevertheless that f is a fibration over each connected component of \mathbb{R}^* ; see Remark 2.2 and Theorem 3.1 below.

Recall that a value y of the map f is called *typical* if f is a C^∞ fibration near y and *atypical* otherwise. The set $B(f)$ of atypical values, called the *bifurcation set of f* , is contained in $\tilde{K}(f)$ but in general is not equal to it—see Example 2.1 and also the following:

Example 2.2 ([22]). The polynomial $2x^2y^3 - 9xy^2 + 12y$ has no critical points i.e., $K_0(f)$ is empty, and $0 \in \tilde{K}_\infty(f)$. But f is a fibration over \mathbb{R} .

Our main goal is to prove that the set of generalized critical values of a semialgebraic mapping $f : \mathbb{K}^n \rightarrow \mathbb{K}^k$ is nowhere dense in \mathbb{K}^k . So we have to modify the definition of the set $\tilde{K}_\infty(f)$. As noticed in [20, Remark 6.1], Theorem 2.1 remains true if, in the definition of $\tilde{K}_\infty(f)$, we replace the condition $\nu(df(x_l)) \rightarrow 0$ by the following one: $\omega(|x_l|)\nu(df(x_l)) \rightarrow 0$, where $\omega : [0, \infty) \rightarrow [0, \infty)$ is a continuous function with $\int_0^\infty \frac{du}{\omega(u)} = \infty$. We take $\omega(t) = 1 + t$. Let $f : \mathbb{K}^n \rightarrow \mathbb{K}^k$ be C^1 mapping, we define *the set of generalized critical values of f* as

$$K(f) = \{y \in \mathbb{K}^k : \exists x_l \in \mathbb{K}^n \\ \text{s.t. } f(x_l) \rightarrow y \text{ and } (1 + |x_l|)\nu(df(x_l)) \rightarrow 0\}.$$

Clearly $K(f) = K_0(f) \cup K_\infty(f)$, where the set

$$K_\infty(f) = \{y \in \mathbb{K}^k : \exists x_l \in \mathbb{K}^n, |x_l| \rightarrow \infty \\ \text{s.t. } f(x_l) \rightarrow y \text{ and } (1 + |x_l|)\nu(df(x_l)) \rightarrow 0\}$$

will be called *the set of asymptotic critical values of f* .

Note that for polynomial functions $f : \mathbb{C}^n \rightarrow \mathbb{C}$ the condition $t_0 \notin K_\infty(f)$ means that t_0 satisfies the Malgrange condition. Let us recall a characterisation of this condition due to Parusiński.

Theorem 2.2 ([16],[17]). *Let $f : \mathbb{C}^n \rightarrow \mathbb{C}$ be a polynomial with isolated singularities at infinity and t_0 a regular value (i.e., $t_0 \notin K_0(f)$). Then the following are equivalent:*

- (i) $t_0 \notin B(f)$.
- (ii) *The Malgrange condition for t_0 is satisfied, i.e., $\exists \delta > 0 : |x||\nabla f(x)| \geq \delta$ for x large and $f(x)$ close to t_0 .*
- (iii) $\chi(f^{-1}(t))$ is constant near t_0 .

In particular, this theorem characterizes the bifurcation set for all polynomials if $n = 2$, in this case we have $K_\infty(f) = \tilde{K}_\infty(f)$ by [5].

In the remainder of this section we discuss some examples and related notions.

Remark 2.2. For the polynomial of Example 2.1 we have $\tilde{K}_\infty(f) = \mathbb{R}$ but $K_\infty(f) = \{0\}$.

To see this consider

$$\nabla f(x, y, z) = (1 + 2xy + 4x^3yz, x^2 + x^4z, x^4y) =: (A, B, C).$$

We already know that $0 \in K_\infty(f)$.

Take any $t \neq 0$. It is easy to see that $x \rightarrow 0$ if $\nabla f \rightarrow 0$: Since B and C must be bounded, A cannot go to 0 if x is unbounded; and y and z must be bounded if x goes to a limit not equal to 0. Now, since $x(1 + xy + x^3yz) \rightarrow t$, if y is bounded then $A = 1 + xy + x^3yz + xy + x^3yz + 2x^3yz \rightarrow \infty$ thus $y \rightarrow \infty$. Put $m = (x, y, z)$ and suppose $|m|\nabla f(m) \rightarrow 0$. The component $|m|C \rightarrow 0$ so $|x^2y| = \epsilon(m) \rightarrow 0$, and since $x + x^2y + x^4yz \rightarrow t$ we also have $x^4yz = t + \delta(m)$ with $\delta(m) \rightarrow 0$. To finish, consider the component $|m|A$; we have $|m|A = |m| + \frac{|m|}{|x|}(4|t| + 4\delta(m) + \epsilon(m)) \geq (5|t| + 1)|m|$ and so $|m|A$ cannot go to 0.

Remark 2.3. For the polynomial of Example 2.2 we have $K(f) = \{0\}$.

Let us recall some classes of polynomials “without critical points at infinity”:

Denote by $M(f) = \{x \in \mathbb{C}^n : \exists \lambda \in \mathbb{C} : \nabla f(x) = \lambda x\}$. A polynomial f is M -tame if for any sequence $x_k \in M(f)$ such that $x_k \rightarrow \infty$ we have $\lim_{k \rightarrow \infty} f(x_k) = \infty$

A polynomial f is called quasi-tame if for any sequence $x_k \in \mathbb{C}^n$ such that $x_k \rightarrow \infty$ and $\nabla f(x_k) \rightarrow 0$ we have $\lim_{k \rightarrow \infty} |f(x_k) - (x_k|\nabla f(x_k))| = \infty$.

It is well known, and easy to prove, that quasi-tame implies Malgrange condition and that Malgrange condition at any point (which is the same as $K(f) = \emptyset$) implies M -tame. Finally we will give an example showing that $K(f)$ is in general different from $B(f)$

Example 2.3 ([19]). The polynomial $x + y - 2x^2y^3 + x^3y^6 + zy^3 - z^2y^5 + ty^5$ is a fibration over \mathbb{C} that is $B(f) = \emptyset$. But it is not M -tame at 0, and thus $K_\infty(f) \neq \emptyset$ —in fact $K_\infty(f) = \{0\}$.

3. Main result. Real case

From now on we will restrict attention to the finite dimensional case. Our aim is to prove the following theorem:

Theorem 3.1. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ be a differentiable semialgebraic map. Then $K(f)$ is a closed semialgebraic set of dimension less than k . Moreover, if f is of class C^2 , then $f : \mathbb{R}^n \setminus f^{-1}(K(f)) \rightarrow \mathbb{R}^k \setminus K(f)$ is a fibration over each connected component of $\mathbb{R}^k \setminus K(f)$. In particular $B(f) \subset K(f)$.*

Here $B(f)$ means the smallest closed subset of \mathbb{R}^k such that $f : \mathbb{R}^n \setminus f^{-1}(B(f)) \rightarrow \mathbb{R}^k \setminus B(f)$ is a fibration over each connected component of $\mathbb{R}^k \setminus B(f)$.

Remark 3.1. The theorem remains valid in the case of a mapping defined on an open subset U of \mathbb{R}^n . That is $K(f)$ is still a closed semialgebraic set of dimension less than k and, if f satisfies Condition (2.3), $f : U - f^{-1}(K(f)) \rightarrow \mathbb{R}^k - K(f)$ is a fibration over each connected component of $\mathbb{R}^k - K(f)$.

The main argument of our proof is based on a fact, due to K. Kurdyka ([10]), that any semialgebraic set $A \subset \mathbb{R}^n$ is a finite union $A = \cup_i L^i$, where each L^i has the *Whitney property with constant M* : any two points $x, y \in L^i$ can be joined in L^i by a piecewise smooth arc of length $\leq M|x - y|$. (Actually any $M > 1$ will do, by [9]). It is crucial for us that the constant $M = M(n)$ depends only on n . In [18] A. Parusinski obtained a similar decomposition of semialgebraic sets, but without any estimate on M .

What we need actually is a uniform version of the above decomposition, for families parameterized by finite dimensional spaces : if $B \subset \mathbb{R}^n \times \mathbb{R}^p$ and $t \in \mathbb{R}^p$, we write $B_t = \{x \in \mathbb{R}^n : (x, t) \in B\}$. Then from the method of [10] (see also [11], [12, Chap. 2]), we obtain the following theorem:

Theorem 3.2. *There exists $M = M(n) > 0$ such that any semialgebraic set $A \subset \mathbb{R}^n \times \mathbb{R}^p$ can be decomposed into a finite (and disjoint) union $A = \cup_{i \in I} L^i$, such that for each $t \in \mathbb{R}^p$, every set L_t^i has the Whitney property with constant M . So, in particular $A_t = \cup_{i \in I} L_t^i$ for each $t \in \mathbb{R}^p$. (Clearly, for a fixed $t \in \mathbb{R}^p$ some of L_t^i may be empty.)*

In fact for our purpose we may use a weaker result due to B. Teissier [21], we explain it after the proof of the main theorem.

The proof of Theorem 3.1 will be given in Section 3.3. The main steps of the proof are as follows:

- We observe that points in $K_\infty(f)$ are associated to sequences x_l satisfying, for some N not depending on y ,

$$(1 + |x_l|)^{1+\frac{1}{N}} \nu(df(x_l)) \rightarrow 0.$$

- We consider, in a sphere with large radius r , the set D_i of points where the Kuo distance $\kappa(df(x))$ is attained by $\nabla f_i(x)$. We decompose $D_i \cap W_b$ relatively to r using Theorem 3.2, where W_b is the level surface $(f_1, \dots, \hat{f}_i, \dots, f_k)^{-1}(b)$.
- We prove that $\text{vol}_k K_\infty(f|_{D_i}) = 0$ using that

$$|\nabla f_i|_{W_b}(x)| \leq |x|^{-(1+\frac{1}{N})}.$$

3.1 Lemma on $K_\infty(f)$

In order to prove our Sard theorem, we shall use the fact that for a fixed mapping f , the convergence of $\nu(df(x_l))$ in the definition of $K_\infty(f)$ is actually faster than $|x_l|^{-1}$. To make this precise, for a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ and any $N \in \mathbb{N}^*$ we define

$$K_\infty^N(f) = \{y \in \mathbb{R}^k : \exists x_l \in \mathbb{R}^n, |x_l| \rightarrow \infty \\ \text{s.t. } f(x_l) \rightarrow y \text{ and } |x_l|^{1+\frac{1}{N}} \nu(df(x_l)) \rightarrow 0\}.$$

We have

Lemma 3.1. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ be a differentiable semialgebraic function. Then, there exists $N \in \mathbb{N}^*$ such that*

$$K_\infty(f) = K_\infty^N(f).$$

Proof. Let $i_n : \mathbb{R}^n \rightarrow S^n \subset \mathbb{R}^{n+1}$ be the inverse of the stereographic projection. Clearly, the graph of i is semialgebraic. Denote $\{\infty\} = S^n \setminus i_n(\mathbb{R}^n)$. In the sequel, we shall consider that $\mathbb{R}^n = i_n(\mathbb{R}^n) \subset S^n$. We compactify also $\mathbb{R}^k = i_k(\mathbb{R}^k) \subset S^k$ and $\mathbb{R} \subset S^1$ in the similar way. Let

$$\sigma(x) = |x| \nu(df(x)) \text{ for } x \in \mathbb{R}^n.$$

By Proposition 2.4, σ is a semialgebraic mapping. Finally, consider the mapping $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^k \times \mathbb{R} \subset S^k \times S^1$ defined by $\varphi(x) = (f(x), \sigma(x))$.

Recall now the classical lemma (cf. [2, Chap. 9], [3])

Lemma 3.2 (Wing Lemma). *Let Ω and B be two semialgebraic subsets of \mathbb{R}^m . Assume that $B = \overline{B} \subset \overline{\Omega} \setminus \Omega$. Then, there exists a semialgebraic set $A \subset \Omega$ such that*

$$B = \overline{A} \cap (\overline{\Omega} \setminus \Omega).$$

We apply the Wing Lemma for $\Omega = \text{graph}(\varphi) \subset S^n \times S^k \times S^1$, and B the closure of $\{\infty\} \times K_\infty(f) \times \{0\}$ in $S^n \times S^k \times S^1$. By Proposition 2.4 and the definition of $K_\infty(f)$ it follows that B is a closed semialgebraic set. Clearly $B \subset \overline{\Omega} \setminus \Omega$. So, there exists a semialgebraic set $A \subset \text{graph}(\varphi)$ such that

$$(3.1) \quad \overline{A} \cap (\{\infty\} \times S^k \times S^1) = B.$$

Let $\Gamma = \pi(A)$, where $\pi : \mathbb{R}^n \times \mathbb{R}^k \times \mathbb{R} \rightarrow \mathbb{R}^n$ is the projection. Equality (3.1) means that $\{\infty\} \times y \times \{0\} \in B$ if and only if there exists a sequence $x_l \in \Gamma$ with $|x_l| \rightarrow \infty$, $f(x_l) \rightarrow y$ and $\sigma(x_l) \rightarrow 0$. Let us define

$$\theta(r) = r \sup_{\substack{x \in \Gamma \\ |x|=r}} \nu(df(x)).$$

with the convention that $\theta(r) = 0$ if $\Gamma \cap S(r) = \emptyset$, where $S(r)$ is the sphere of radius r centered at 0. Clearly θ is semialgebraic. Note that for any sequence $x_l \in \Gamma$, $|x_l| \rightarrow \infty$ we have $\sigma(x_l) \rightarrow 0$, since B is compact. This implies that $\theta(r) \rightarrow 0$ as $r \rightarrow \infty$. So, by Puiseux, $\theta(r) \leq cr^{-\alpha}$ at ∞ for some $\alpha \in \mathbb{Q}_+^*$ and $c > 0$. Let $N \in \mathbb{N}^*$ be such that $\frac{1}{N} < \alpha$, then of course

$$r^{\frac{1}{N}} \theta(r) \rightarrow 0 \text{ when } r \rightarrow \infty.$$

So $K_\infty(f) \subset K_\infty^N(f)$, this completes the proof of Lemma 3.1 since obviously the reverse inclusion is satisfied. q.e.d.

3.2 Semialgebraic arcs at infinity

We shall use the following version of the curve selection lemma for semialgebraic sets (it can be easily obtained using a semialgebraic compactification of \mathbb{R}^n and the classical curve selection lemma, see [2], [3]).

Lemma 3.3 (Curve selection at infinity). *Let $A \subset \mathbb{R}^n$ and let $\phi : A \rightarrow \mathbb{R}^q$ be a semialgebraic map. Assume that there exists a sequence $x_l \in A$ such that $|x_l| \rightarrow \infty$ and $\phi(x_l) \rightarrow y$, for some $y \in \mathbb{R}^q$. Then there exists a semialgebraic arc $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^n$ such that $\gamma(t) \in A$, $\lim_{t \rightarrow \beta} |\gamma(t)| = +\infty$ and $\lim_{t \rightarrow \beta} \phi(\gamma(t)) = y$.*

Now consider such a semialgebraic arc $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^n$. Since $|\gamma'(t)| > 0$ for t close to β , we may reparametrize γ in such a way that $\beta = +\infty$ and $|\gamma(r)| = r$. Under this assumption we have:

Lemma 3.4. $\lim_{r \rightarrow \infty} |\gamma'(r)| = 1$; in particular, $\gamma'(r)$ is bounded for $r > 0$ large enough.

Proof. Since γ is semialgebraic, $\lim_{r \rightarrow \infty} \frac{\gamma(r)}{|\gamma(r)|}$ and $\lim_{r \rightarrow \infty} \frac{\gamma'(r)}{|\gamma'(r)|}$ exist. Hence, it is easily seen that these limits are equal. In other words $\cos \alpha(r) \rightarrow 1$, as $r \rightarrow \infty$, where $\alpha(r)$ is the angle between $\frac{\gamma(r)}{|\gamma(r)|} = \frac{\gamma(r)}{|r|}$ and $\frac{\gamma'(r)}{|\gamma'(r)|}$. Differentiation of $|\gamma(r)|^2 = r^2$ yields $|\gamma'(r)| = \frac{1}{\cos \alpha(r)}$. This implies the lemma. q.e.d.

3.3 Proof of main Theorem 3.1

Let us fix $N \in \mathbb{N}$ such that $K_\infty(f) = K_\infty^N(f)$ —cf. Lemma 3.1. By Proposition 2.6, we may replace the distance ν of Rabier by the distance κ of Kuo. For each $i \in \{1, \dots, k\}$, we define

$$D_i = \{x \in \mathbb{R}^n : \kappa(df(x)) = \text{dist}(\nabla f_i(x), V_i(x))\}$$

where $V_i(x)$ is the vector space generated by $\nabla f_j(x)$, $j = 1, \dots, k$ and $j \neq i$.

Clearly each D_i is semialgebraic in \mathbb{R}^n and $\mathbb{R}^n = \cup_{i=1}^k D_i$, so

$$K_\infty(f) = \bigcup_{i=1}^k K_\infty(f|_{D_i})$$

where

$$K_\infty(f|_{D_i}) = \{y \in \mathbb{R}^k : \exists x_l \in D_i, |x_l| \rightarrow \infty, f(x_l) \rightarrow y \\ \text{and } |x_l| \nu(df(x_l)) \rightarrow 0\}.$$

We shall prove the following

Lemma 3.5. $\text{vol}_k(K_\infty(f|_{D_i})) = 0$ for each $i \in \{1, \dots, k\}$. In particular $\dim K_\infty(f) < k$.

Proof. We will give the proof for $i = 1$, we write $D = D_1$, $\bar{f} = (f_2, \dots, f_k)$.

Let us fix B an open ball in \mathbb{R}^{k-1} and (α, β) an open bounded interval in \mathbb{R} . The lemma is clearly a consequence of

$$(3.2) \quad \text{vol}_k(K_\infty(f|_D) \cap (\alpha, \beta) \times B) = 0.$$

In order to prove Equality (3.2), we construct a family of sets Δ_r such that

$$\overline{\Delta_r} \supset K_\infty(f|_D) \cap (\alpha, \beta) \times B \text{ and } \text{vol}_k(\overline{\Delta_r}) \rightarrow 0 \text{ as } r \rightarrow \infty.$$

We first define

$$\tilde{\Sigma}_r = \{x \in D : |x| \geq r, f_1(x) \in (\alpha, \beta), \bar{f}(x) \in B \text{ and } |x|^{1+\frac{1}{N}} \kappa(df(x)) \leq 1\}$$

where $r > 0$, and put

$$\Delta_r = f(\tilde{\Sigma}_r) \text{ and finally } \Delta = \bigcap_{r>0} \overline{\Delta_r}.$$

Every Δ_r is semialgebraic, hence we have $\text{vol}_k(\overline{\Delta_r}) = \text{vol}_k(\Delta_r)$ and consequently

$$\text{vol}_k(\Delta) = \lim_{r \rightarrow \infty} \text{vol}_k(\Delta_r)$$

since the family $(\Delta_r)_{r>0}$ is decreasing with respect to $r \rightarrow \infty$.

It is clear that

$$K_\infty(f|_D) \cap (\alpha, \beta) \times B \subset \Delta,$$

so it is enough to prove that $\text{vol}_k(\Delta) = 0$. First, using Fubini's theorem we write

$$\text{vol}_k(\Delta_r) = \int_B m_r(b) db$$

where db stands for the Lebesgue measure on \mathbb{R}^{k-1} , and

$$m_r(b) = \text{vol}_1(\{y_1 \in \mathbb{R} : (y_1, b) \in \Delta_r\}).$$

Clearly, each m_r is measurable. Moreover, for fixed $b \in B$, the function $r \mapsto m_r(b) \geq 0$ is decreasing. Let

$$m(b) = \lim_{r \rightarrow \infty} m_r(b).$$

By Lebesgue's theorem on bounded convergence, we obtain

$$\text{vol}_k(\Delta) = \int_B m(b) db.$$

Now the final point in the proof of Equality (3.2) is the fact that $m \equiv 0$, which follows from the next lemma. q.e.d.

Lemma 3.6. *There exists a constant $c > 0$ such that, for r large enough*

$$m_r(b) \leq cr^{-\frac{1}{N}}.$$

Proof. To prove Lemma 3.6, we introduce the semialgebraic family

$$\Sigma_{r,b} = \tilde{\Sigma}_r \cap \bar{f}^{-1}(b) \cap S(r),$$

where $b \in B$, $r > 0$, and next we write

$$\tilde{\Sigma}_{r,b} = \tilde{\Sigma}_r \cap \bar{f}^{-1}(b) = \bigcup_{s \geq r} \Sigma_{s,b}.$$

Note that

$$m_r(b) = \text{vol}_1(f_1(\tilde{\Sigma}_{r,b})).$$

It follows from Theorem 3.2 that there exists a finite family $L^i \subset \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^{k-1}$, $i \in I$ of semialgebraic sets such that

$$\Sigma_{r,b} = \bigcup_{i \in I} L_{r,b}^i.$$

Each $L_{r,b}^i$ has the Whitney property with constant M (some of $L_{r,b}^i$ may be empty).

Recall that the condition $|x|^{1+\frac{1}{N}} \kappa(df(x)) \leq 1$ for $x \in \bar{f}^{-1}(b) = W_b$ means that

$$(3.3) \quad |\nabla f_1|_{W_b}(x)| \leq |x|^{-(1+\frac{1}{N})}.$$

Hence, by the mean value theorem $f_1(L_{r,b}^i)$ is a segment of length $d(r)$ where

$$(3.4) \quad d(r) \leq 2Mr \sup_{L_{r,b}^i} |\nabla f_1|_{W_b}| \leq 2Mr^{-\frac{1}{N}}.$$

Fix $b \in B$, $i \in I$ and assume that $L_{r,b}^i \neq \emptyset$ for any r large enough. Applying the [curve selection lemma at infinity](#), we obtain a semialgebraic arc $\gamma : [r, +\infty) \rightarrow \mathbb{R}^n$ such that $\gamma(\zeta) \in L_{\zeta,b}^i$. In particular, $\gamma(\zeta) \in \bar{f}^{-1}(b) = W_b$ and $|\gamma(\zeta)| = \zeta$.

By Lemma 3.4, we may suppose that $|\gamma'(\zeta)| \leq 2$. So we can easily compute length of $f_1 \circ \gamma([r, +\infty))$; namely, by (3.3) we have

$$(3.5) \quad \int_r^{+\infty} |(f_1 \circ \gamma)'(\zeta)| d\zeta \leq 2 \int_r^{+\infty} \zeta^{-(1+\frac{1}{N})} d\zeta = 2Nr^{-\frac{1}{N}}.$$

Thus, by (3.4) and (3.5), $f_1(\bigcup_{\zeta \geq r} L_{\zeta, b}^i)$ is contained in a segment of length

$$(4M + 2N)r^{-\frac{1}{N}}.$$

Therefore $f_1(\tilde{\Sigma}_{r, b})$ is contained in $\#I$ segments of this length. Put $c = (\#I)(4M + 2N)$; we have

$$m_r(b) \leq cr^{-\frac{1}{N}}$$

and Lemma 3.6 follows. q.e.d.

As we already seen the map $\varphi = (f, \sigma)$ is semialgebraic, and so $K(f)$ is a semialgebraic subset of \mathbb{R}^k .

From Lemma 3.5 and the usual semialgebraic Sard's theorem (see [3]) it follows that $\dim K(f) < k$.

The fact that f is a fibration is a consequence of Theorem 2.1. This ends the proof of Theorem 3.1.

3.4 An alternative proof of Lemma 3.6 via Teissier's theorem

Let us recall the following result due to B. Teissier [21]. We state only the semialgebraic version which is of our interest

Theorem 3.3 (Teissier). *Let $B \subset \mathbb{R}^n \times \mathbb{R}^p$ be a semialgebraic set. Assume that for any $t \in \mathbb{R}^p$ the set $B_t = \{x \in \mathbb{R}^n : (x, t) \in B\}$ is contained in the unit closed ball in \mathbb{R}^n . Then there exists a constant $M > 0$, depending on B , such for any $t \in \mathbb{R}^p$: any two points in the same connected component of B_t can be joined in B_t by a piecewise smooth curve of length at most M .*

The original proof of Teissier (for subanalytic sets) used Hironaka's resolution of singularities, but now there are many simplified proofs by R. Hardt, the first named author, Y. Yomdin [23]. S. K. Donaldson [4] obtained an explicit estimate for M in some special cases.

We obtain the following from Theorem 3.3 as an immediate corollary:

Corollary 3.1. *Let $A \subset \mathbb{R}^n \times \mathbb{R}^p$ be a semialgebraic set. Assume that for any $t \in \mathbb{R}^p$ the set $A_t = \{x \in \mathbb{R}^n : (x, t) \in A\}$ is contained in the ball $\{x \in \mathbb{R}^n : |x| \leq r(t)\}$, where $r : \mathbb{R}^p \rightarrow \mathbb{R}$ is some semialgebraic (not necessarily continuous) positive function. Then there exists a constant $M > 0$, depending on A , such that, for any $t \in \mathbb{R}^p$, any two points in the same connected component of A_t can be joined in A_t by a piecewise smooth curve of length at most $Mr(t)$.*

Indeed it is enough to apply Teissier's theorem to the semialgebraic family

$$B_t = \frac{1}{r(t)} A_t, \quad t \in \mathbb{R}^p.$$

Let us come back to the proof of Lemma 3.6. Recall that

$$\Sigma_{r,b} = \tilde{\Sigma}_r \cap \bar{f}^{-1}(b) \cap S(r).$$

It follows from Hardt's semialgebraic triviality theorem (see [2, Chap. 9]) that there exists a finite family $L^i \subset \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^{k-1}$, $i \in I$ of semialgebraic sets such that

$$\Sigma_{r,b} = \bigcup_{i \in I} L_{r,b}^i.$$

But this time $L_{r,b}^i$ denote a connected component of $\Sigma_{r,b}$ or an empty set. By Corollary 3.1 any two points in $L_{r,b}^i$ can be joined (in $L_{r,b}^i$) by a piecewise smooth curve of length at most Mr . So we can obtain estimate (3.4) by the same argument as before. The rest of the proof of Lemma 3.6 need not to be modified.

Remark 3.2. We have actually proved that, for generic $b \in \mathbb{R}^{k-1}$, the set $\Delta \cap \{x \in \mathbb{R}^k : (x_2, \dots, x_k) = b\}$ has at most s points, where s is a bound for the number of connected components of any $\Sigma_{r,b}$. This number s can be easily estimated from above (cf. [3]) in terms of the degrees of the polynomials describing the graph of f . In particular, if f_1, \dots, f_k are polynomials of degree less than d , then, by [3, Proposition 4.4.5],

$$s = s(n, d, k) \leq (2d + 1)(4d - 3)^n.$$

Hence, if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a polynomial of degree d then

$$\#K_\infty(f) \leq (2d + 1)(4d - 3)^n.$$

3.5 The set J_f of Z. Jelonek

To end this section we will prove that, in some cases, our $K_\infty(f)$ is equal to the set J_f of Z. Jelonek.

Let us consider a continuous mapping $f : \mathbb{K}^n \rightarrow \mathbb{K}^n$. We say (cf. [7]) that f is *proper at a point* $y \in \mathbb{K}^n$ if there exists an open neighborhood U of y such that the restriction $f|_{f^{-1}(U)} : f^{-1}(U) \rightarrow U$ is a proper map. We denote by J_f the set of points at which f is not proper, we always have the inclusion $K_\infty(f) \subset J_f$.

Now we prove the following:

Proposition 3.1. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a C^1 semialgebraic mapping, then $J_f = K_\infty(f)$.*

Proof. Observe that $y \in J_f$ implies that there exists a sequence $x_l \in \mathbb{R}^n$ such that $|x_l| \rightarrow \infty$ and $f(x_l) \rightarrow y$. Since f is semialgebraic, using a standard curve selection (see Subsection 3.2) we can find a C^1 semialgebraic arc $\gamma : [a, \infty) \rightarrow \mathbb{R}^n$ such that $|\gamma(r)| = r \rightarrow \infty$ and $f(\gamma(r)) \rightarrow y$. Hence in particular $|\gamma'(r)| \geq 1$. Note that a continuous semialgebraic arc $\eta : [\alpha, \beta) \rightarrow \mathbb{R}^n$ has a limit (in \mathbb{R}^n) at β , if and only if η is of finite length.

Assume that $y \in J_f$ and $y \notin K_\infty(f)$. It means that for some $b, c > 0$ we have

$$|df(\gamma(r))(\gamma'(r))| \geq \frac{c}{r}$$

for any $r \geq b$. But this implies that length of $f(\gamma)$ is not less than $c \int_b^\infty \frac{dr}{r} = \infty$, since f is injective on γ for r large enough. So $f(\gamma(r))$ has no limit in \mathbb{R}^n as $r \rightarrow \infty$, which is a contradiction. q.e.d.

We shall use the above proposition in a discussion (in the next section) on the dimension of $K_\infty(f)$ in the complex case.

Recall that we have always $K_\infty(f) \subset \tilde{K}_\infty(f) \subset J_f$. Thus Proposition 3.1 yields

Corollary 3.2. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a C^1 semialgebraic mapping, then $K_\infty(f) = \tilde{K}_\infty(f)$.*

This is in strong contrast with a case, discussed in Section 2, where the dimension of the target was smaller than the dimension of the source. Corollary 3.2 seems to be a new result also in the complex case (even for $n = 2$) and may be of some interest for Jacobian Conjecture.

Proposition 3.1 holds actually in a more general case. It is enough to assume that the generic fibers of $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ are compact. More precisely we have

Theorem 3.4. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$, $k \leq n$ be a C^1 semialgebraic mapping. Assume that the set of regular points of f is dense and that $f^{-1}(y)$ is compact for any $y \in \mathbb{R}^k \setminus K_0(f)$. Then $K_\infty(f) = \tilde{K}_\infty(f) = J_f$.*

Proof. Clearly, by previous remarks, it is enough to prove that $J_f \subset K_\infty(f)$. Assume this is not the case, it means that there exists $y \notin K_\infty(f)$ and a sequence $x_l \in \mathbb{R}^n$ such that $|x_l| \rightarrow \infty$ and $f(x_l) \rightarrow y$. By the curve selection lemma we obtain a C^1 semialgebraic arc

$\gamma : [a, \infty) \rightarrow \mathbb{R}^n$ such that $|\gamma(r)| \rightarrow \infty$ and $f(\gamma(r)) \rightarrow y$. For a large enough the set $\Gamma = f(\gamma(a, \infty))$ is a smooth curve of finite length. Moreover, since $y \notin K_\infty(f)$ there exists a constant $c > 0$ such that

$$(3.6) \quad (1 + |x|)\nu(df(x)) \geq c$$

for any x such that $f(x) \in \Gamma$ and $|x| \geq R$, where $R > 0$ is large enough. Since the set of regular values of f is dense we can suppose that $\Gamma \cap K_0(f) = \emptyset$. Recall that $f^{-1}(f(\gamma(a)))$ is compact, so we may assume that it is contained in $B(0, R)$.

We are going to prove that $f^{-1}(\Gamma)$ is bounded which will be a contradiction.

Denote by g the map $f|_{f^{-1}(\Gamma)}$, and by $\delta(g(x))$ the unit tangent vector field to Γ in the direction of y . Using the right inverse $s(x)$ of $dg(x)$ (see Section 4 of [20]), we can lift δ to a C^0 vector field $X(x) = s(x)\delta(g(x))$ on $f^{-1}(\Gamma)$. In fact $X = \frac{\nabla g}{|\nabla g|^2}$, where ∇g is the gradient of g with respect to the induced metric on $f^{-1}(\Gamma)$.

Since $\nu(dg(x)) = \nu(df(x)|_{f^{-1}(\Gamma)}) \geq \nu(df(x))$ (use Proposition 2.3), Condition (3.6) with f replaced by g is still true outside of $B(0, R)$.

Take a point $z \in (\mathbb{R}^n \setminus B(0, R)) \cap f^{-1}(\Gamma)$, by Peano's theorem there exists a local integral curve of the field $-X$ starting at z ; denote it by $\phi(t)$ with $t \in [\alpha, \beta)$. Using Gronwall lemma and the bound

$$|X(x)| \leq \frac{2(1 + |x|)}{c}$$

which comes from (3.6), we easily obtain that

$$|\phi(t)| \leq |z| \exp(\beta - \alpha) + \exp\left(\frac{2}{c}(\beta - \alpha)\right) - 1$$

for $t \in [\alpha, \beta)$. This inequality implies that ϕ is Lipschitz on $[\alpha, \beta)$ and so ϕ has a limit at β and can be extended.

Observe that t is the arc length on Γ , since $dg(\phi(t))\phi'(t) = \delta_{g(\phi(t))}$. This implies that ϕ can be extended in such a way that in finite time it goes into $B(0, R)$, since $g^{-1}(g(\gamma(a))) \subset B(0, R)$. But, reversing time, we see (again by Gronwall lemma) that $\phi(t)$ has to stay in the ball of radius $R \exp(\text{length}(\Gamma)) + \exp\left(\frac{2}{c} \text{length}(\Gamma)\right) - 1$.

Thus $f^{-1}(f(\gamma(b)))$ is uniformly bounded irrespective of b . Hence the set $f^{-1}(\Gamma)$ is bounded, which is a contradiction. \square

Remark 3.3. P. J. Rabier pointed out to us that the same result is valid for general C^2 mappings from \mathbb{R}^n to \mathbb{R}^k , if $f^{-1}(y)$ is compact for y in a dense subset of \mathbb{R}^k : By standard transversality arguments one can construct a perturbation with compact support of f for which y becomes a regular value. This does not change $K_\infty(f)$ and J_f near y , thus we can conclude using Rabier's fibration theorem. Combining this with our proof we obtain that, in fact, Theorem 3.4 is valid for any C^1 mappings with generically compact fibers.

Remark 3.4. In a recent paper [8] Z. Jelonek proved that if a polynomial mapping satisfies the assumptions of Theorem 3.4, then the corresponding semialgebraic set $K_\infty(f) = J_f$ is \mathbb{R} -uniruled—it means that by every point pass a curve (in J_f) with polynomial parametrization.

4. The complex case

Suppose now that $f : \mathbb{C}^n \rightarrow \mathbb{C}^k$ is a complex polynomial mapping. We know, by Theorem 3.1, that $K_\infty(f)$ is a nowhere dense semialgebraic subset of \mathbb{C}^k . But in this case, we prove more:

Theorem 4.1. *If $f : \mathbb{C}^n \rightarrow \mathbb{C}^k$ is a complex polynomial mapping, then $K_\infty(f)$ is a complex algebraic set and $\dim_{\mathbb{C}} K_\infty(f) < k$.*

Proof. We shall prove that $K_\infty(f)$ is a complex algebraic set. We can assume that f is dominant, that is the generic rank of f is k , as otherwise $K_\infty(f)$ is equal to the closure of the image of f . Since the closure is the same in the strong and in Zariski topology, we deduce that $K_\infty(f)$ is a complex algebraic set.

As before we replace Rabier's ν invariant by the invariant κ of Kuo. Let $f = (f_1, \dots, f_k)$. For any $j = 1, \dots, k$ we denote

$$\tilde{f}_j = (f_1, \dots, f_{j-1}, f_{j+1}, \dots, f_k) : \mathbb{C}^n \rightarrow \mathbb{C}^{k-1}.$$

Let $V_j(z)$ be the kernel of $d\tilde{f}_j(z)$ and let $w_j(z)$ denote the restriction of $df_j(z)$ to the linear subspace $V_j(z)$. Recall that by the definition (2.2) of κ

$$(4.1) \quad \kappa(df(z)) = \min_{1 \leq j \leq k} \|w_j(z)\|.$$

Let $z = (z_1, \dots, z_n) \in \mathbb{C}^n$. For each $s = 1, \dots, n$ we will use the following

change of coordinates:

$$\tau_s(z) = \left(\frac{z_1}{z_s}, \dots, \frac{z_{s-1}}{z_s}, \frac{1}{z_s}, \frac{z_{s+1}}{z_s}, \dots, \frac{z_n}{z_s} \right),$$

which sends $(z_s = 0)$ to ∞ . Denote by $\mathbb{G} = \mathbb{G}_{n-k+1}(\mathbb{C}^n \times \mathbb{C})$ the Grassmanian of $(n - k + 1)$ -planes in $\mathbb{C}^n \times \mathbb{C}$. For fixed j and s we consider rational mappings

$$A_s^j : \mathbb{C}^n \setminus \{z_s = 0\} \rightarrow \mathbb{C}^k \times \mathbb{G}$$

given by the formula

$$A_s^j(z) = (f(\tau_s(z)), W_s^j(z)).$$

Here $W_s^j(z)$ stands for the graph of $\frac{1}{z_s}w_j(z)$. To be more precise $A_s^j(z)$ is well defined for those z for which $V_j(z)$ is of dimension $(n - k + 1)$. Recall that we assumed that f is dominant, hence this condition holds outside a nowhere dense algebraic set. Let $\Lambda = \mathbb{G}_{n-k+1}(\mathbb{C}^n \times 0) \subset \mathbb{G}$. Note that each $\lambda \in \Lambda$ can be seen as a graph of constant (equal to 0) mapping on the $(n - k + 1)$ -plane λ . Put

$$B_s^j = \overline{\text{graph}(A_s^j)} \cap \{z \in \mathbb{C}^n : z_s = 0\} \times \mathbb{C}^k \times \Lambda.$$

The closure is taken in the strong topology, but it is equal to the Zariski closure. Hence B_s^j is an algebraic set. Let $\pi : \mathbb{C}^n \times \mathbb{C}^k \times \mathbb{G} \rightarrow \mathbb{C}^k$ denote the projection. Finally let

$$K_s^j = \pi(B_s^j).$$

Clearly each K_s^j is a constructible set. Assume that we have proved:

Lemma 4.1.
$$K_\infty(f) = \bigcup_{s=1, j=1}^{n, k} K_s^j.$$

This, of course, implies that $K_\infty(f)$ is constructible. On the other hand, the set $K_\infty(f)$ is closed, so it must be algebraic.

So we are left with proving Lemma 4.1. Suppose that $y \in K_s^j$, then we can find a real Puiseux arc $\eta = (\eta_1, \dots, \eta_n) : (0, \varepsilon) \rightarrow \mathbb{C}^n$, $|\eta_s(t)| = t$ such that

$$f \circ \tau_s \circ \eta(t) \rightarrow y, \text{ when } t \rightarrow 0$$

and

$$W_s^j(\tau_s(\eta(t))) \rightarrow W_s^j, \text{ when } t \rightarrow 0$$

for some $W_s^j \in \Lambda$. By (4.1) this means that $y \in K_\infty(f)$.

The reverse inclusion is easily obtained. Take $y \in K_\infty(f)$, then there is an arc going to infinity, say $z(t) = (z_1(t), \dots, z_n(t))$, such that $f(z(t)) \rightarrow y$ and

$$(4.2) \quad |z(t)|\kappa(df(z(t))) \rightarrow 0.$$

Choose a coordinate z_s in such a way that $z_s(t)$ goes to infinity faster than the other coordinates, then choose j such that $\kappa(df(z(t))) = \|w_j(z(t))\|$ and proceed using A_s^j . Since \mathbb{G} is compact there exists a limit W_s^j of graphs of $z_s(t)w_j(z(t))$. It follows, by (4.2), that $W_s^j \in \Lambda$. So $(0, y, W_s^j) \in B_s^j$ and consequently $y \in K_s^j$. q.e.d.

Recall that, by a result of Z. Jelonek [7], if $f : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is a polynomial dominant mapping with nonempty J_f , then J_f is a hypersurface (actually this is a \mathbb{C} -uniruled hypersurface, its degree is effectively bounded). So, by our Proposition 3.1, $K_\infty(f)$ is also a \mathbb{C} -uniruled hypersurface in this case. One could conjecture that, for a general polynomial dominant mapping $f : \mathbb{C}^n \rightarrow \mathbb{C}^k$ with $k < n$, $K_\infty(f)$ must be a hypersurface or an empty set. The following example (suggested by Z. Jelonek) shows that this not the case.

Example 4.1. Consider the polynomial mapping $f : \mathbb{C}^3 \rightarrow \mathbb{C}^2$, given by $f(x, y, z) = (xy, xz)$. Using Proposition 2.5 it is easily seen that $\nu(df(\xi)) = |x|$ for $\xi = (x, y, z)$. Let $\xi_n = (x_n, y_n, z_n)$ be a sequence which tends to infinity and assume that

$$|\xi_n||x_n| = |\xi_n|\nu(df(\xi_n)) \rightarrow 0.$$

Hence $|x_n| \rightarrow 0$ and therefore $f(\xi_n) \rightarrow 0$. So 0 is the only asymptotic critical value of f .

However we may still conjecture that $K_\infty(f)$ is \mathbb{C} -uniruled in the sense that by each point of $K_\infty(f)$ passes a curve which has a polynomial (possibly constant) parametrization.

Remark 4.1. After the first version of the paper was written (november 1999) we learned that T. Gaffney [6] studied the set $K_\infty(f)$ in the complex setting. In particular he proved (without using Rabier's result) that the mapping is a locally trivial fibration over the complement of $K_\infty(f)$. However his definition of ν is not exactly the same as

ours. We leave to the reader to prove that in fact both definitions are equivalent.

References

- [1] D. d'Acunto, *Valeurs critiques asymptotiques de fonctions définissables dans une structure o-minimale*, Ann. Polon. Math. (to appear).
- [2] J. Bochnak, M. Coste & M.-F. Roy, *Real algebraic geometry*, E.M.G., Springer, Vol. 36, 1998.
- [3] R. Benedetti & J.-J. Risler, *Real algebraic and semialgebraic sets*, Hermann, 1990.
- [4] S. K. Donaldson, *Symplectic submanifolds and almost-complex geometry*, J. Differential Geom. **44** (1996) 666–705.
- [5] H.V. Hà, *Nombres de Lojasiewicz et singularités à l'infini des polynômes de deux variables complexes*, C. R. Acad. Sci. Paris, Série I Math. **311** (1990) 429–432.
- [6] T. Gaffney, *Fibers of polynomial mappings at infinity and a generalized Malgrange Condition*, Compositio Math. **119** (1999) 157–167.
- [7] Z. Jelonek, *Testing sets for polynomial mapping*, Math. Ann. **315** (1999) 1–35.
- [8] ———, *Geometry of real polynomial mappings*, Math. Z. (to appear).
- [9] K. Kurdyka & P. Orro, *Distance géodésique sur un sous-analytique*, Revista matemática de la Universidad Computense de Madrid, Vol. 10, 1997, num. supl.
- [10] K. Kurdyka, *On a subanalytic stratification satisfying a Whitney property with exponent 1*, Real algebraic geometry, Proc., Rennes 1991, L.N.M. **1524** (1992) 316–322.
- [11] ———, *Linear bounds for resolutions of 1-boundaries in subanalytic sets*, Real analytic and algebraic geometry, Proc., Trento 1992, Walter de Gruyter, 1995, 189–196.
- [12] K. Kurdyka & T. Mostowski, *On the gradient conjecture of R. Thom*, Preprint, 1998.
- [13] K. Kurdyka, T. Mostowski & A. Parusiński, *Proof of the gradient conjecture of R. Thom*, Ann. of Math. **152** (2000) 763–792.
- [14] T. C. Kuo, *Characterizations of V-sufficiency of jets*, Topology **11** (1972) 115–131.
- [15] R. S. Palais, *Lusternik-Schnirelman theory on Banach manifolds*, Topology **5** (1966) 115–132.

- [16] A. Parusiński, *On the bifurcation set of complex polynomial with isolated singularities at infinity*, Compositio Math. **97** (1995) 369–384.
- [17] ———, *A note on singularities at infinity of complex polynomials*, Banach center publication, Vol. 39, 1997, 131–141.
- [18] ———, *Lipschitz properties of semianalytic sets*, Ann. Inst. Fourier **38** (1988) 189–213.
- [19] L. Paunescu & A. Zaharia, *Remarks on the Milnor fibration at infinity*, Preprint, 1999.
- [20] P. J. Rabier, *Ehresmann’s fibrations and Palais-Smale conditions for morphisms of Finsler manifolds*, Ann. of Math. **146** (1997) 647–691.
- [21] B. Teissier, *Sur trois questions de finitude en géométrie analytique réelle, appendix to F. Tréves, on the local solvability and the local integrability of systems of vector fields*, Acta Math. **151** (1983) 38–48.
- [22] M. Tibăr & A. Zaharia, *Asymptotic behaviour of families of curves*, Manuscripta Math. **99** (1999) 383–393.
- [23] Y. Yomdin, *The geometry of critical and near-critical values of differentiable mappings*, Math. Ann. **4** (1983) 495–515.

UNIVERSITÉ DE SAVOIE, CAMPUS SCIENTIFIQUE, FRANCE
UNIwersytet Jagielloński, POLAND