

Linear independence between odd and even periods of modular forms

Hui Xue

ABSTRACT. We investigate the linear dependence between an odd period and an even period of modular forms. We show that two periods of different parity are linearly independent provided that the even period has index at least 6 or the odd period has index at least 7.

CONTENTS

1. Introduction and statements of results	521
2. The proofs	524
Acknowledgment	531
References	531

1. Introduction and statements of results

For each even integer $k \geq 4$, let M_k be the space of modular forms of weight k for $SL_2(\mathbb{Z})$, and let S_k be its subspace of cuspforms. For each $0 \leq t \leq k - 2$, the t^{th} period of $f \in S_k$ is defined as [6]

$$r_t(f) := \int_0^{i\infty} f(z)z^t dz = \frac{t!}{(-2\pi i)^{t+1}} L(f, t + 1). \quad (1)$$

Here, the L -series of a cuspform $f(z) = \sum_{n=1}^{\infty} a_n(f)e^{2\pi inz} \in S_k$ is $L(f, s) = \sum_{n=1}^{\infty} \frac{a_n(f)}{n^s}$. Each r_t defines a linear map from S_k to \mathbb{C} , that is $r_t \in S_k^*$ (the dual space of S_k).

The set of odd periods $\{r_{2i+1}\}_{i=0}^{k/2-2}$ and the set of even periods $\{r_{2i}\}_{i=0}^{k/2-1}$ behave differently, and they are subject to many linear dependence relations, called the Eichler-Shimura relations; see Manin [6] for more details. However, not much is known about the linear independence of a subset of the periods: the first work in this direction seems to be [1], in which Fukuhara found an explicit subset of odd periods that forms a basis for S_k^* . As a corollary, Fukuhara [2] found a special basis for M_k consisting of products of two Eisenstein series.

Received June 7, 2023.

2010 *Mathematics Subject Classification.* 11F67, 11F30.

Key words and phrases. Eisenstein series, Rankin's identity, linear independence, periods of modular forms.

Most recently, Lei et al. [5, 4] have provided some evidence for the linear independence of odd periods and even periods, respectively. The main theme of [5, 4] is that odd or even periods of modular forms are linearly independent unless forced by dimension considerations. On the other hand, very little seems to be known about the relationship between even and odd periods: for instance, the Eichler-Shimura relations only address them separately. In the present paper, we will extend the ideas of [5, 4] to provide some evidence for the linear independence between odd and even periods. More precisely, we will show the following.

Theorem 1.1. *Let ℓ and ℓ' be positive even integers such that $\ell < \frac{k}{2} - 1$ and $\ell' \leq \frac{k}{2}$, and suppose that $\ell \geq 6$ or $\ell' \geq 8$. If $\dim S_k \geq 2$, then the even period r_ℓ and the odd period $r_{\ell'-1}$ are linearly independent.*

The restriction to periods r_ℓ and $r_{\ell'-1}$ for even integers $\ell < \frac{k}{2} - 1$ and $\ell' \leq \frac{k}{2}$ is due to the Eichler-Shimura relations $r_\ell + r_{k-2-\ell} = 0$ and $r_{\ell'-1} - r_{k-1-\ell'} = 0$. In fact, numerical computation done by Daozhou Zhu ([10]) shows that Theorem 1.1 holds true for all positive even integers ℓ and ℓ' for $k \leq 100$ and $\dim S_k \geq 2$. So, we propose the following natural speculations.

Conjecture 1.2. (1) *Let ℓ and ℓ' be positive even integers such that $\ell < \frac{k}{2} - 1$ and $\ell' \leq \frac{k}{2}$. If $\dim S_k \geq 2$, then the even period r_ℓ and the odd period $r_{\ell'-1}$ are linearly independent.*

(2) *More generally, suppose $2 \leq \ell_1 < \ell_2 < \dots < \ell_a < \frac{k}{2} - 1$ and $2 \leq \ell'_1 < \dots < \ell'_b \leq \frac{k}{2}$ are even integers. If $a + b \leq \dim S_k$, then the set of periods $\{r_{\ell_1}, \dots, r_{\ell_a}, r_{\ell'_1-1}, \dots, r_{\ell'_b-1}\}$ is linearly independent.*

We now give an account of the main idea of the proof. For an even integer $k \geq 2$, let $E_k(z)$ denote the normalized Eisenstein series of weight k given by

$$E_k(z) = 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n)q^n,$$

where B_k is the k -th Bernoulli number, $\sigma_{k-1}(n) = \sum_{d|n} d^{k-1}$ and $q = e^{2\pi iz}$. It should be noted that $E_2(z)$, although holomorphic, is not a modular form; it is a quasi-modular form instead [9].

Let $f \in M_k$ and $g \in M_\ell$. For some integer $d \geq 0$, the d -th Rankin-Cohen bracket of f and g is defined as [8, (1)]:

$$[f, g]_d = \sum_{0 \leq r \leq d} (-1)^r \binom{d+k-1}{d-r} \binom{d+\ell-1}{r} f^{(r)} g^{(d-r)},$$

where $f^{(r)} := \frac{1}{(2\pi i)^r} \frac{d^r f}{dz^r}$ is the normalized r th derivative of f with respect to z . In this paper, we are only interested in the cases when both f and g are Eisenstein

series and $d = 0, 1$. In order to include E_2 in the Rankin-Cohen brackets, we also define ([3, p. 214 (ii)] and [4, (1.2)])

$$[E_k, E_2]_d := \sum_{0 \leq r \leq d} (-1)^r \binom{d+k-1}{d-r} \binom{d+\ell-1}{r} E_k^{(r)} E_2^{(d-r)} \tag{2}$$

$$- (-1)^d \frac{12}{k+d} E_k^{(d+1)}$$

for $k > 2$, where $E_2^{(i)}$ on the right hand side is the normalized i^{th} derivative of E_2 with respect to the variable z , for $0 \leq i \leq d$. Then, $[E_k, E_2]_d$ is a modular form in M_{k+2+2d} .

Next, we recall the Rankin’s identity for the two cases $d = 0, 1$. Let $k > \ell \geq 2$ and $k' \geq \ell' \geq 2$ be even integers such that $K := k + \ell + 2 = k' + \ell'$. Then, by ([7, (77)], [3, pp. 213-215]) we have the following formulas for the Petersson inner products

$$\langle g, E_{k'} E_{\ell'} - E_K \rangle = (-1)^{\frac{k'}{2}} \frac{\Gamma(K-1)\Gamma(k')}{(4\pi)^{K-1}(2\pi)^{k'}} \frac{2k'}{B_{k'}} \frac{2\ell'}{B_{\ell'}} L(g, K-1)L(g, k') \tag{3}$$

and

$$\langle g, [E_k, E_\ell]_1 \rangle = (-1)^{\frac{k}{2}+1} \frac{\Gamma(K-1)\Gamma(k+1)}{(4\pi)^{K-1}(2\pi)^k} \frac{2k}{B_k} \frac{2\ell}{B_\ell} L(g, K-2)L(g, k+1) \tag{4}$$

For later application we need to normalize $[E_k, E_\ell]_1$ and $E_k E_\ell - E_K$, so that their q -coefficients become 1. It follows from [4, Section 2] that

$$a_1(E_{k'} E_{\ell'} - E_K) = \begin{cases} -\frac{2k'}{B_{k'}} - \frac{2\ell'}{B_{\ell'}} + \frac{2K}{B_K} & \text{if } \ell' \geq 4 \\ -24 \left(1 + \frac{K-2}{12B_{K-2}} - \frac{1}{B_{K-2}} - \frac{K}{12B_K} \right) & \text{if } \ell' = 2 \end{cases}, \tag{5}$$

and

$$a_1([E_k, E_\ell]_1) = \begin{cases} \frac{2k\ell}{B_k} - \frac{2k\ell}{B_\ell} & \text{if } \ell \geq 4 \\ \frac{4k}{B_k} - \frac{24k}{(k+1)B_k} - \frac{4k}{B_2} & \text{if } \ell = 2 \end{cases}. \tag{6}$$

We normalize $[E_k, E_\ell]_1$, denoted $\Delta_{k,\ell}^1(z)$, so that its q -coefficient $a_1(\Delta_{k,\ell}^1)$ becomes 1. Similarly, we normalize $E_k E_\ell - E_K$, denoted $\Delta_{k',\ell'}^1(z)$, so that $a_1(\Delta_{k',\ell'}^1) = 1$. The following result follows immediately from (3)-(6).

Proposition 1.3. *Let $k > \ell \geq 2$ and $k' \geq \ell' \geq 2$ be even integers such that $K := k + \ell + 2 = k' + \ell'$. Let \mathcal{H}_K denote the set of normalized Hecke eigenforms in S_K . Then*

$$\Delta_{k,\ell}^1 = A_{k,\ell}^1 \cdot \sum_{g \in \mathcal{H}_K} \frac{L(g, K-2)L(g, k+1)}{\langle g, g \rangle} g,$$

where

$$A_{k,\ell}^1 := (-1)^{\frac{k}{2}+1} \frac{\Gamma(K-1)\Gamma(k+1)}{(4\pi)^{K-1}(2\pi)^k} \frac{2k}{B_k} \frac{2\ell}{B_\ell} \cdot \frac{1}{a_1([E_k, E_\ell]_1)}$$

with $a_1([E_k, E_\ell]_1)$ given by (6). Also,

$$\Delta_{k', \ell'} = A_{k', \ell'} \cdot \sum_{g \in \mathcal{H}_K} \frac{L(g, K-1)L(g, k')}{\langle g, g \rangle} g,$$

where

$$A_{k', \ell'} := (-1)^{\frac{k'}{2}} \frac{\Gamma(K-1)\Gamma(k')}{(4\pi)^{K-1}(2\pi)^{k'}} \frac{2k' 2\ell'}{B_{k'} B_{\ell'}} \cdot \frac{1}{a_1(E_{k'}E_{\ell'} - E_K)}$$

with $a_1(E_{k'}E_{\ell'} - E_K)$ given in (5).

Remark 1.4. We want to point out that the actual values of $A_{k, \ell}^1$ and $A_{k', \ell'}$ are not important, as long as they are nonzero and are independent of $g \in \mathcal{H}_K$; see Section 2. It is also important to note that for each $g \in \mathcal{H}_K$ the value $L(g, k+1)$ is positive because $k+1 > \frac{K+1}{2}$ is within the region of absolute convergence for the Euler product of $L(g, s)$.

Now, assume that ℓ and ℓ' satisfy the conditions of Theorem 1.1, and that r_ℓ and $r_{\ell'}$ are linearly dependent. Our strategy is to compare the a_2 Fourier coefficients of $\Delta_{k, \ell}^1$ and $\Delta_{k', \ell'}$, to reach a contradiction. On one hand, using results from [4, 5], we show (Lemma 2.2) that for $\ell \geq 6$ or $\ell' \geq 8$ and $K \geq 100$

$$|a_2(\Delta_{k, \ell}^1) - a_2(\Delta_{k', \ell'})| > 18. \quad (7)$$

On the other hand, by some detailed analysis on the L -values $L(g, K-1)$ and $L(g, K-2)$ for $g \in \mathcal{H}_K$, we obtain (Proposition 2.6) for $K \geq 100$

$$|a_2(\Delta_{k, \ell}^1) - a_2(\Delta_{k', \ell'})| < 16.007. \quad (8)$$

These arguments enable us to finish the proof for $K \geq 100$. The case $K < 100$ has been verified numerically by Daozhou Zhu; see [10]. Altogether, the proof of Theorem 1.1 is complete.

2. The proofs

We shall first establish a lower bound for $|a_2(\Delta_{k, \ell}^1) - a_2(\Delta_{k', \ell'})|$. In order to do this we first recall the following results on the a_2 -coefficients of $\Delta_{k, \ell}^1$ and $\Delta_{k', \ell'}$ for large K .

Proposition 2.1. *We have*

$$\lim_{k \rightarrow \infty} \frac{a_2(\Delta_{k, \ell}^1)}{2(1 + 2^{\ell-1})} = 1, \quad \text{and} \quad \lim_{k' \rightarrow \infty} \frac{a_2(\Delta_{k', \ell'})}{1 + 2^{\ell'-1}} = 1.$$

Moreover, when $K \geq 100$, we have

$$a_2(\Delta_{k, \ell}^1) = (2 + 2^\ell)(1 + \delta^1), \quad \text{and} \quad a_2(\Delta_{k', \ell'}) = (1 + 2^{\ell'-1})(1 + \delta),$$

where $|\delta^1| < 0.11294$, and $|\delta| < 0.21703$.

Proof. The bound for $|\delta|$ is obtained by plugging $K = 100$ in the calculations in [5, Lemma 3.8]. The bound for $|\delta^1|$ is obtained by plugging $K = 100$ into the calculations in [4, Proposition 3.4]. \square

Lemma 2.2. *Let $K \geq 100$. If $\ell \geq 6$ or $\ell' \geq 8$ are even, then*

$$|a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'})| > 18.$$

Proof. Let us first assume that $\ell' \geq 8$. Then there are two cases to consider.

Case 1: $\ell \geq \ell'$. By Proposition 2.1

$$\begin{aligned} |a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'})| &= |(2 + 2^\ell)(1 + \delta^1) - (1 + 2^{\ell'-1})(1 + \delta)| \\ &= (2 + 2^\ell) \left| (1 + \delta^1) - \frac{(1 + 2^{\ell'-1})}{2(1 + 2^{\ell-1})}(1 + \delta) \right| \\ &> (2 + 2^\ell)(1 - 0.11294 - 0.5(1 + 0.21763)) \\ &> (2 + 2^8) \cdot 0.27825 \\ &> 71. \end{aligned}$$

Case 2: $\ell < \ell'$. Note that $(2 + 2^\ell)/(1 + 2^{\ell'-1})$ maximizes at $\ell = 6$ and $\ell' = 8$. Thus

$$\begin{aligned} |a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'})| &= |(2 + 2^\ell)(1 + \delta^1) - (1 + 2^{\ell'-1})(1 + \delta)| \\ &= (1 + 2^{\ell'-1}) \left| \frac{2 + 2^\ell}{1 + 2^{\ell'-1}}(1 + \delta^1) - (1 + \delta) \right| \\ &> (1 + 2^{\ell'-1}) \left(1 - |\delta| - \frac{2 + 2^6}{1 + 2^7}(1 + |\delta^1|) \right) \\ &> (1 + 2^{\ell'-1}) \cdot \left(1 - 0.21763 - \frac{66}{129}(1 + 0.11294) \right) \\ &> 27. \end{aligned}$$

Now, assume that $\ell \geq 6$. Similarly, when $\ell \geq \ell'$, we have

$$\begin{aligned} |a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'})| &> (2 + 2^\ell)(1 - 0.11294 - 0.5(1 + 0.21763)) \\ &> (2 + 2^6) \cdot 0.27825 \\ &> 18. \end{aligned}$$

When $\ell < \ell'$, then $\ell' \geq 8$ and we have

$$\begin{aligned} |a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'})| &> (1 + 2^{\ell'-1}) \cdot \left(1 - 0.21763 - \frac{66}{129}(1 + 0.11294) \right) \\ &> 27. \end{aligned}$$

The proof is now complete. \square

Remark 2.3. If neither condition of Lemma 2.2 is met, for instance if $\ell' = 6$ and $\ell = 4$, then

$$|2 + 2^4 - 1 - 2^5| = 15 < 16,$$

which does not contradict the upper bound obtained in Proposition 2.6. Therefore, it seems that Lemma 2.2 is optimal.

Next, we shall establish some estimates on the values $L(g, K-1)$ and $L(g, K-2)$ for each Hecke eigenform $g \in \mathcal{H}_K$ and for $K \geq 100$.

By Deligne's bound $|a_n(g)| \leq d(n)n^{(K-1)/2}$ with $d(n)$ being the number of divisor function and the fact that $\sum_{n=1}^{\infty} \frac{d(n)}{n^s} = \zeta(s)^2$, we get

$$\left| L(g, K-1) - 1 - \frac{a_2(g)}{2^{K-1}} \right| \leq \sum_{n=3}^{\infty} \frac{d(n)}{n^{(K-1)/2}} = \zeta\left(\frac{K-1}{2}\right)^2 - 1 - 2^{1-(K-1)/2}.$$

Noting that

$$\begin{aligned} \zeta\left(\frac{K-1}{2}\right) &= 1 + 2^{-(K-1)/2} + 3^{-(K-1)/2} + \sum_{n=4}^{\infty} n^{-(K-1)/2} \\ &\leq 1 + 2^{-(K-1)/2} + 3^{-(K-1)/2} + \int_3^{\infty} t^{-(K-1)/2} dt \\ &= 1 + 2^{-(K-1)/2} + 3^{-(K-1)/2} + \frac{6}{K-3} \cdot 3^{-(K-1)/2} \\ &= 1 + 2^{-(K-1)/2} + \frac{K+3}{K-3} 3^{-(K-1)/2}. \end{aligned}$$

This means that

$$\begin{aligned} &\zeta\left(\frac{K-1}{2}\right)^2 - 1 - 2^{1-(K-1)/2} \\ &= 2 \cdot 3^{-(K-1)/2} \frac{K+3}{K-3} + 2^{-(K-1)} + 2^{1-(K-1)/2} 3^{-(K-1)/2} \frac{K+3}{K-3} \\ &\quad + \left(\frac{K+3}{K-3}\right)^2 3^{-(K-1)}. \end{aligned}$$

Thus, for $K \geq 100$, we may write

$$L(g, K-1) = 1 + \frac{a_2(g)}{2^{K-1}} + \delta_{K-1} \cdot 3^{-(K-1)/2}, \quad (9)$$

where

$$\begin{aligned} &|\delta_{K-1}| \\ &\leq 2 \frac{K+3}{K-3} + \left(\frac{3}{4}\right)^{(K-1)/2} + 2^{1-(K-1)/2} \frac{K+3}{K-3} + \left(\frac{K+3}{K-3}\right)^2 3^{-(K-1)/2} \\ &< 2.12372. \end{aligned} \quad (10)$$

Similarly, for $L(g, K-2)$ we have

$$\left| L(g, K-2) - 1 - \frac{a_2(g)}{2^{K-2}} \right| \leq \sum_{n=3}^{\infty} \frac{d(n)}{n^{(K-3)/2}} = \zeta\left(\frac{K-3}{2}\right)^2 - 1 - 2^{1-(K-3)/2},$$

and for $K \geq 100$ we obtain

$$L(g, K - 2) = 1 + \frac{a_2(g)}{2^{K-2}} + \delta_{K-2} \cdot 3^{-(K-3)/2}, \quad (11)$$

where

$$\begin{aligned} & |\delta_{K-2}| \quad (12) \\ & \leq 2 \frac{K+1}{K-5} + \left(\frac{3}{4}\right)^{(K-3)/2} + 2^{1-(K-3)/2} \frac{K+1}{K-5} + \left(\frac{K+1}{K-5}\right)^2 3^{-(K-3)/2} \\ & < 2.12632. \end{aligned}$$

From now on, let us assume on the contrary that the odd period $r_{\ell'-1}$ and the even period r_ℓ are linearly dependent. By (1) and the Eichler-Shimura relations $r_\ell + r_{K-2-\ell} = 0$ and $r_{\ell'-1} - r_{K-1-\ell'} = 0$, this means that there is some constant c such that for all $g \in \mathcal{H}_K$, where $K = k + \ell + 2 = k' + \ell'$, such that

$$L(g, k') = c \cdot L(g, k + 1). \quad (13)$$

Our strategy is to derive a contradiction from (13) on the value

$$|a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'})|.$$

A lower bound of it has been established in Lemma 2.2. Our next task is to find an upper bound. We shall first derive some information from the a_1 -coefficients. By Proposition 1.3 and (11), we obtain

$$\begin{aligned} 1 &= A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, K-2)L(g, k+1)}{\langle g, g \rangle} \\ &= A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)(1 + a_2(g)2^{2-K} + \delta_{K-2} \cdot 3^{-(K-3)/2})}{\langle g, g \rangle}. \end{aligned}$$

As $|a_2(g)2^{2-K}| \leq 2 \cdot 2^{(3-K)/2}$, by (12), for $K \geq 100$

$$|a_2(g)2^{2-K}| + |\delta_{K-2} \cdot 3^{-(K-3)/2}| < 2.0001 \cdot 2^{(3-K)/2}.$$

Since $L(g, k + 1)$ is positive for all g (Remark 1.4), for $K \geq 100$

$$\begin{aligned} 1 &= \left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)(1 + a_2(g)2^{2-K} + \delta_{K-2} \cdot 3^{-(K-3)/2})}{\langle g, g \rangle} \right| \\ &\geq |A_{k,\ell}^1| \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)(1 - |a_2(g)2^{2-K}| - |\delta_{K-2} \cdot 3^{-(K-3)/2}|)}{\langle g, g \rangle} \\ &\geq \left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} \right| \cdot (1 - 2.0001 \cdot 2^{(3-K)/2}) \end{aligned}$$

and

$$\begin{aligned}
1 &= \left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)(1 + a_2(g)2^{2-K} + \delta_{K-2} \cdot 3^{-(K-3)/2})}{\langle g, g \rangle} \right| \\
&\leq |A_{k,\ell}^1| \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)(1 + |a_2(g)2^{2-K}| + |\delta_{K-2} \cdot 3^{-(K-3)/2}|)}{\langle g, g \rangle} \\
&\leq \left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} \right| \cdot (1 + 2.0001 \cdot 2^{(3-K)/2}).
\end{aligned}$$

Thus, for $K \geq 100$

$$A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} = 1 + \epsilon_K^1(1), \quad (14)$$

for some $|\epsilon_K^1(1)| < 2.001 \cdot 2^{(3-K)/2}$.

Similarly, by Proposition 1.3, (11) and taking (13) into account, we get

$$\begin{aligned}
1 &= A_{k',\ell'} \sum_{g \in \mathcal{H}_K} \frac{L(g, K-1)L(g, k')}{\langle g, g \rangle} \\
&= cA_{k',\ell'} \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)(1 + a_2(g)2^{1-K} + \delta_{K-1} \cdot 3^{-(K-1)/2})}{\langle g, g \rangle}.
\end{aligned}$$

For $K \geq 100$, as

$$|a_2(g)2^{1-K}| \leq 2 \cdot 2^{(1-K)/2}$$

and

$$|a_2(g)2^{1-K} + \delta_{K-1} \cdot 3^{-(K-1)/2}| < 2.0001 \cdot 2^{(1-K)/2},$$

we obtain analogously

$$cA_{k',\ell'} \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} = 1 + \epsilon_K(1) \quad (15)$$

for some $|\epsilon_K(1)| \leq 2.001 \cdot 2^{(1-K)/2}$.

We next investigate and compare the a_2 -coefficients of $\Delta_{k,\ell}^1$ and $\Delta_{k,\ell}$. Again, by Proposition 1.3 and (11)

$$\begin{aligned}
a_2(\Delta_{k,\ell}^1) &= A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, K-2)L(g, k+1)}{\langle g, g \rangle} a_2(g) \\
&= A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2(g) + A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2^2(g)2^{2-K} \\
&\quad + A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} \cdot \delta_{K-2} \cdot 3^{-(K-3)/2} a_2(g).
\end{aligned} \quad (16)$$

We denote the last term of (16) by $\epsilon_K^1(2)$. Then, for $K \geq 100$, by (12)

$$\begin{aligned} |\epsilon_K^1(2)| &= \left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} \cdot \delta_{K-2} \cdot 3^{-(K-3)/2} a_2(g) \right| \\ &\leq \left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} \right| \cdot |\delta_{K-2}| \cdot 2 \cdot \left(\frac{2}{3}\right)^{(K-3)/2} \\ &< 0.0001. \end{aligned}$$

Here we have again used the fact that $L(g, k+1)$ is positive, see Remark 1.4. Analogously, for $K \geq 100$ we have

$$\begin{aligned} &a_2(\Delta_{k',\ell'}) \tag{17} \\ &= cA_{k',\ell'} \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2(g) + cA_{k',\ell'} \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2^2(g) 2^{1-K} \\ &\quad + \epsilon_K(2), \end{aligned}$$

where

$$|\epsilon_K(2)| < 0.0001. \tag{18}$$

Lemma 2.4. *Let $k > \ell \geq 2$ and $k' \geq \ell' \geq 2$ be even integers such that $K := k + \ell + 2 = k' + \ell'$. Assume that (13): $L(g, k') = cL(g, k+1)$ holds for all $g \in \mathcal{H}_K$. Then for $K \geq 100$*

$$\left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2^2(g) 2^{1-K} \right| \leq 4(1 + |\epsilon_K^1(1)|).$$

Proof. This is due to the Deligne's bound $|a_2(g)| \leq 2 \cdot 2^{(K-1)/2}$, (14) and positivity of $L(g, k+1)$. \square

Lemma 2.5. *Retain the assumptions of Lemma 2.4. Then for $K \geq 100$*

$$\left| (A_{k,\ell}^1 - cA_{k',\ell'}) \left(\sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2(g) \right) \right| < 12.006,$$

and

$$\left| A_{k,\ell}^1 - cA_{k',\ell'} \right| \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2^2(g) 2^{1-K} \leq 4(|\epsilon_K^1(1)| + |\epsilon_K(1)|).$$

Proof. By (14) and (15), we get

$$\left| (A_{k,\ell}^1 - cA_{k',\ell'}) \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} \right| \leq |\epsilon_K^1(1)| + |\epsilon_K(1)|.$$

Remembering that $|a_2(g)| \leq 2 \cdot 2^{(K-1)/2}$, $|\epsilon_K^1(1)| \leq 4.002 \cdot 2^{(1-K)/2}$, $|\epsilon_K(1)| \leq 2.001 \cdot 2^{(1-K)/2}$, and that $L(g, k+1) > 0$, thus

$$\begin{aligned} & \left| (A_{k,\ell}^1 - cA_{k',\ell'}^1) \left(\sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2(g) \right) \right| \\ & \leq 2 \cdot 2^{(K-1)/2} \left| (A_{k,\ell}^1 - cA_{k',\ell'}^1) \sum_{g \in \mathcal{X}} \frac{L(g, k+1)}{\langle g, g \rangle} \right| \\ & \leq 2 \cdot 2^{(K-1)/2} \cdot (|\epsilon_K^1(1)| + |\epsilon_K(1)|) \\ & \leq 12.006, \end{aligned}$$

giving the first inequality. The second inequality follows exactly the same way by noticing that $|a_2(g)|^2 2^{1-K} \leq 4$. \square

We are now ready to establish an upper bound for $|\Delta_{k,\ell}^1 - \Delta_{k',\ell'}^1|$. Note that this result does not require that ℓ or ℓ' satisfy the conditions in Lemma 2.2.

Proposition 2.6. *Let $k > \ell \geq 2$ and $k' \geq \ell' \geq 2$ be even integers such that $K := k + \ell + 2 = k' + \ell'$. Assume that (13): $L(g, k') = cL(g, k+1)$ holds for all $g \in \mathcal{H}_K$. Then, for $K \geq 100$*

$$|a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'}^1)| < 16.007.$$

Proof. By (16) and (17)

$$\begin{aligned} & |a_2(\Delta_{k,\ell}^1) - a_2(\Delta_{k',\ell'}^1)| \\ & \leq \left| A_{k,\ell}^1 - cA_{k,\ell}^1 \right| \left| \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2(g) \right| + \left| A_{k,\ell}^1 - cA_{k',\ell'}^1 \right| \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{2^{K-1} \langle g, g \rangle} a_2^2(g) \\ & \quad + \left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2^2(g) 2^{1-K} \right| + |\epsilon_K^1(2)| + |\epsilon_K(2)|. \end{aligned}$$

Now, by Lemma 2.4

$$\left| A_{k,\ell}^1 \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2^2(g) 2^{1-K} \right| \leq 4(1 + |\epsilon_K^1(1)|).$$

By Lemma 2.5

$$\left| (A_{k,\ell}^1 - cA_{k',\ell'}^1) \left(\sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2(g) \right) \right| < 12.006,$$

and hence

$$\left| A_{k,\ell}^1 - cA_{k',\ell'}^1 \right| \sum_{g \in \mathcal{H}_K} \frac{L(g, k+1)}{\langle g, g \rangle} a_2^2(g) 2^{1-K} \leq 4(|\epsilon_K^1(1)| + |\epsilon_K(1)|).$$

The desired upper bound then follows by taking into consideration of the bounds $|\epsilon_K^1(1)| < 2.001 \cdot 2^{(3-K)/2}$, $|\epsilon_K(1)| < 2.001 \cdot 2^{(1-K)/2}$, $|\epsilon_K^1(2)|$, $|\epsilon_K(2)| < 0.0001$. \square

As having been explained in the introduction, the lower bound Lemma 2.2 and the upper bound Proposition 2.6 imply that Theorem 1.1 holds true for $K \geq 100$. The remaining finitely many cases have been numerically verified by Daozhou Zhu ([10]).

Acknowledgment

The author is grateful to the referee for careful reading of the paper and valuable suggestions and comments.

References

- [1] FUKUHARA, SHINJI. Explicit formulas for Hecke operators on cusp forms, Dedekind symbols and period polynomials. *J. Reine Angew. Math.* **607** (2007), 163–216. MR2338123, Zbl 1137.11029, arXiv:math/0506373, doi: 10.1515/CRELLE.2007.048. 521
- [2] FUKUHARA, SHINJI. A basis for the space of modular forms. *Acta Arith.* **151** (2012), no. 4, 421–427. MR2861775, Zbl 1260.11031, arXiv:1008.4008, doi: 10.4064/aa151-4-5. 521
- [3] KOHNEN, WINFRIED; ZAGIER, DON. Modular forms with rational periods. *Modular forms* (Durham, 1983), 197–249. Ellis Horwood Ser. Math. Appl.: Statist. Oper. Res. *Ellis Horwood Ltd., Chichester*, 1984. ISBN: 0-85312-669-0. MR0803368, Zbl 0618.10019. 523
- [4] LEI, AUSTIN; NI, TIANYU; XUE, HUI. Linear independence of even periods of modular forms. *J. Number Theory* **248** (2023), 120–139. MR4556159, Zbl 07672968, doi: 10.1016/j.jnt.2023.01.004. 522, 523, 524, 525
- [5] LEI, AUSTIN; NI, TIANYU; XUE, HUI. Linear independence of odd periods of modular forms. *Res. Number Theory* **9** (2023), no. 2, Paper No. 33, 20 pp. MR4578514, Zbl 07691096, doi: 10.1007/s40993-023-00439-9. 522, 524, 525
- [6] MANIN, JU. I. Periods of cusp forms, and p -adic Hecke series. Lecture Notes in Mathematics, 1374. *Mat. Sb. (N.S.)* **92(134)** (1973), 378–401, 503. MR0345909. Zbl 0293.14007, 521
- [7] ZAGIER, DON. Modular forms whose Fourier coefficients involve zeta-functions of quadratic fields. *Modular functions of one variable, VI* (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), 105–169. Lecture Notes in Math., 627. *Springer-Verlag, Berlin-New York*, 1977. ISBN: 3-540-08530-0. MR0485703, Zbl 0372.10017. 523
- [8] ZAGIER, DON. Modular forms and differential operators. *Proc. Indian Acad. Sci. Math. Sci.* **104** (1994), no. 1, 57–75. MR1280058, Zbl 0806.11022, doi: 10.1007/BF02830874. 522
- [9] ZAGIER, DON. Elliptic modular forms and their applications. *The 1-2-3 of modular forms*, 1–103. Universitext. *Springer, Berlin*, 2008. ISBN: 978-3-540-74117-6. MR2409678, Zbl 1259.11042. doi: 10.1007/978-3-540-74119-0_1. 522
- [10] ZHU, DAOZHOU. Code for independence between odd and even periods. <https://github.com/hxue-clemson/Independence-between-even-and-odd-periods>. 522, 524, 531

(Hui Xue) SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, CLEMSON UNIVERSITY, CLEMSON, SC 29634-0975, USA
huixue@clemson.edu

This paper is available via <http://nyjm.albany.edu/j/2024/30-22.html>.