

\mathcal{PT} Symmetric Schrödinger Operators: Reality of the Perturbed Eigenvalues*

Emanuela CALICETI[†], Francesco CANNATA[‡] and Sandro GRAFFI[†]

[†] *Dipartimento di Matematica, Università di Bologna, and INFN, Bologna, Italy*

E-mail: caliceti@dm.unibo.it, graffi@dm.unibo.it

[‡] *INFN, Via Irnerio 46, 40126 Bologna, Italy*

E-mail: Francesco.Cannata@bo.infn.it

Received November 03, 2009, in final form January 14, 2010; Published online January 20, 2010

doi:[10.3842/SIGMA.2010.009](https://doi.org/10.3842/SIGMA.2010.009)

Abstract. We prove the reality of the perturbed eigenvalues of some \mathcal{PT} symmetric Hamiltonians of physical interest by means of stability methods. In particular we study 2-dimensional generalized harmonic oscillators with polynomial perturbation and the one-dimensional $x^2(ix)^\epsilon$ for $-1 < \epsilon < 0$.

Key words: \mathcal{PT} symmetry; real spectra; perturbation theory

2010 Mathematics Subject Classification: 47A55; 47A75; 81Q15; 34L40; 35J10

1 Introduction and statement of the results

A basic mathematical question in the framework of \mathcal{PT} symmetric quantum mechanics concerns the reality of the spectrum of the considered Hamiltonian. One crucial issue is to formulate conditions for the reality of (part of) the spectrum of Hamiltonians depending on a real parameter ϵ of the form

$$H(\epsilon) = H_0 + \epsilon V, \quad \epsilon \in \mathbb{R}. \quad (1)$$

In [1, 2] results have been obtained in the case when the unperturbed Hamiltonian H_0 is self-adjoint with discrete spectrum, and in [3] in the more general case when H_0 is \mathcal{PT} symmetric but not necessarily selfadjoint, with discrete spectrum. The common framework of these papers is perturbation theory, which allows to obtain the reality result only for small values of the coupling constant ϵ . However this is not a limitation: in many cases as ϵ varies, critical phenomena occur, such as a spontaneous breaking of the \mathcal{PT} symmetry and the appearance of complex eigenvalues caused by the crossing of energy levels of H_0 . The most elementary example of this phenomenon is represented by a 2×2 matrix of the form

$$H_1(\epsilon) = \begin{pmatrix} e_1 & i\epsilon \\ i\epsilon & e_2 \end{pmatrix} \quad (2)$$

with $e_1, e_2 \in \mathbb{R}$, recently examined in [4, 5]. $H_1(\epsilon)$ is \mathcal{PT} symmetric, i.e. $H_1(\epsilon)(\mathcal{PT}) = (\mathcal{PT})H_1(\epsilon)$, if the parity operator \mathcal{P} is the unitary involution defined by

$$\mathcal{P} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

*This paper is a contribution to the Proceedings of the 5-th Microconference “Analytic and Algebraic Methods V”. The full collection is available at <http://www.emis.de/journals/SIGMA/Prague2009.html>

and \mathcal{T} is the antilinear involution of complex conjugation. The eigenvalues of $H_1(\epsilon)$ are

$$\lambda(\epsilon) = \frac{1}{2}(e_1 + e_2) \pm \frac{1}{2}\sqrt{(e_1 - e_2)^2 - 4\epsilon^2}$$

and they are real if and only if $|\epsilon| \leq \frac{|e_1 - e_2|}{2}$. The matrix (2) can be rewritten in the perturbative form (1)

$$H_1(\epsilon) = \begin{pmatrix} e_1 & 0 \\ 0 & e_2 \end{pmatrix} + i\epsilon \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

so that it can be put in the framework of Theorem 1.2 of [1], whose applicability goes far beyond the case of finite dimensional operators (it includes in fact classes of Schrödinger operators with bounded perturbation potentials, also in dimension greater than one): this theorem ensures the reality of the spectrum of $H_1(\epsilon)$ for $|\epsilon| < \frac{|e_1 - e_2|}{2\|W\|}$ where $\|W\|$ denotes the norm of the bounded perturbation operator W , which in the present case is 1 since $W = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. The conclusion is that the perturbation argument of [1] yields actually the reality of the spectrum for all the allowed values of ϵ .

In a similar way one can analyze \mathcal{PT} symmetric matrices of the form

$$H'_1(\epsilon) = \begin{pmatrix} e + i\epsilon & b \\ b & e - i\epsilon \end{pmatrix} = \begin{pmatrix} e & b \\ b & e \end{pmatrix} + i\epsilon \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (3)$$

with $e, b \in \mathbb{R}$, which are \mathcal{PT} symmetric with

$$\mathcal{P} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

For a physical interpretation of these matrices see [4, 5] for applications to optics, and Section 4 of the present paper for an application to classical mechanics. Complex Hamiltonians of type (2) are a particular case of those considered in [6]. In turn the Hamiltonians (3) are a particular case of those examined in [7].

The results obtained in [1, 2] deal with the case when the perturbation potential V is bounded and, under suitable assumptions on H_0 and V , they guarantee the reality of the entire spectrum of $H(\epsilon)$ for $|\epsilon| < \delta/\|W\|$, where $\delta = \frac{1}{2} \inf_{n \neq m} |\lambda_n - \lambda_m|$ and $\sigma(H_0) = \{\lambda_n : n \in \mathbb{N}\}$ is the (discrete) spectrum of H_0 . In some cases, not only the reality of the eigenvalues can be proved, but also the similarity to a selfadjoint operator [8, 9]. In [3] the authors analyze the case when V is unbounded (and in general not even relatively bounded with respect to H_0), obtaining a weaker result which guarantees that the perturbed eigenvalues of $H(\epsilon)$, converging to those of H_0 as $\epsilon \rightarrow 0$, are real for $|\epsilon|$ sufficiently small. In [10] this result has been extended to classes of Hamiltonians of the form

$$H(\epsilon) = H + iW_\epsilon, \quad \epsilon \geq 0$$

acting in $L^2(\mathbb{R})$, where $H = -\frac{d^2}{dx^2} + x^2$ denotes the operator associated with the one dimensional harmonic oscillator and $W_\epsilon \in C^0(\mathbb{R})$ is an odd real-valued function: $W_\epsilon(-x) = -W_\epsilon(x) \in \mathbb{R}$, $\forall x \in \mathbb{R}$, $\forall \epsilon \geq 0$. Such generalization of the result of [3] is based on the fact that in W_ϵ the dependence on the perturbation parameter ϵ is the most general one and not only of linear type as in (1). The aim of this paper is to show how simple extensions of the results of [3, 10] allow us to treat models of considerable interest in the context of \mathcal{PT} symmetric quantum mechanics. More precisely, extending the results of [3] to the case of Schrödinger operators in dimension greater than 1 it is possible to examine models of the following type

$$H_2(\epsilon) = -\frac{d^2}{dx_1^2} - \frac{d^2}{dx_2^2} + \omega_1^2 x_1^2 + \omega_2^2 x_2^2 + i\epsilon x_1^r x_2^s, \quad x = (x_1, x_2) \in \mathbb{R}^2, \quad (4)$$

where we assume $r, s \in \mathbb{N}$, $r + s$ odd, and $\omega_1, \omega_2 > 0$. This type of Hamiltonians, which represent a natural generalization of the quantum Henon–Heiles model, has been studied also in [11, 12]. We will analyze the Hamiltonian (4) in Section 2 and we will prove the reality of the eigenvalues $\lambda_{n_1, n_2}(\epsilon)$ of $H_2(\epsilon)$ generated by the unperturbed ones, i.e. by the eigenvalues $\lambda_{n_1, n_2}(0) = (2n_1 + 1)\omega_1 + (2n_2 + 1)\omega_2$, $\forall n_1, n_2 \in \mathbb{N}_0 := \{0, 1, 2, \dots\}$, such that $\lambda_{n_1, n_2}(\epsilon) \rightarrow \lambda_{n_1, n_2}(0)$ as $\epsilon \rightarrow 0$, in the case of non-resonant frequencies ω_1, ω_2 .

Then, generalizing also the results of [10] we will examine the basic model

$$H_3(\epsilon) = -\frac{d^2}{dx^2} + x^2(ix)^\epsilon, \quad -1 < \epsilon < 1. \tag{5}$$

The Hamiltonian (5) has been examined and its physical interest discussed by several authors [13, 14, 15, 16] and the reality of its spectrum for $\epsilon > 0$ has been proved in [17] (see also [18]); for $\epsilon < 0$ numerical results indicate the appearance of complex eigenvalues (for a recent discussion see e.g. [14]), in the sense that, as $|\epsilon|$ increases while ϵ moves from 0 to -1 , the perturbed eigenvalues of $H_3(\epsilon)$, generated by the unperturbed ones $\lambda_n(0) = 2n + 1$, $\forall n \in \mathbb{N}_0$, become complex. In Section 3 we will prove that as long as $|\epsilon|$ stays sufficiently small, those eigenvalues remain real. We will actually treat the case of small $|\epsilon|$ independently of the sign of ϵ : for $\epsilon > 0$ we recover the known result proved in [17].

Remark 1. As in [3, 10], the results of Sections 2 and 3 are obtained by proving the stability of the unperturbed eigenvalues. For the notion of stability see e.g. [19, 20]. In particular, let us recall that if $\{H(\epsilon) : |\epsilon| < \epsilon_0\}$ is a family of closed \mathcal{PT} symmetric operators in a Hilbert space \mathcal{H} and if a simple (i.e., non degenerate) eigenvalue λ of $H(0)$ is stable with respect to the family $\{H(\epsilon) : |\epsilon| < \epsilon_0\}$, then for $|\epsilon|$ sufficiently small there exists one and only one eigenvalue $\lambda(\epsilon)$ of $H(\epsilon)$ near λ such that

$$\lambda(\epsilon) \rightarrow \lambda, \quad \text{as } \epsilon \rightarrow 0.$$

Now, recalling that the eigenvalues of a \mathcal{PT} symmetric operator come in pairs of complex conjugate values, we can assert that the uniqueness of $\lambda(\epsilon)$ implies its reality: in fact, if $\lambda(\epsilon)$ is not real then there are two distinct eigenvalues, $\lambda(\epsilon)$ and $\overline{\lambda(\epsilon)}$, and not just one as stated above.

Assume now that $\mathcal{H} = L^2(\mathbb{R}^d)$, $d \geq 1$, and that $H(\epsilon)$, $|\epsilon| < \epsilon_0$, is formally given by

$$H(\epsilon) = p^2 + V_1 + W_\epsilon,$$

where $p^2 = -\Delta = -\sum_{k=1}^d \frac{d^2}{dx_k^2}$, $x = (x_1, \dots, x_k) \in \mathbb{R}^d$ and $V_1, W_\epsilon \in L^2_{\text{loc}}(\mathbb{R}^d)$. Moreover we assume that $H(\epsilon)$ has discrete spectrum, i.e. each point in the spectrum is an isolated eigenvalue with finite multiplicity (for a review of the notion of multiplicity of an eigenvalue see also [3]). Let $H^*(\epsilon)$ denote the adjoint of $H(\epsilon)$ and assume that both $H(\epsilon)$ and $H^*(\epsilon)$ have $C_0^\infty(\mathbb{R}^d)$ as a core.

As recalled in [10] the stability of the eigenvalues of $H(0)$ in the framework of the stability theory developed by Hunziker and Vock in [21] holds if the following three conditions are satisfied.

- (1) For all $u \in C_0^\infty(\mathbb{R}^d)$

$$\lim_{\epsilon \rightarrow 0} \|H(\epsilon)u - H(0)u\| = 0 \tag{6}$$

and

$$\lim_{\epsilon \rightarrow 0} \|H^*(\epsilon)u - H^*(0)u\| = 0. \tag{7}$$

(2) *There exist constants $a, b > 0$ and γ , $|\gamma| < \pi/2$, such that*

$$\langle u, p^2 u \rangle \leq a \{ \cos \gamma \Re \langle u, H(\epsilon) u \rangle + \sin \gamma \Im \langle u, H(\epsilon) u \rangle + b \langle u, u \rangle \} \quad (8)$$

for all $u \in C_0^\infty(\mathbb{R}^d)$.

(3) *For any $z \in \mathbb{C}$, there exist positive constants δ , n_0 and ϵ_1 such that*

$$d_n(z, H(\epsilon)) \geq \delta > 0, \quad (9)$$

for all $n > n_0$ and $|\epsilon| < \epsilon_1$, where $d_n(z, H(\epsilon)) := \text{dist}(z, \mathcal{N}_n(\epsilon))$ and

$$\mathcal{N}_n(\epsilon) := \{ \langle u, H(\epsilon) u \rangle : u \in D(H(\epsilon)), \|u\| = 1, u(x) = 0 \text{ for } |x| < n \}$$

is the so called “numerical range at infinity” (see [3]).

Here $D(H(\epsilon))$ denotes the domain of $H(\epsilon)$ and $\langle u, v \rangle := \int_{\mathbb{R}^d} u(x) \overline{v(x)} dx$ denotes the standard scalar product in $L^2(\mathbb{R}^d)$. For future reference we summarize this result in the following criterion, proved in [21] (see also [10]).

Proposition 1. *The eigenvalues of $H(0)$ are stable w.r.t. the family $H(\epsilon)$, $|\epsilon| < \epsilon_0$, if the above conditions (1)–(3) are satisfied.*

In Sections 2 and 3 we will prove that the operator families $H_2(\epsilon)$ and $H_3(\epsilon)$ satisfy this criterion and, in view of Remark 1 above, this is enough to ensure the reality of the perturbed eigenvalues of $H_2(\epsilon)$ and $H_3(\epsilon)$.

2 The operator family $H_2(\epsilon)$: stability and reality of the perturbed eigenvalues

The Schrödinger operator $H_2(\epsilon)$, $\epsilon \in \mathbb{R}$, introduced in Section 1 is defined as the closure in $L^2(\mathbb{R}^2)$ of the minimal operator defined on $C_0^\infty(\mathbb{R}^2)$, we denote its domain $D(H_2(\epsilon))$, and its adjoint is given by $H_2(-\epsilon)$. It has discrete spectrum (see e.g. [20], Theorem XIII.64; we note that the extension to the present non-selfadjoint case is straightforward) and the unperturbed eigenvalues, i.e. the eigenvalues of $H_2(0)$, are given by

$$\lambda_{n_1, n_2}(0) = (2n_1 + 1)\omega_1 + (2n_2 + 1)\omega_2, \quad \forall n_1, n_2 \in \mathbb{N}_0.$$

We assume that the frequencies $\omega_1, \omega_2 > 0$ are non-resonant, i.e. the equation $k_1\omega_1 + k_2\omega_2 = 0$, $k_1, k_2 \in \mathbb{Z}$, is satisfied if and only if $k_1, k_2 = 0$. Then each eigenvalue $\lambda_{n_1, n_2}(0)$ is simple, i.e. the corresponding eigenspace has dimension 1. Moreover we assume that $r, s \in \mathbb{N}$ are not both even. Then $H_2(\epsilon)$ is \mathcal{PT} symmetric, where \mathcal{T} is complex conjugation and the parity operator \mathcal{P} is defined as follows: if r and s are both odd \mathcal{P} changes the sign of one coordinate, i.e. it can be either $(\mathcal{P}_1 u)(x_1, x_2) = u(-x_1, x_2)$ or $(\mathcal{P}_2 u)(x_1, x_2) = u(x_1, -x_2)$, $\forall x = (x_1, x_2) \in \mathbb{R}^2$, $\forall u \in L^2(\mathbb{R}^2)$. If r and s are not both odd a suitable choice for \mathcal{P} is $(\mathcal{P}_3 u)(x) = u(-x)$, $\forall x \in \mathbb{R}^2$, $\forall u \in L^2(\mathbb{R}^2)$.

We now prove that conditions (1)–(3) are satisfied.

Proposition 2. *Conditions (1)–(3) are satisfied by the operator family $H_2(\epsilon)$, $\epsilon \in \mathbb{R}$.*

Proof. (1) $\forall u \in C_0^\infty(\mathbb{R}^2)$ let $K = \text{supp}(u)$ denote the support of u ; since K is compact in \mathbb{R}^2 , there exists a constant $M > 0$ such that $|x_1^r x_2^s| \leq M$, $\forall x = (x_1, x_2) \in K$. Therefore

$$\begin{aligned} \|H_2(\epsilon)u - H_2(0)u\|^2 &= \|H_2^*(\epsilon)u - H_2^*(0)u\|^2 = \int_K |i\epsilon x_1^r x_2^s u(x)|^2 dx \\ &\leq \epsilon^2 M^2 \int_K |u(x)|^2 dx = \epsilon^2 M^2 \|u\|^2, \quad \forall \epsilon \in \mathbb{R}, \end{aligned}$$

and this yields (6) and (7).

(2) $\forall u \in C_0^\infty(\mathbb{R}^2)$ we have

$$\Re\langle u, H_2(\epsilon)u \rangle = \langle u, p^2u \rangle + \omega_1^2\langle u, x_1^2u \rangle + \omega_2^2\langle u, x_2^2u \rangle \geq \langle u, p^2u \rangle.$$

Thus, (8) is satisfied with $\gamma = 0$, $a = 1$, $b = 0$.

(3) Let $z \in \mathbb{C}$ and $\omega := \min\{\omega_1, \omega_2\}$. Let $u \in D(H_2(\epsilon))$ be such that $\|u\| = 1$ and $u(x) = 0$ for $|x| < n$, $n \in \mathbb{N}$. Then

$$\begin{aligned} |z - \langle u, H_2(\epsilon)u \rangle| &\geq |\langle u, H_2(\epsilon)u \rangle| - |z| \geq \Re\langle u, H_2(\epsilon)u \rangle - |z| \\ &\geq \omega_1^2\langle u, x_1^2u \rangle + \omega_2^2\langle u, x_2^2u \rangle - |z| \geq \omega^2\langle u, |x|^2u \rangle - |z| \geq \omega^2n^2 - |z|. \end{aligned}$$

Thus, $\text{dist}(z, \mathcal{N}_n(\epsilon)) \geq \omega^2n - |z|$, $\forall \epsilon \in \mathbb{R}$, $\forall n \in \mathbb{N}$, whence $\lim_{n \rightarrow \infty} d_n(z, H(\epsilon)) = +\infty$, $\forall \epsilon \in \mathbb{R}$ and this proves (9). ■

Corollary 1. *Near each unperturbed eigenvalue $\lambda_{n_1, n_2}(0)$, $n_1, n_2 \in \mathbb{N}_0$, of $H_2(0)$ there exists one and only one eigenvalue $\lambda_{n_1, n_2}(\epsilon)$ of $H_2(\epsilon)$ for $\epsilon \in \mathbb{R}$, $|\epsilon|$ small, converging to $\lambda_{n_1, n_2}(0)$ as $\epsilon \rightarrow 0$. Moreover $\lambda_{n_1, n_2}(\epsilon)$ is real.*

Proof. The first statement follows from Proposition 1 (applicable after Proposition 2), the definition of stability of eigenvalues and the fact that each eigenvalue $\lambda_{n_1, n_2}(0)$ of $H_2(0)$ is simple. The reality of $\lambda_{n_1, n_2}(\epsilon)$ follows from its uniqueness, as anticipated in Remark 1. ■

3 The operator family $H_3(\epsilon)$: stability and reality of the perturbed eigenvalues

We first consider the minimal operator in $L^2(\mathbb{R})$ defined on $C_0^\infty(\mathbb{R})$ by the formal expression (5) for $|\epsilon| < 1$, i.e. $\forall u \in C_0^\infty(\mathbb{R})$ we set

$$H_3(\epsilon)u = -u'' + x^2(ix)^\epsilon u. \tag{10}$$

Then (10) can be rewritten as follows

$$H_3(\epsilon)u = -u'' + e^{(\text{sign } x)i\epsilon\frac{\pi}{2}}|x|^{2+\epsilon}u = -u'' + \cos\left(\frac{\pi}{2}\epsilon\right)|x|^{2+\epsilon}u + (\text{sign } x)i\sin\left(\frac{\pi}{2}\epsilon\right)|x|^{2+\epsilon}u,$$

where $\text{sign } x = \begin{cases} 1, & \text{if } x \geq 0 \\ -1, & \text{if } x < 0. \end{cases}$

Then $H_3(\epsilon)$ is closable and the domain of its closure (still denoted $H_3(\epsilon)$) is $D(H_3(\epsilon)) = H^2(\mathbb{R}) \cap D(|x|^{2+\epsilon})$, $\forall \epsilon: |\epsilon| < 1$. Moreover the closed operator $H_3(\epsilon)$ has compact resolvent and therefore discrete spectrum, and it is \mathcal{PT} symmetric if \mathcal{T} is, once again, the complex conjugation operator and \mathcal{P} is the parity operator defined by $(\mathcal{P}u)(x) = u(-x)$, $\forall u \in L^2(\mathbb{R})$.

The unperturbed operator $H_3(0) = p^2 + x^2$, where $p^2 = -\frac{d^2}{dx^2}$, corresponds to the one-dimensional harmonic oscillator and its eigenvalues $\lambda_n(0) = 2n + 1$, $n \in \mathbb{N}_0$, are simple. Let us now proceed in analogy with Section 2 and prove that the operator family $H_3(\epsilon)$ satisfies conditions (1)–(3).

Proposition 3. *Conditions (1)–(3) are satisfied by the operator family $H_3(\epsilon)$, $-1 < \epsilon < 1$.*

Proof. (1) $\forall u \in C_0^\infty(\mathbb{R})$ let $K = \text{supp}(u)$. Then

$$\|H_3(\epsilon)u - H_3(0)u\|^2 = \int_K (e^{(\text{sign } x)i\epsilon\frac{\pi}{2}}|x|^{2+\epsilon} - x^2)^2 |u(x)|^2 dx. \tag{11}$$

Since K is compact there exists a constant $c > 0$ such that the integrand expression in the right hand side of (11) can be bounded from above by $c|u(x)|^2$, $\forall x \in K$, $\forall \epsilon \in]-1, 1[$. Then, ■

by Lebesgue's dominated convergence theorem the r.h.s. of (11) converges to zero as $\epsilon \rightarrow 0$ and this proves (6). A similar argument proves (7).

(2) $\forall u \in C_0^\infty(\mathbb{R}^2)$ we have

$$\Re\langle u, H_3(\epsilon)u \rangle = \langle u, p^2u \rangle + \cos\left(\frac{\pi}{2}\epsilon\right) \langle u, |x|^{2+\epsilon}u \rangle \geq \langle u, p^2u \rangle.$$

Thus, as for the case of $H_2(\epsilon)$, (8) is satisfied with $\gamma, b = 0$ and $a = 1$.

(3) Again, in analogy with the argument used for $H_2(\epsilon)$, let $z \in \mathbb{C}$ and $u \in D(H_3(\epsilon))$ be such that $\|u\| = 1$ and $u(x) = 0$ for $|x| < n$, $n \in \mathbb{N}$. Then

$$|z - \langle u, H_3(\epsilon)u \rangle| \geq \Re\langle u, H_3(\epsilon)u \rangle - |z| \geq \cos\left(\frac{\pi}{2}\epsilon\right) \langle u, |x|^{2+\epsilon}u \rangle - |z| \geq \frac{n^2}{2} - |z|$$

for $|\epsilon| < \epsilon_1 := \frac{2}{3}$. ■

Now, with an argument analogous to that used to prove Corollary 1 we obtain the following result.

Corollary 2. *Near each eigenvalue $\lambda_n(0) = 2n + 1$, $n \in \mathbb{N}_0$, of $H_3(0)$ there exists one and only one eigenvalue $\lambda_n(\epsilon)$ of $H_3(\epsilon)$ for $\epsilon \in]-1, 1[$, $|\epsilon|$ small, converging to $\lambda_n(0)$ as $\epsilon \rightarrow 0$. Moreover $\lambda_n(\epsilon)$ is real.*

4 Conclusions

Concerning Section 1 we remark that 2×2 matrices can characterize complexified classical mechanical systems (see e.g. [22]) related to Schrödinger operators (4) for $r = s = 1$, i.e.

$$p_1^2 + p_2^2 + \omega_1^2 x_1^2 + \omega_2^2 x_2^2 + i\epsilon x_1 x_2.$$

Indeed the classical equation of motion can be written as

$$\begin{pmatrix} \ddot{x}_1 \\ \ddot{x}_2 \end{pmatrix} = -2 \begin{pmatrix} 2\omega_1^2 & i\epsilon \\ i\epsilon & 2\omega_2^2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad (12)$$

where \ddot{x}_k , $k = 1, 2$, denotes the second time derivative of x_k . It is straightforward to study the eigenvalue problem for the 2×2 matrix in (12), which corresponds to (2) with $\epsilon_k = 2\omega_k^2$, $k = 1, 2$. The eigenvalues are

$$\lambda_{\pm}(\epsilon) = (\omega_1^2 + \omega_2^2) \pm \sqrt{(\omega_1^2 - \omega_2^2)^2 - \epsilon^2}.$$

We see that the condition $|\epsilon| < |\omega_1^2 - \omega_2^2|$, which yields the reality of the spectrum in quantum mechanics, also yields the reality of the frequencies of the normal oscillation modes in classical mechanics. The quantum eigenvalues are given by

$$\lambda_{n_1, n_2}(\epsilon) = (2n_1 + 1)\lambda_+(\epsilon) + (2n_2 + 1)\lambda_-(\epsilon), \quad \forall n_1, n_2 \in \mathbb{N}_0.$$

Expanding the functions $\lambda_{\pm}(\epsilon)$ in powers of ϵ we obtain the Rayleigh–Schrödinger perturbation expansion (RSPE) for the eigenvalues $\lambda_{n_1, n_2}(\epsilon)$. All expansions clearly converge for $|\epsilon| < |\omega_1^2 - \omega_2^2|$, therefore the radius of convergence of the RSPE coincides with the threshold of transition between real and complex spectrum.

In a similar way one proves that the complexified classical Hamiltonian corresponding to the 2×2 matrix (3) for $e, b \in \mathbb{R}$

$$p_1^2 + p_2^2 + \frac{1}{2}(e + i\epsilon)x_1^2 + \frac{1}{2}(e - i\epsilon)x_2^2 + bx_1x_2$$

admits real normal modes with real frequencies if $|\epsilon| < |b|$.

As far as Section 2 is concerned we remark that a suitable generalization regards polynomial perturbations of the d -dimensional harmonic oscillator, $d > 2$. In addition, still in dimension $d = 2$ one could try to study the case of resonant frequencies, not with stability methods (which fail because of the degeneracy of the unperturbed eigenvalues), but proving the reality of the Rayleigh–Schrödinger perturbation expansion and its summability.

Finally, concerning Section 3 an open question is the rigorous proof of the existence of complex eigenvalues for $-1 < \epsilon < 0$, supporting the existing numerical results, and the analysis of the case $\epsilon < -1$.

Acknowledgement

We wish to thank R. Tateo for useful correspondence.

References

- [1] Caliceti E., Graffi S., Sjöstrand J., Spectra of \mathcal{PT} -symmetric operators and perturbation theory, *J. Phys. A: Math. Gen.* **38** (2005), 185–193, [math-ph/0407052](#).
- [2] Caliceti E., Graffi S., Sjöstrand J., \mathcal{PT} symmetric non-selfadjoint operators, diagonalizable and non-diagonalizable, with a real discrete spectrum, *J. Phys. A: Math. Theor.* **40** (2007), 10155–10170, [arXiv:0705.4218](#).
- [3] Caliceti E., Cannata F., Graffi S., Perturbation theory of \mathcal{PT} symmetric Hamiltonians, *J. Phys. A: Math. Gen.* **39** (2006), 10019–10027, [math-ph/0607039](#).
- [4] Rotter I., Real eigenvalues in non-Hermitian quantum physics, [arXiv:0909.1232](#).
- [5] Guo A., Salamo G.J., Duchesne D., Morandotti R., Volatier-Ravat M., Aimez V., Siviloglou G.A., Chistodoulides D.N., Observation of \mathcal{PT} -symmetry breaking in complex optical potentials, *Phys. Rev. Lett.* **103** (2009), 093902, 4 pages.
- [6] Seyranian A.P., Kirillov O.N., Mailybaev A.A., Coupling of eigenvalues of complex matrices at diabolic and exceptional points, *J. Phys. A: Math. Gen.* **38** (2005), 1732–1740, [math-ph/0411024](#).
- [7] Kirillov O.N., Mailybaev A.A., Seyranian A.P., Unfolding of eigenvalue surfaces near a diabolic point due to a complex perturbation, *J. Phys. A: Math. Gen.* **38** (2005), 5531–5546, [math-ph/0411006](#).
- [8] Albeverio S., Motovilov A.K., Shkalikov A.A., Bounds on variation of spectral subspaces under J -self-adjoint perturbations, *Integral Equations Operator Theory* **64** (2009), 455–486, [arXiv:0808.2783](#).
- [9] Albeverio S., Motovilov A.K., Tretter C., Bounds on the spectrum and reducing subspaces of a J -self-adjoint operator, [arXiv:0909.1211](#).
- [10] Caliceti E., Cannata F., Graffi S., An analytic family of \mathcal{PT} -symmetric Hamiltonians with real eigenvalues, *J. Phys. A: Math. Theor.* **41** (2008), 244008, 6 pages.
- [11] Nanayakkara A., Real eigenspectra in non-Hermitian multidimensional Hamiltonians, *Phys. Lett. A* **304** (2002), 67–72.
- [12] Nanayakkara A., Classical motion of complex 2-D non-Hermitian Hamiltonian systems, *Czechoslovak J. Phys.* **54** (2004), 101–107.
- [13] Bender C.M., Boettcher S., Real spectra in non-Hermitian Hamiltonians having \mathcal{PT} symmetry, *Phys. Rev. Lett.* **80** (1998), 5243–5246, [physics/9712001](#).
- [14] Dorey P., Dunning C., Tateo R., From \mathcal{PT} -symmetric quantum mechanics to conformal field theory, *Pramana J. Phys.* **73** (2009), 217–239, [arXiv:0906.1130](#).
- [15] Dorey P., Dunning C., Lishman A., Tateo R., \mathcal{PT} symmetry breaking and exceptional points for a class of inhomogeneous complex potentials, *J. Phys. A: Math. Theor.* **42** (2009), 465302, 41 pages. [arXiv:0907.3673](#).
- [16] Bender C.M., Besseghir K., Jones H.F., Yin X., Small- ϵ behavior of the non-Hermitian \mathcal{PT} -symmetric Hamiltonian $H = p^2 + x^2(ix)^\epsilon$, *J. Phys. A: Math. Theor.* **42** (2009), 355301, 10 pages, [arXiv:0906.1291](#).
- [17] Dorey P., Dunning C., Tateo R., Supersymmetry and the spontaneous breakdown of \mathcal{PT} symmetry, *J. Phys. A: Math. Gen.* **34** (2001), L391–L400, [hep-th/0104119](#).
- [18] Shin K.C., On the reality of the eigenvalues for a class of \mathcal{PT} -symmetric oscillators, *Comm. Math. Phys.* **229** (2002), 543–564, [math-ph/0201013](#).

- [19] Kato T., Perturbation theory for linear operators, 2nd ed., Springer-Verlag, Berlin – New York, 1976.
- [20] Reed M., Simon B., Methods of modern mathematical physics. IV. Analysis of operators, Academic Press, New York – London, 1978.
- [21] Vock E., Hunziker W., Stability of Schrödinger eigenvalue problems, *Comm. Math. Phys.* **83** (1982), 281–302.
- [22] Bender C.M., Holm D.D., Hook D.W., Complexified dynamical systems, *J. Phys. A: Math. Theor.* **40** (2007), F793–F804, [arXiv:0705.3893](https://arxiv.org/abs/0705.3893).