

## A NOTE ON INJECTIVE HULLS OF POSEMIGROUPS

CHANGCHUN XIA, SHENGWEI HAN, BIN ZHAO

ABSTRACT. In this note, we prove the existence of  $\mathcal{E}_{\leq}$ -injective hulls in the category  $\mathbf{PoSgr}_{\leq}$  of posemigroups and their submultiplicative order-preserving maps; here  $\mathcal{E}_{\leq}$  denotes the class of those morphisms  $h: A \rightarrow B$  for which  $h(a_1) \cdots h(a_n) \leq h(a)$  always implies  $a_1 \cdots a_n \leq a$ . The result of this note subsumes the results given by Lambek et al. (2012) and by Zhang and Laan (2014).

### 1. Introduction

The injective hull, though not under that name, was first obtained in [Baer (1940)]. The best-known early discussion of injective hulls is in [Eckmann and Schopf (1953)], who introduced them as maximal essential extensions. Banaschewski and Bruns [1967] characterized the injective hulls in the category of posets. In 1970, Bruns and Lakser characterized the injective hulls in the category of semilattices. As we all know, the injective objects in the category of pomonoids with pomonoid homomorphisms are trivial [Lambek et al. (2012)]. Hence, Lambek et al. changed a few of the definitions of pomonoid homomorphisms, chose submultiplicative order-preserving maps as morphisms of the category  $\mathbf{PoMon}_{\leq}$  of pomonoids to investigate the injectivity in  $\mathbf{PoMon}_{\leq}$ , where a *submultiplicative map* between pomonoids is a map  $f: (A, \cdot, \leq) \rightarrow (B, *, \leq)$  such that  $f(a) * f(a') \leq f(a \cdot a')$  for all  $a, a' \in A$ . Also, they proved that every pomonoid has an injective hull and gave the concrete form of the injective hulls. Based on the work of Lambek et al., dropping out the unit, Zhang and Laan intended to consider the injective hulls in the category  $\mathbf{PoSgr}_{\leq}$  of posemigroups, and constructed the injective hulls for a certain class of posemigroups with respect to a class of order embedding. However, for the question whether every posemigroup has an  $\mathcal{E}_{\leq}$ -injective hull in  $\mathbf{PoSgr}_{\leq}$ , Zhang and Laan did not give an answer. In this note, we shall give an affirmative answer and obtain the concrete form of injective hulls for general posemigroups. Particularly, when a posemigroup  $S$  is a pomonoid, the injective hull of  $S$  in  $\mathbf{PoSgr}_{\leq}$  we obtain is exactly the injective hull of  $S$  in  $\mathbf{PoMon}_{\leq}$  Lambek et al. gave.

A *partially ordered semigroup* (posemigroup for short) is a semigroup  $(S, \cdot)$  with a partial order  $\leq$  on  $S$  which is compatible with the partial order  $\leq$ , that is,  $x \leq y \Rightarrow x \cdot z \leq y \cdot z$  and  $z \cdot x \leq z \cdot y$  for any  $z$  in  $S$ . A *quantale* is a complete lattice  $Q$  with an associative binary operation  $\&$

---

We would like to thank the National Natural Science Foundation of China (11531009), the Natural Science Program for Basic Research of Shaanxi Province, China (2015JM1020) and the Fundamental Research Funds for the Central Universities (GK201501001).

Received by the editors 2017-01-19 and, in final form, 2017-02-01.

Transmitted by Walter Tholen. Published on 2017-02-08.

2010 Mathematics Subject Classification: 06F05, 06F07, 08B30.

Key words and phrases: partially ordered semigroup, quantale, quantic nucleus, injective hull.

© Changchun Xia, Shengwei Han, Bin Zhao, 2017. Permission to copy for private use granted.

satisfying  $a \& (\bigvee_{i \in I} b_i) = \bigvee_{i \in I} (a \& b_i)$  and  $(\bigvee_{i \in I} b_i) \& a = \bigvee_{i \in I} (b_i \& a)$  for any  $a \in Q$  and  $\{b_i\}_{i \in I} \subseteq Q$ . A *quantic nucleus*  $j$  on a quantale  $Q$  is a submultiplicative closure operator. It is easy to show that  $j(a \& b) = j(a \& j(b)) = j(j(a) \& b) = j(j(a) \& j(b))$  for all  $a, b \in Q$ . Furthermore, we can also prove that the set  $Q_j$  of fixed points of  $j$  is a quantale with  $a \otimes b = j(a \& b)$  and  $\bigvee_{i \in I} a_i = j(\bigvee_{i \in I} a_i)$ . Clearly, every quantale is necessarily a posemigroup. For a given semigroup  $(S, \cdot)$ , we denote by  $\mathcal{P}(S)$  the set of all subsets of  $S$ . Clearly,  $(\mathcal{P}(S), \bullet)$  forms a quantale under the inclusive order, where  $A \bullet B = \{a \cdot b : a \in A, b \in B\}$ . When  $A = \{a\}$ , we write  $A \bullet B$  and  $B \bullet A$  simply as  $a \cdot B$  and  $B \cdot a$ .

Let  $\mathcal{C}$  be a category and let  $\mathcal{M}$  be a class of morphisms in  $\mathcal{C}$ . Then an object  $S$  in  $\mathcal{C}$  is  *$\mathcal{M}$ -injective* in  $\mathcal{C}$  provided that for any morphism  $h: A \rightarrow B$  in  $\mathcal{M}$  and any morphism  $f: A \rightarrow S$  in  $\mathcal{C}$  there exists a morphism  $g: B \rightarrow S$  such that  $g \circ h = f$ . A morphism  $\eta: A \rightarrow B$  in  $\mathcal{M}$  is called  *$\mathcal{M}$ -essential* provided that every morphism  $\psi: B \rightarrow C$  in  $\mathcal{C}$  for which the composite  $\psi \circ \eta$  is in  $\mathcal{M}$  is itself in  $\mathcal{M}$ . An object  $H \in \mathcal{C}$  is said to be an  *$\mathcal{M}$ -injective hull* of an object of  $S$  provided that  $H$  is  $\mathcal{M}$ -injective and there exists an  $\mathcal{M}$ -essential morphism  $S \rightarrow H$ .

For notions and concepts concerned, but not explained, please refer to [Lambek et al. (2012)], [Zhang and Laan (2014)] and [Adámek et al. (2004)].

## 2. Main results

In this section, we shall show that every posemigroup has an  $\mathcal{E}_{\leq}$ -injective hull, and give its concrete form. Firstly, we review some main results on injective hulls in the category **PoMon** $_{\leq}$  and **PoSgr** $_{\leq}$ .

2.1. THEOREM. [Lambek et al. (2012)] *A pomonoid is injective (with respect to embeddings) if and only if it is a quantale.*

2.2. THEOREM. [Lambek et al. (2012)] *The map  $\eta(x \mapsto x \downarrow): A \rightarrow \mathcal{Q}(A)$  is an essential embedding and thus is an embedding into an injective hull.*

2.3. THEOREM. [Zhang and Laan (2014)] *Let  $(S, \cdot, \leq)$  be a posemigroup. Then  $S$  is  $\mathcal{E}_{\leq}$ -injective in **PoSgr** $_{\leq}$  if and only if  $S$  is a quantale.*

2.4. THEOREM. [Zhang and Laan (2014)] *Let  $S$  be a posemigroup such that  $cl(s \downarrow) = s \downarrow$  for every  $s \in S$ . Then  $\mathcal{D}(S)$  is an  $\mathcal{E}_{\leq}$ -injective hull in **PoSgr** $_{\leq}$ .*

In what follows we shall prove our main result.

Let  $(S, \cdot, \leq)$  be a posemigroup and  $a \in S$ . Then we denote by  $a \downarrow$  the set  $\{b \in S : b \leq a\}$ . For any subset  $X$  of  $S$ , we define its closure by  $X^*$

$$X^* \triangleq X^{\text{ul}} \cap X^L \cap X^R \cap X^T$$

where

$$X^{\text{ul}} \triangleq \{s \in S : \forall b \in S, X \subseteq b \downarrow \Rightarrow s \leq b\}$$

$$X^L \triangleq \{s \in S : \forall a, b \in S, X \cdot a \subseteq b \downarrow \Rightarrow s \cdot a \leq b\}$$

$$X^R \triangleq \{s \in S : \forall a, b \in S, a \cdot X \subseteq b \downarrow \Rightarrow a \cdot s \leq b\}$$

$$X^T \triangleq \{s \in S : \forall a, b, c \in S, a \cdot X \cdot c \subseteq b \downarrow \Rightarrow a \cdot s \cdot c \leq b\}$$

It is easy to verify that  $X^* = X^T$  when  $S$  is a pomonoid.

2.5. LEMMA. [Xia et al. (2016)] *Let  $(S, \cdot, \leq)$  be a posemigroup. Then  $(\cdot)^*$  is a quantic nucleus on the power-set quantale  $(\mathcal{P}(S), \bullet)$ , that is,  $(\mathbf{S}^*, \otimes)$  is a quantic quotient of  $(\mathcal{P}(S), \bullet)$ , where  $\mathbf{S}^* = \{A \in \mathcal{P}(S) : A^* = A\}$ ,  $X \otimes Y = (X \bullet Y)^*$ .*

In [Xia et al. (2016)], we prove that  $(\cdot)^*$  is also topological, that is,  $\{x\}^* = x \downarrow$  for all  $x \in S$ . We define now a map  $\eta_S : S \rightarrow \mathbf{S}^*$  as follows  $\eta_S(x) = \{x\}^*$  for all  $x \in S$ . In [Xia et al. (2016)], we prove that  $(\eta_S, \mathbf{S}^*)$  is the least quantale completion of  $S$ . In the following we shall see that  $\eta_S$  is in fact  $\mathcal{E}_{\leq}$ -essential in  $\mathbf{PoSgr}_{\leq}$ , that is,  $\mathbf{S}^*$  is an  $\mathcal{E}_{\leq}$ -injective hull of  $S$  in  $\mathbf{PoSgr}_{\leq}$ .

2.6. THEOREM. *Let  $(S, \cdot, \leq)$  be a posemigroup. Then  $\mathbf{S}^*$  is an  $\mathcal{E}_{\leq}$ -injective hull of  $S$  in  $\mathbf{PoSgr}_{\leq}$ .*

PROOF. One can easily see that  $(x \downarrow)^* = x \downarrow$  for any  $x \in S$ . In the following we shall prove that  $\eta_S$  is an  $\mathcal{E}_{\leq}$ -essential morphism in  $\mathbf{PoSgr}_{\leq}$ . Clearly,  $\eta_S$  belongs to  $\mathcal{E}_{\leq}$ . By Theorem 2.3,  $\mathbf{S}^*$  is  $\mathcal{E}_{\leq}$ -injective. It remains to show that  $\eta_S$  is  $\mathcal{E}_{\leq}$ -essential. Let  $\psi : \mathbf{S}^* \rightarrow (B, \leq, *)$  be a morphism in  $\mathbf{PoSgr}_{\leq}$  such that  $\psi \circ \eta_S \in \mathcal{E}_{\leq}$ . We shall verify that  $\psi \in \mathcal{E}_{\leq}$ . Suppose that  $\psi(X_1) * \psi(X_2) * \cdots * \psi(X_n) \leq \psi(Y)$  in  $B$ , where  $X_1, X_2, \dots, X_n, Y \in \mathbf{S}^*$ . We need to show that  $X_1 \otimes X_2 \otimes \cdots \otimes X_n \subseteq Y$ . In fact, we only need to prove that  $X_1 \bullet X_2 \bullet \cdots \bullet X_n \subseteq Y^{\text{ul}} \cap Y^L \cap Y^R \cap Y^T$ .

$$(1) X_1 \bullet X_2 \bullet \cdots \bullet X_n \subseteq Y^{\text{ul}}.$$

Assume that  $x_1 \in X_1, \dots, x_n \in X_n$ . We let  $b \in S$  and  $Y \subseteq b \downarrow$ . Then we have

$$\begin{aligned} (\psi \circ \eta_S)(x_1) * \cdots * (\psi \circ \eta_S)(x_n) &= \psi(x_1 \downarrow) * \cdots * \psi(x_n \downarrow) \\ &\leq \psi(X_1) * \cdots * \psi(X_n) \\ &\leq \psi(Y) \\ &\leq \psi(b \downarrow) \\ &= (\psi \circ \eta_S)(b). \end{aligned}$$

It follows from the case that  $\psi \circ \eta_S \in \mathcal{E}_{\leq}$  that  $x_1 \cdot x_2 \cdot \cdots \cdot x_n \leq b$ . Hence,  $X_1 \bullet X_2 \bullet \cdots \bullet X_n \subseteq b \downarrow$ , which implies  $X_1 \bullet X_2 \bullet \cdots \bullet X_n \subseteq Y^{\text{ul}}$ .

$$(2) X_1 \bullet X_2 \bullet \cdots \bullet X_n \subseteq Y^L.$$

Suppose that  $x_1 \in X_1, \dots, x_n \in X_n$ . We let  $a, b \in S$  and  $Y \cdot a \subseteq b \downarrow$ . Then we have

$$\begin{aligned} (\psi \circ \eta_S)(x_1) * \cdots * (\psi \circ \eta_S)(x_n) &= \psi(x_1 \downarrow) * \cdots * \psi(x_n \downarrow) * \psi(a \downarrow) \\ &\leq \psi(X_1) * \cdots * \psi(X_n) * \psi(a \downarrow) \\ &\leq \psi(Y) * \psi(a \downarrow) \\ &\leq \psi(Y \otimes a \downarrow) \\ &\leq \psi((Y \cdot a)^*) \\ &\leq \psi(b \downarrow) \\ &= (\psi \circ \eta_S)(b), \end{aligned}$$

It follows from the case that  $\psi \circ \eta_S \in \mathcal{E}_{\leq}$  that  $x_1 \cdot x_2 \cdot \dots \cdot x_n \cdot a \leq b$ . Hence,  $(X_1 \bullet X_2 \bullet \dots \bullet X_n) \cdot a \subseteq b \downarrow$ , which implies  $X_1 \bullet X_2 \bullet \dots \bullet X_n \subseteq Y^L$ .

(3) Similar to (2), we can prove that  $X_1 \bullet X_2 \bullet \dots \bullet X_n \subseteq Y^R$  and  $X_1 \bullet X_2 \bullet \dots \bullet X_n \subseteq Y^T$ .

By (1)-(3), we conclude that  $X_1 \bullet X_2 \bullet \dots \bullet X_n \subseteq Y^{\text{ul}} \cap Y^L \cap Y^R \cap Y^T$ , that is,  $X_1 \bullet X_2 \bullet \dots \bullet X_n \subseteq Y$ . Therefore,  $\mathbf{S}^*$  is an  $\mathcal{E}_{\leq}$ -injective hull of  $S$  in  $\mathbf{PoSgr}_{\leq}$ . ■

2.7. REMARK.  $\mathbf{S}^*$  is the injective hull of  $S$  in  $\mathbf{PoMon}_{\leq}$  whenever  $S$  is a pomonoid.

## References

- J. Adámek, H. Herrlich and G.E. Strecker (2004), *Abstract and Concrete Categories: The joy of cats*, Wiley, New York.
- B. Banaschewski and G. Bruns (1967). Categorical characterization of the MacNeille completion. *Archiv der Mathematik*. **18**, 369–377.
- R. Baer (1940). Abelian groups which are direct summands of every containing group. *Bull. Mmer. Math. Soc.* **46**, 800–806.
- G. Bruns, H. Lakser (1970), Injective hulls of semilattices. *Canad. Math. Bull.* **13**, 115–118.
- B. Eckmann and A. Schopf (1953), Über injektive Moduln. *Archiv der Mathematik* **4**, 75–78.
- J. Lambek, M. Barr, J.F. Kennison and R. Raphael (2012), Injective hulls of partially ordered monoids. *Theory Appl. Categ.* **26**, 338–348.
- C.C. Xia, B. Zhao and S.W. Han (2016), Some further results of quantale completions of ordered semigroup. Submitted to *Acta Mathematica Sinica, Chinese Series*.
- X. Zhang, V. Laan (2014), Injective hulls for posemigroups. *Proc. Est. Acad. Sci.* **63**, 372–378.

*Department of Mathematics, Shaanxi Normal University,  
Xi'an 710119, China*

Email: xiachangchun@snnu.edu.cn

hansw@snnu.edu.cn

zhaobin@snnu.edu.cn

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available from the journal's server at <http://www.tac.mta.ca/tac/>. It is archived electronically and in printed paper format.

**SUBSCRIPTION INFORMATION** Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to [tac@mta.ca](mailto:tac@mta.ca) including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca).

**INFORMATION FOR AUTHORS** The typesetting language of the journal is  $\text{\TeX}$ , and  $\text{\LaTeX}2\text{e}$  is required. Articles in PDF format may be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

**MANAGING EDITOR.** Robert Rosebrugh, Mount Allison University: [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca)

**$\text{\TeX}$ NICAL EDITOR.** Michael Barr, McGill University: [barr@math.mcgill.ca](mailto:barr@math.mcgill.ca)

**ASSISTANT  $\text{\TeX}$  EDITOR.** Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: [gavin\\_seal@fastmail.fm](mailto:gavin_seal@fastmail.fm)

**TRANSMITTING EDITORS.**

Clemens Berger, Université de Nice-Sophia Antipolis: [cberger@math.unice.fr](mailto:cberger@math.unice.fr)

Richard Blute, Université d' Ottawa: [rblute@uottawa.ca](mailto:rblute@uottawa.ca)

Lawrence Breen, Université de Paris 13: [breen@math.univ-paris13.fr](mailto:breen@math.univ-paris13.fr)

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva, Nuance Communications Inc: [valeria.depaiva@gmail.com](mailto:valeria.depaiva@gmail.com)

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: [kathryn.hess@epfl.ch](mailto:kathryn.hess@epfl.ch)

Martin Hyland, University of Cambridge: [M.Hyland@dpms.cam.ac.uk](mailto:M.Hyland@dpms.cam.ac.uk)

Anders Kock, University of Aarhus: [kock@imf.au.dk](mailto:kock@imf.au.dk)

Stephen Lack, Macquarie University: [steve.lack@mq.edu.au](mailto:steve.lack@mq.edu.au)

F. William Lawvere, State University of New York at Buffalo: [wlawvere@buffalo.edu](mailto:wlawvere@buffalo.edu)

Tom Leinster, University of Edinburgh: [Tom.Leinster@ed.ac.uk](mailto:Tom.Leinster@ed.ac.uk)

Ieke Moerdijk, Utrecht University: [i.moerdijk@uu.nl](mailto:i.moerdijk@uu.nl)

Susan Niefield, Union College: [niefiels@union.edu](mailto:niefiels@union.edu)

Robert Paré, Dalhousie University: [pare@mathstat.dal.ca](mailto:pare@mathstat.dal.ca)

Jiri Rosicky, Masaryk University: [rosicky@math.muni.cz](mailto:rosicky@math.muni.cz)

Giuseppe Rosolini, Università di Genova: [rosolini@disi.unige.it](mailto:rosolini@disi.unige.it)

Alex Simpson, University of Ljubljana: [Alex.Simpson@fmf.uni-lj.si](mailto:Alex.Simpson@fmf.uni-lj.si)

James Stasheff, University of North Carolina: [jds@math.upenn.edu](mailto:jds@math.upenn.edu)

Ross Street, Macquarie University: [ross.street@mq.edu.au](mailto:ross.street@mq.edu.au)

Walter Tholen, York University: [tholen@mathstat.yorku.ca](mailto:tholen@mathstat.yorku.ca)

Myles Tierney, Université du Québec à Montréal: [tierney.myles4@gmail.com](mailto:tierney.myles4@gmail.com)

R. J. Wood, Dalhousie University: [rjwood@mathstat.dal.ca](mailto:rjwood@mathstat.dal.ca)